

Transport and adsorption of antibiotics by marine sediments in a dynamic environment

Weihai Xu,^{1,2} Gan Zhang,³ Onyx W.H.Wai,¹ Shichun Zou,⁴ Xiangdong Li^{1*}

¹ Department of Civil and Structural Engineering, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong

² South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China

³ State Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

⁴ School of Oceanology, Sun Yat-sen University, Guangzhou 510250, China

Abstract

Background, aim, and scope Bed-sediments are the major sink for many contaminants in aquatic environments. With increasing knowledge of and research on the environmental occurrence of antibiotics, there has been growing interest in their behaviour and fate in aquatic environments. However, there is little information about the behaviour of antibiotics in a dynamic water/sediment environment, such as river and coastal marine water. Therefore, the aims of the present study were: (1) to study the transport and distribution of four common antibiotics between water and sediment in both dynamic and quiescent water/sediment systems; (2) to understand the persistence and possible degradation of the four antibiotics in the two different systems.

Materials and methods A Lid-driven Elongated Annular Flume (LEAF), designed to reduce the centrifugal effect, was used to simulate a dynamic water environment. In addition, a quiescent water/sediment experiment was conducted for comparison with the dynamic water system. The seawater and sediment, used in both experiments of flowing and quiescent water/sediment systems, were collected from Victoria Harbour, a dynamic coastal environment in an urban setting. The four antibiotics selected in this study were ofloxacin

* Corresponding authors (cexdli@polyu.edu.hk)

(OFL), roxithromycin (RTM), erythromycin (ETM), and sulfamethoxazole (SMZ), the most commonly used antibiotics in south China.

Results and discussion Antibiotics in an overlying solution decreased very quickly in the flume system due to the sorption to suspended particles and surface sediment. There were significant differences in the adsorption of the four antibiotics in sediment. OFL showed a high tendency to be adsorbed by sediment with a high K_d value (2980 L/Kg), while the low K_d values of SMZ indicated that there was a large quantity in water. The four antibiotics reached a depth of 20–30 mm in the sediment over a period of 60 days in the flume system. However, the compounds were only found in surface sediment (above 10 mm) in the quiescent system, indicating the influence of the dynamic flume system on the distribution of antibiotics in sediment. OFL showed a moderate persistence in the dynamic flume system, while other three antibiotics had less persistence in sediment. However, all of the four compounds showed moderate persistence in the quiescent system.

Recommendations and perspectives The study showed the rapid diffusive transfer of antibiotics from water to sediment in the dynamic flume system. The four antibiotics exhibited larger differences in their adsorption to sediment in both dynamic and quiescent systems due to their different K_d values. The high sorption of antibiotics to marine sediment may reduce their availability to benthic invertebrates.

Keywords: Antibiotics · transport · adsorption · persistence · sediments · dynamic water environment · South China.

1 Background, aim, and scope

The occurrence and potential adverse effects of pharmaceutical residuals in aquatic environments have generated growing interest in recent years due to their potential threat to the balance of the ecosystem and the risk they pose to the health of humans and animals. Antibiotics rank among the most important classes of pharmaceuticals because of the large amounts used in medicines for humans and animals, and in aquaculture.

One of the major pathways of antibiotics to the aquatic environment is via municipal sewage treatment plants (STPs). The removal of antibiotics by STPs has been shown to be incomplete (Miao et al. 2002; Xu et al. 2007b), and the effects on antibiotics and antibiotic-resistant bacteria during the wastewater treatment process is largely unknown. Various groups of antibiotics and some of their metabolites have been detected in the effluents from municipal STPs (Golet et al. 2003; Ternes et al. 2002). It is known that, as a result, considerable quantities of antibiotics enter surface water environments, such as river water and sediments (Golet et al. 2001; Hirsch et al. 1999; Lindberg et al. 2004; Sacher et al. 2001; Xu et al. 2007a), and coastal water and groundwater (Daughton and Ternes 1999; Heberer 2002; Ternes 1998).

With increasing knowledge of the environmental occurrence of antibiotics, interest is now being focused on their behaviour and fate in the environment (Brannon et al. 2005). For example, exposure to antibiotics might induce resistance (Kummerer 2004), and lead to the horizontal transfer of resistance genes in field bacterial populations (Dantas et al. 2008; Davison 1999; Pruden et al. 2006). Once introduced into surface waters, antibiotics may also undergo biodegradation and adsorption to sediment. Aquatic sediment is the most important sink of pharmaceuticals and other contaminants. The distribution of a particular compound between sediment/suspended particulate matter and water is largely dependent on the lipophilicity of the compound and on the sorption properties of the sediment. In order to investigate the distribution kinetics between water and sediment, and the environmental fate of contaminants, many test systems have been established under a variety of relevant environmental conditions (Freitag et al. 1985; Freitag et al. 1982; Suzuki et al. 1998). One

drawback of most of these systems is that they are based on the quiescent system. Hence, these systems cannot provide high comparability and reproducibility with dynamic water systems under real environmental conditions (Sabaliunas et al. 2003). Experimental flumes with sediment and an overlying solution have proven to be good at mimicking these dynamic riverine and coastal environments (Allan et al. 2004; Chan and Wai 2004; Wai 2003). So far, very little work has been conducted on the transport and distribution of antibiotics in dynamic water environments, such as the discharge points of wastewater effluents in riverbanks and coastal zones.

China has the largest population in the world, and antibiotics are in very common use, with the annual consumption being over 25,000 tonnes (Kummerer 2003; Xu et al. 2007a). The use of antibiotics in the fast-developing Pearl River Delta (PRD) region of south China is especially high (Richardson et al. 2005). The four antibiotics selected in the current study are representative of three classes of antibiotics that are commonly used in the PRD region. They have been detected in the Pearl River and other coastal waters in the PRD region at maximum concentrations of up to 1000 ng/L (Gulkowska et al. 2007; Xu et al. 2007a). The objectives of this study were: (1) to investigate the distribution of four common antibiotics between water and sediment in both dynamic and quiescent water/sediment systems; (2) to study the vertical distribution of antibiotics in 50-mm-bed sediment layers in the flume system, and to compare it with that in a quiescent system; (3) to understand the persistence and degradation of the four antibiotics in the two water systems.

2 Materials and methods

2.1 Collecting sediment and seawater

Seawater and sediment were collected from Hong Kong's Victoria Harbour in March 2006. The seawater was transported to the laboratory in 50-litre plastic containers. Sediment was obtained from a depth of < 10 cm in the harbour. Stones, branches, and other solid materials in the sediment were carefully removed, and the sediment was then thoroughly mixed before being introduced into the flume and quiescent systems. The characteristics of the seawater and sediment are shown in Table 1.

2.2 Methods

2.2.1 *Dynamic water/sediment system*

A Lid-driven Elongated Annular Flume (LEAF), designed to reduce the centrifugal effect, was used to perform the dynamic water experiment (Fig. 1). The flume has two identical 3-m long straight sections, which are meant to create a uniform flow environment. The inner side of the flume is made of glass in order to reduce the sorption of antibiotics. The depth of the water and vertical position of the lid can be adjusted. The lid is driven by an adjustable-speed motor. The water is re-circulated using the butterfly lid controlled by an electromotor. An ultrasonic velocity monitor was installed in the straight sections of the flume. The flume was housed in a large laboratory, with the temperature of the room kept at 20 ± 2 °C. About 60 mm of sediment were laid evenly at the bottom of the flume, and 400 litres of seawater were then slowly added to the flume. The flume was kept running for a week with a lid rotational speed (RS) of 0.6 m/s (the water velocity was about 20 cm/s) before antibiotics were added to the water. Then, 16 mg of each of the four selected antibiotics were dissolved and spiked into the flume system. Following this, the flume started to work at a water velocity

of 20 cm/s and the run ended after 60 days. At each sampling period, one litre of surface seawater (3 cm below the surface) and bottom seawater (3 cm above the sediment), as well as sediment in the middle of the straight sections of the flume, were collected. After each sampling programme, two litres of seawater were added to compensate for what had been lost due to water sampling. Ultra-pure water was also added to keep the volume of the water constant every day. At the end of the experiment (60 days later), after the overlying seawater had been cautiously removed, the sediment in the straight section was longitudinally sectioned at 5-mm intervals down to 20 mm below the sediment–water interface, then 10 mm down to 30 mm.

2.2.2 *Quiescent water/sediment system*

A quiescent water/sediment system was designed using a tank in order to compare the results with the environmental fate of antibiotics in the dynamic flume channel. About 60 mm of sediment were laid evenly at the bottom, before 40 litres of seawater were added to the tank. The antibiotics were added as they had been in the flume experiments. Surface seawater and sediment samples were collected at each sampling period. Appropriate amounts of seawater and ultra-pure water were added to keep constant the volume of the water in the system.

2.2.3 *Extraction and analysis of antibiotics*

Seawater samples: The extraction of antibiotics from the seawater samples was performed mainly using the method described by Xu et al. (2007a), based on solid phase extraction (SPE).

Sediment samples: Samples of approximately 5 g were accurately weighed (200 ng $^{13}\text{C}_3$ -caffeine being added as a surrogate) and then placed into a 50-ml polypropylene

centrifuge tube into which 10 ml of extraction buffer had been added. The extraction buffer consisted of a 2:1:1 mixture of methanol, 0.1 M of a citric acid buffer with the pH adjusted to 6.0, and a 10 mM Na₂EDTA buffer with the pH adjusted to 6.0. The tubes were vortex mixed for 1 min and were then placed into an ultrasonic bath for 15 min (water temperature <40°C). The tubes were then centrifuged (Eppendorf Centrifuge 5810 R) for 10 min at 3000×g. The supernatant was decanted into a 500 ml glass bottle and the sediment residue was extracted one more time. The supernatant was combined and diluted to approximately 500 ml with ultra-pure water. SAX-HLB SPE cartridges were set up in tandem, and pre-conditioned sequentially with 6.0 ml of methanol, 6.0 ml of ultra-pure water, and 6.0 ml of a 10 mM Na₂EDTA buffer (pH 6.0). The samples were then passed through the SPE cartridges and SPE columns at a flow rate of approximately 5 ml/min. After this, the SAX cartridges were removed and the HLB cartridges washed with ultra-pure water (10 ml) before being dried with a flow of nitrogen gas for 1 h. Each cartridge was then eluted with three 2-ml vol of methanol. The analytes were collected in 10 ml brown glass vials, concentrated under a flow of N₂ gas to about 20 µl, and then dissolved in 40% aqueous methanol to a final volume of 1.0 ml.

The four antibiotics were analysed using high-performance liquid chromatography-electrospray ionization tandem mass spectrometry. A quantitative analysis of each compound was performed using LC-ESI-MS/MS with the MRM mode, using the two highest characteristic precursor ion/product ion transitions. Together with the retention times, the characteristic ions were used to ensure correct peak assignment and peak purity. ¹³C₃-caffeine was added as a surrogate standard to all samples prior to the enrichment

of the control to avoid possible losses during the analytical procedure. These spiked antibiotics in seawater and sediment were recovered at mean percentages ranging from 68% to 87% and from 65% to 72%, respectively. The limit of quantification (LOQ) for each compound in seawater and sediment are from 1 to 10 ng/L and 10 to 50 ng/g, respectively.

3 Results and discussion

3.1 Hydrodynamic characteristics of the LEAF

To check the uniformity of the flow field in the LEAF, a Preston tube was used to measure bed shear stress in the straight section of the flume. The stress was calculated according to the equations described in a previous study ([Patel 1965](#)). It was found that the lid rotational speed (RS) had a quadratic relationship with the flow velocity and the bottom shear stress. Fig. 2 shows the relationship of RS with the flow velocity and shear stress. The maximum bed shear stress that the LEAF could generate was around 1 N/m². Therefore, rather broad energy ranges (induced by shear stress from 0 to 1 N/m²) can be obtained through the LEAF, with these energy levels considered to be typical of near bottom shear stresses induced by a tide or flowing river water ([Bokuniewicz et al. 1991](#)). In this study, the RS was set up at 0.6 m/s. Thus, the induced bed shear stress and water velocity were about 0.1 N/m² and 20 cm/s, respectively. According to Chan and Wai ([2004](#)), the energy level induced in our experimental conditions (0.1 N/m²) was below the critical shear stress of typical non-cohesive sediment (0.15 N/m²). Wai ([2003](#)) showed that the concentration of sediment (turbidity signal) increased and decreased in response to changes in the flow field in the LEAF. This indicated that the flume that was used was indeed suitable for the study of the erosion and deposition

activities of sediment near the sediment-water interface. In addition, the LEAF was also suitable for carrying out long-term chemical-sediment sorption experiments because it had a well-controlled environment, and was made of non-reactive materials.

3.2 Sediment adsorption of antibiotics in both dynamic and quiescent environments

The changes over time in the concentration of antibiotics in the overlying seawater, including the surface and bottom seawater and sediment, in the dynamic flume system are shown in Fig. 3. The original spiked concentrations of the four antibiotics in the flume were 40 µg/L. However, the concentrations in seawater of all four antibiotics, detected at the first sampling event (30 min), were much lower than the initial spiked concentrations because of the rapid sorption to suspended particles and sediment. The antibiotic concentrations in the original seawater were mainly close to or lower than LOQ, and those in sediment were all below LOQ (see Table 1). Hence, the concentrations of the original antibiotics in water and sediment were much lower than the spiked concentrations, and need not be a cause of concern.

Concentration profiles in the overlying water suggested that the diffusive transfer of antibiotics into sediment was a quick process, with the compounds generally detected in surface sediment at a maximum concentration of more than 3000 ng/g at a very short sampling interval. Since the antibiotics were spiked into the seawater, their degradation, especially photodegradation, in the water phase was a competitive process between the sorption to sediment and a chemical transformation. It is often difficult to distinguish between sorption and degradation in a natural environment. However, the photodegradation function can be evaluated in this system based on previous studies under certain controlled conditions. The half-life times of OFL and AMX (amoxicillin) in a solution of water were 2.4 and 10.6 d,

respectively in a solar experiment (Andreozzi et al. 2003), and 7.0-17 d for ETM in sludge (Wu et al. 2008). Generally, the intensity of sunlight is about 50 times greater than the light emitted by common fluorescent lamps. In addition, the relatively low temperature in the laboratory may affect the reactivity of antibiotics with radicals formed by photons (Yamamoto et al. 2009). Together with the shielding from the apparatus, the photodegradation rates of the antibiotics in this experiment would certainly be greatly reduced. In a river environment biodegradation would occur at a less rapid rate than would be the case in photodegradation (Kummerer 2001; Yamamoto et al. 2009). Therefore, the quick changes in the concentrations of antibiotics in the overlying water at the initial sampling times (i.e., the first 10 days) were mainly due to sediment adsorption.

The concentrations in sediment reached several hundred ng/g at the first sampling event. This suggests that a water velocity of 20 cm/s can mobilize the small particles in the surface sediment so that the antibiotic compounds can be adsorbed rapidly to suspended particles, and then to surface sediment. It can be seen from Table 1 that the small particles (clay content) in suspended matter increased from 28.4% to 34.7% in comparison with the level in bulk sediment. On the contrary, the sand content decreased from 12.4% to 1.1%. The adsorption capacity of suspended particles is strongly related to their size, with larger particles providing additional surfaces for sorption (Clymo et al. 2005; Pouliquen and LeBris 1996; Thiele-Bruhn et al. 2004). An X-ray diffraction analysis of clay showed that the sorption of antibiotics can widen the interlayer spacing of clay. Hence, an increase in the content of clay can lead to an increase in adsorption capacity (Pouliquen and LeBris 1996). This rapid and extensive sorption of antibiotics into sediment, which had previously been reported for marine

sediments ([Cannavan et al. 2000](#); [Loffler et al. 2005](#)), is mainly attributable to the lipophilicity of these compounds. The concentrations of antibiotics in the water of the quiescent environment at the first sampling event were much higher than those in the flume (see Fig. 4). Correspondingly, the concentrations of antibiotics in sediment were lower than those in the flume system at the same sampling event due to the lack of dynamic interaction between water and sediment.

In both the flume and quiescent systems, large discrepancies were seen in the adsorption to sediment of the four antibiotics. The highest concentrations of 3730 ng/g and 1880 ng/g of OFL in water were detected in the flume and quiescent systems, respectively. However, the concentrations of SMZ were only 1036 ng/g and 629 ng/g, respectively. The adsorption of OFL into sediment was particularly strong, while the adsorption of SMZ was found to be weak. Similar findings were found in previous studies ([Drillia et al. 2005](#); [Sukul et al. 2008](#)).

3.3 Vertical profiles of antibiotics in sediments

In the flume system, the concentrations of the four selected antibiotics were found to be highest at the top layer of sediment, and to decrease sharply with depth (Fig. 5). At the end of the experiment, OFL persisted with a residual concentration of 542 ng/g in the top layer (0-5 mm), while the respective figures were 434 ng/g for RTM, 393 ng/g for ETM, and 55 ng/g for SMZ. The results revealed a pattern of diffusive distribution of the selected antibiotics into the sediment, due to the dynamic interaction between water and sediments in the flume system. A different pattern was seen in the quiescent system. Except for approximately 20 ng/g of OFL, no antibiotics were detected below 10 mm in the sediment profile. Allan et al. investigated the diffusion of the synthetic pyrethroid permethrin into sediment using flume

channels (Allan et al. 2005). Their results clearly showed that a large quantity of permethrin accumulated in the top layer (0-3 mm). Little permethrin was found in the sediment at a depth of 3 mm below the sediment-water interface. In our work, a high water velocity certainly affected the normal diffuse boundary layer, which may have resulted in mass transfers and increased the overall fluxes of antibiotics into the sediment. On the whole, the concentrations of antibiotics decreased by about one hundred ng/g for every 5 mm in the sediment profile to a depth of 25 mm, with the exception of the SMZ. No SMZ was found in the sediment at a depth below 15 mm, probably due to its low distribution coefficient and fast degradation rate (Holtge and Kreuzig 2007; Wu et al. 2009).

3.4 Effects of adsorption/desorption between seawater and sediment

The functions of the adsorption/desorption of antibiotics between water and sediment are rather complex. The distribution coefficient (K_d) and the normalized distribution coefficient (K_{oc}) with respect to the organic content (OC) (%) of the solid matrix have often been used to describe the effects of adsorption between water and sediment. K_d and K_{oc} were calculated according to:

$$K_d = C_s / C_{aq}$$

$$K_{oc} = 100 K_d / OC,$$

where C_s is the antibiotics equilibrium concentration in a solid matrix, and C_{aq} is the equilibrium concentration in an aqueous solution. In the present study, the K_d and K_{oc} values were determined using the concentrations of antibiotics in water and solid after 1 d or later, and should therefore be close to equilibrium conditions. That no degradation took place before 1 d was also taken into account. It is known that pharmaceuticals display a wide range

276 of mobility ($0.2 < K_d < 6000$ L/Kg), and that the variations in K_d for a given compound in
 277 different soils and sediments can be significant (Tolls 2001). In the present study, the K_d and
 278 K_{oc} values of each antibiotic did indeed vary significantly (Table 2). The values of K_d and
 279 K_{oc} decreased in the following order: OFL>RTM>ETM>SMZ. The adsorption of all
 280 compounds was generally higher in sediments with a higher total organic content (TOC)
 281 (Drillia et al. 2005). Hence, it is believed that the high sorption capacity (high K_d) to marine
 282 sediment may reduce the availability of antibiotics to benthic invertebrates. The calculated
 283 values showed that OFL has a high tendency to be adsorbed by sediment or solid particles,
 284 while SMZ, with a pK_a value of 1.69, has a low affinity for sediment. Studies have shown that
 285 the interaction of antibiotics with Ca^{2+} at clay surfaces is the prevalent sorption mechanism at
 286 low pH levels (Nowara et al. 1997). However, in neutral and weak alkaline pH conditions,
 287 this mechanism cannot play a leading role in the interaction of antibiotics with solids. This
 288 indicates that the interaction of deprotonated carboxylic acid with a clay surface can
 289 contribute significantly to the sorption of fluoroquinolones antibiotics (Nowara et al. 1997). An
 290 X-ray diffraction analysis of clay showed that the sorption of antibiotics can widen the clay
 291 interlayer spacing. Hence, due to the high clay content of the marine sediment in this study,
 292 the mechanism for the fluoroquinolones may be through cation bridging in the diffuse double
 293 layer at the surfaces of the clay. The possible sorption mechanism for the fluoroquinolones
 294 agrees with the high K_d obtained in previous studies (Tolls 2001). It has been suggested that
 295 electrochemical affinity and hydrophobic interaction can play important roles in the sorption
 296 of macrolides and sulfonamides to sediment/soil (Liu et al. 2002; Pan et al. 2009; Yamamoto
 297 et al. 2009).

The fate and mobility of six pharmaceuticals, including ofloxacin and sulfamethoxazole, were investigated in two types of soils with different values of TOC ([Drillia et al. 2005](#)). Ofloxacin had the highest K_d among the six pharmaceuticals. The values of K_d decreased in the following order: ofloxacin > propranolol > diclofenac > carbamazepine > sulfamethoxazole > clofibric acid. The details of the K_d and K_{oc} of the ofloxacin (OFL) and the sulfamethoxazole (SMZ) are also given in Table 2. It should be noted that the K_d and K_{oc} values in the present study were obtained under dynamic flume conditions similar to those of a subtropical river or coastal environment. Therefore, the K_d and K_{oc} values may be different from those obtained in a steady water/sediment system.

3.5 The persistence and fate of antibiotics in dynamic and quiescent environments

Antibiotics are designed to have a biological effect, and can persist in the human or animal body after administration. For easy absorption, most antibiotics are made to be water-soluble. These chemicals can degrade in the body more easily than in the environment. The persistence of antibiotics in the aquatic environment is a rather complex process, governed by biodegradation, sunlight photolysis, and other abiotic transformations, such as hydrolysis. Many antibiotics are relatively resistant to degradation under environmental conditions and pass through the STP treatment process ([Putschew et al. 2001](#); [Ternes 1998](#); [Ternes and Hirsch 2000](#)). At the end of the LEAF experiment (after 60 d), the final average concentrations of the four antibiotics in the surface water and sediment ranged from 0.26-1.27 $\mu\text{g/L}$ and 36-461 ng/g , respectively.

The degradation rate, often expressed as DT_{50} and DT_{90} (the time at which 50% and 90% of the parent compound has disappeared from sediment or water by transformation or

degradation, respectively), has been used to characterize the degradation of pharmaceuticals. Table 3 shows the degradation rate (DT_{50} and DT_{90}) of the four antibiotics in the human body, in the flume system, and in the quiescent system. The DT_{50} values generally varied from several hours to a day in the human body. However, in the flume environment, the maximum DT_{50} values in seawater and sediment exceeded 10 days. Thus, the transformation of the selected antibiotics in the human body is very different from that in the environment. Hence, appropriate experimental studies and field observations are indispensable for obtaining reliable data to assess the environmental fate of antibiotics. Great differences in the DT_{50} values of OFL and SMZ were found in the water and sediment samples due to the different degradation rates and partitioning process between water and sediment. The expectation is that OFL is adsorbed relatively quickly by solid matrices in the environment. As for SMZ, it is likely that a large amount stays in water.

The DT_{50} values, together with the DT_{90} values, are often used to show the persistence of antibiotics in the environment because the single DT_{50} value cannot exactly describe the rate of degradation. With the exception of SMZ, the DT_{90} values of the other three compounds (> 60 d) in sediment were longer than the values in water. Many antibiotic compounds photodegrade in liquids ([Halling-Sorensen et al. 2003](#)). In addition, photodegradation in sediment can only occur at the surface interface and in the first millimeters of depth. The chemical removal of antibiotics from sediments is done mainly through scouring or diffusion processes across the sediment-water interface. The persistence of antibiotics in sediment has become an important concern in the context of their long-term accumulation in aquatic environments ([Williams et al. 1999](#)).

According to the method described by Hollis (Hollis 1991), with regard to their persistence in sediment, antibiotics can be grouped into the following four classes: impersistent - $DT_{50} < 5$ d; slightly persistent - DT_{50} 5-21 d; moderately persistent - DT_{50} 22-60 d; and very persistent - $DT_{50} > 60$ d. By this classification, OFL was moderately persistent and the other three compounds were impersistent. However, in the quiescent system, the DT_{50} values ranged from 12.9 to 29 d, and from 24.3 to 41.1 d in seawater and sediment, respectively. The DT_{90} values were all >60 d for both seawater and sediment. Therefore, all four antibiotics displayed moderately persistent behaviour in the quiescent system.

4 Conclusions

The dynamic environment that we simulated gave some insight into the environmental behaviours of antibiotic compounds when they are introduced into aquatic environments. The results showed that the diffusive transfer of antibiotic into sediment was a quick process in the flume system. The four antibiotics exhibited larger differences in their adsorption to sediment in both dynamic and quiescent systems due to their different K_d values. With a high K_d value, OFL showed a high tendency to be adsorbed by sediment, while the low K_d value of SMZ indicated that a large quantity would remain in water. The experiments revealed that their high sorption capacity (high K_d) to marine sediment may reduce the availability of antibiotics to benthic invertebrates. In the flume system, the four antibiotics reached sediment layers of 20–30 mm over a period of 60 days. However, in the quiescent system the compounds were only found in surface sediment (above 10 mm). In the quiescent water system, the four compounds displayed moderate persistence, with DT_{50} values ranging from 24.3 to 41.1 d.,

and DT₉₀ values of ≥ 60 d for most of the compounds. In the dynamic flume system, OFL displayed a moderate persistence, with DT₅₀ values of ≥ 22 d in sediment, while the other three antibiotics displayed impersistence. Furthermore, the experiment indicated that antibiotics can resist degradation, with low concentrations persisting in sediment.

Acknowledgements: This work was funded by the Natural Science Foundation of China (No. 40672212) and The Hong Kong Polytechnic University (G-U300). The research for this work was also supported by the CAS/SAFEA International Partnership Programme for Creative Research Teams (KZCX2-YW-T001), the Research Grants Council of Hong Kong (PolyU5152/03E), the Area of Excellence (AoE) project under Grant No. AoE/P-04/2004 from the University Grants Council of Hong Kong, and the China Postdoctoral Science Foundation (No. 20070420149).

References

- Allan IJ, House WA, Parker A, Carter JE (2004) Transport and distribution of lindane and simazine in a riverine environment: measurements in bed sediments and modelling. *Pest Manage Sci* 60:417-433
- Allan IJ, House WA, Parker A, Carter JE (2005) Diffusion of the synthetic pyrethroid permethrin into bed-sediments. *Environ Sci Technol* 39:523-530
- Andreozzi R, Raffaele M, Nicklas P (2003) Pharmaceuticals in STP effluents and their solar photodegradation in aquatic environment. *Chemosphere* 50:1319-1330
- Bokuniewicz H, McTiernan L, Davis W (1991) Measurement of Sediment Resuspension Rates in Long-Island Sound. *Geo Mar Lett* 11:159-161
- Brannon JM, Price CB, Yost SL, Hayes C, Porter B (2005) Comparison of environmental fate and transport process descriptors of explosives in saline and freshwater systems. *Mar Pollut Bull* 50:247-251
- Cannavan A, Coyne R, Kennedy DG, Smith P (2000) Concentration of 22,23-dihydroavermectin B-1a detected in the sediments at an Atlantic salmon farm using orally administered ivermectin to control sea-lice infestation. *Aquaculture* 182:229-240
- Chan WY, Wai OWH (2004): Hydrodynamic and sediment transport properties in a Lid-driven Elongated Annular Flume (LEAF), Proceedings of the 4th International Symposium on Environmental Hydraulics & 14th Congress of IAHR-APD, Hong Kong, pp. 2195-2200

395 Chan WY, Wai OWH, Li YS. 2006. Critical shear stress for deposition of cohesive sediments in Mai Po.
396 Proceedings of the Conference of Global Chinese Scholars on Hydrodynamics:300-305.

397 Clymo AS, Shin JY, Holmen BA (2005) Herbicide sorption to fine particulate matter suspended
398 downwind of agricultural operations: Field and laboratory investigations. *Environ Sci Technol*
399 39:421-430

400 Dantas G, Sommer MOA, Oluwasegun RD, Church GM (2008) Bacteria subsisting on antibiotics.
401 *Science* 320:100-103

402 Daughton CG, Ternes TA (1999) Pharmaceuticals and personal care products in the environment:
403 Agents of subtle change? *Environ Health Perspect* 107:907-938

404 Davison J (1999) Genetic exchange between bacteria in the environment. *Plasmid* 42:73-91

405 Drillia P, Stamatelatou K, Lyberatos G (2005) Fate and mobility of pharmaceuticals in solid matrices.
406 *Chemosphere* 60:1034-1044

407 Freitag D, Geyer H, Kraus A, Viswanathan R, Kotzias D, Attar A, Klein W, Korte F (1982)
408 Ecotoxicological Profile Analysis .7. Screening Chemicals for Their Environmental Behavior
409 by Comparative-Evaluation. *Ecotoxicol Environ Saf* 6:60-81

410 Freitag D, Ballhorn L, Geyer H, Korte F (1985) Environmental-Hazard Profile of Organic-Chemicals -
411 an Experimental-Method for the Assessment of the Behavior of Organic-Chemicals in the
412 Ecosphere by Means of Simple Laboratory Tests with C-14-Labeled Chemicals. *Chemosphere*
413 14:1589-1616

414 Gobel A, Thomsen A, McArdell CS, Joss A, Giger W (2005) Occurrence and Sorption Behavior of
415 Sulfonamides, Macrolides, and Trimethoprim in Activated Sludge Treatment. *Environ Sci*
416 *Technol* 39:3981-3989

417 Golet EM, Alder AC, Hartmann A, Ternes TA, Giger W (2001) Trace Determination of
418 Fluoroquinolone Antibacterial Agents in Urban Wastewater by Solid-Phase Extraction and
419 Liquid Chromatography with Fluorescence Detection. *Anal Chem* 73:3632-3638

420 Golet EM, Xifra I, Siegrist H, Alder AC, Giger W (2003) Environmental Exposure Assessment of
421 Fluoroquinolone Antibacterial Agents from Sewage to Soil. *Environ Sci Technol*
422 37:3243-3249

423 Gulkowska A, He YH, So MK, Yeung LWY, Leung HW, Giesy JP, Lam PKS, Martin M, Richardson BJ
424 (2007) The occurrence of selected antibiotics in Hong Kong coastal waters. *Mar Pollut Bull*
425 54:1287-1293

426 Halling-Sorensen B, Sengelov G, Ingerslev F, Jensen LB (2003) Reduced antimicrobial potencies of
427 oxytetracycline, tylosin, sulfadiazin, streptomycin, ciprofloxacin, and olaquinox due to
428 environmental processes. *Arch Environ Contam Toxicol* 44:7-16

429 Heberer T (2002) Occurrence, fate, and removal of pharmaceutical residues in the aquatic environment:
430 a review of recent research data. *Toxicol Lett* 131:5-17

431 Hirsch R, Ternes T, Haberer K, Kratz K-L (1999) Occurrence of antibiotics in the aquatic environment.
432 *Sci Total Environ* 225:109-118

433 Hollis JM (1991) Mapping the Vulnerability of Aquifers and Surface Waters to Pesticide
434 Contamination at the National Regional Scale. *Pestic Soils Water: Current Perspectives*
435 47:165-174

436 Holtge S, Kreuzig R (2007) Laboratory testing of sulfamethoxazole and its metabolite
437 acetyl-sulfamethoxazole in soil. *Clean-Soil Air Water* 35:104-110

438 Jones OAH, Voulvoulis N, Lester JN (2002) Aquatic environmental assessment of the top 25 English

439 prescription pharmaceuticals. *Water Res* 36:5013-5022
 440 Kummerer K (2001) Emission and biodegradability of pharmaceuticals, contrast media, disinfectants
 441 and AOX from hospitals. *Pharmaceuticals in the Environment - Sources, Fate, Effects, and*
 442 *Risks*:29-41
 443 Kummerer K (2003) Significance of antibiotics in the environment. *J Antimicrob Chemother* 52:5-7
 444 Kummerer K (2004) Resistance in the environment. *J Antimicrob Chemother* 54:311-320
 445 Lindberg R, Jarnheimer P-A, Olsen B, Johansson M, Tysklind M (2004) Determination of antibiotic
 446 substances in hospital sewage water using solid phase extraction and liquid
 447 chromatography/mass spectrometry and group analogue internal standards. *Chemosphere*
 448 57:1479-1488
 449 Liu GD, Yu HF, Yan HS, Shi ZQ, He BL (2002) Utilization of synergetic effect of weak interactions in
 450 the design of polymeric sorbents with high sorption selectivity. *J Chromatogr A* 952:71-78
 451 Löffler D, Rombke J, Meller M, Ternes TA (2005) Environmental fate of pharmaceuticals in
 452 water/sediment systems. *Environ Sci Technol* 39:5209-5218
 453 Miao XS, Koenig BG, Metcalfe CD (2002) Analysis of acidic drugs in the effluents of sewage
 454 treatment plants using liquid chromatography-electrospray ionization tandem mass
 455 spectrometry. *J Chromatogr A* 952:139-147
 456 Nowara A, Burhenne J, Spiteller M (1997) Binding of fluoroquinolone carboxylic acid derivatives to
 457 clay minerals. *J Agric Food Chem* 45:1459-1463
 458 Pan B, Ning P, Xing BS (2009) Part V-sorption of pharmaceuticals and personal care products. *Environ*
 459 *Sci Pollut Res* 16:106-116
 460 Patel VC (1965) Calibration of Preston Tube and Limitations on Its Use in Pressure Gradients. *J Fluid*
 461 *Mech* 23:185-208
 462 Pouliquen H, LeBris H (1996) Sorption of oxolinic acid and oxytetracycline to marine sediments.
 463 *Chemosphere* 33:801-815
 464 Pruden A, Pei R, Storteboom H, Carlson KH (2006) Antibiotic Resistance Genes as Emerging
 465 Contaminants: Studies in Northern Colorado. *Environ Sci Technol* 40:7445-7450
 466 Putschew A, Schittko S, Jekel M (2001) Quantification of triiodinated benzene derivatives and X-ray
 467 contrast media in water samples by liquid chromatography-electrospray tandem mass
 468 spectrometry. *J Chromatogr A* 930:127-134
 469 Richardson BJ, Lam PKS, Martin M (2005) Emerging chemicals of concern: Pharmaceuticals and
 470 personal care products (PPCPs) in Asia, with particular reference to Southern China. *Mar*
 471 *Pollut Bull* 50:913-920
 472 Sabaliunas D, Webb SF, Hauk A, Jacob M, Eckhoff WS (2003) Environmental fate of Triclosan in the
 473 River Aire Basin, UK. *Water Res* 37:3145-3154
 474 Sacher F, Lange FT, Brauch H-J, Blankenhorn I (2001) Pharmaceuticals in groundwaters: Analytical
 475 methods and results of a monitoring program in Baden-Württemberg, Germany. *J Chromatogr*
 476 *A* 938:199-210
 477 Sukul P, Lamshoft M, Zuhlke S, Spiteller M (2008) Sorption and desorption of sulfadiazine in soil and
 478 soil-manure systems. *Chemosphere* 73:1344-1350
 479 Suzuki N, Yasuda M, Sakurai T, Nakanishi J (1998) Model simulation of environmental profile
 480 transformation and fate of polychlorinated dibenzo-p-dioxins and polychlorinated
 481 dibenzofurans by the multimedia environmental fate model. *Chemosphere* 37:2239-2250
 482 Ternes TA (1998) Occurrence of drugs in German sewage treatment plants and rivers. *Water Res*

483 32:3245-3260
 484 Ternes TA, Hirsch R (2000) Occurrence and behavior of X-ray contrast media in sewage facilities and
 485 the aquatic environment. *Environ Sci Technol* 34:2741-2748
 486 Ternes TA, Meisenheimer M, McDowell D, Sacher F, Brauch HJ, Haist-Gulde B, Preuss G, Wilme U,
 487 Zulei-Seibert N (2002) Removal of Pharmaceuticals during Drinking Water Treatment.
 488 *Environ Sci Technol* 36:3855-3863
 489 Thiele-Bruhn S, Seibicke T, Schulten HR, Leinweber P (2004) Sorption of sulfonamide pharmaceutical
 490 antibiotics on whole soils and particle-size fractions. *J Environ Qual* 33:1331-1342
 491 Tolls J (2001) Sorption of veterinary pharmaceuticals in soils: A review. *Environ Sci Technol*
 492 35:3397-3406
 493 Wai OWH (2003) A Lid-Driven Elongated Annular Flume (LEAF) for the determination of sediment
 494 transport properties. *Sedimentation and Sediment Transport, Proceedings*:241-244
 495 Williams RJ, Jurgens MD, Johnson AC (1999) Initial predictions of the concentrations and distribution
 496 of 17 beta-oestradiol, oestrone and ethinyl oestradiol in 3 English rivers. *Water Res*
 497 33:1663-1671
 498 Wu CX, Spongberg AL, Witter JD (2008) Determination of the persistence of pharmaceuticals in
 499 biosolids using liquid-chromatography tandem mass spectrometry. *Chemosphere* 73:511-518
 500 Wu CX, Spongberg AL, Witter JD (2009) Sorption and biodegradation of selected antibiotics in
 501 biosolids. *J Environ Sci Health Part A-Toxic/Hazard Subst Environ Eng* 44:454-461
 502 Xu WH, Zhang G, Zou SC, Li XD, Liu YC (2007a) Determination of selected antibiotics in the Victoria
 503 Harbour and the Pearl River, South China using high-performance liquid
 504 chromatography-electrospray ionization tandem mass spectrometry. *Environ Pollut*
 505 145:672-679
 506 Xu WH, Zhang G, Li XD, Zou SC, Li P, Hu ZH, Li J (2007b) Occurrence and elimination of antibiotics
 507 at four sewage treatment plants in the Pearl River Delta (PRD), South China. *Water Res*
 508 41:4526-4534
 509 Yamamoto H, Nakamura Y, Moriguchi S, Nakamura Y, Honda Y, Tamura I, Hirata Y, Hayashi A,
 510 Sekizawa J (2009) Persistence and partitioning of eight selected pharmaceuticals in the aquatic
 511 environment: Laboratory photolysis, biodegradation, and sorption experiments. *Water Res*
 512 43:351-362
 513
 514
 515

List of tables and figure captions

Table 1 Information about bulk seawater and sediment

Table 2 The K_d and K_{oc} values of four selected antibiotics

Table 3 The DT_{50} and DT_{90} values of the four antibiotics in seawater and sediment

Fig. 1 The setting of the experimental flume (LEAF)

Fig. 2 Relationship of the lid rotational speed (RS) with the averaged flow velocity ($\langle u \rangle$) and the bed shear stress (τ_b) (adopted from Chan et al. 2006)

Fig. 3 Temporal changes of the four antibiotics in surface water (a), bottom water (b), and sediment (c) of the flume system

Fig. 4 Temporal changes of the four antibiotics in surface water (a) and sediment (b) of the quiescent system

Fig. 5 Concentration profiles of the four antibiotics in sediment layers of the flume system

Table 1 Information about bulk seawater and sediment

Type	Organic content	Grain size (%)			Concentration of antibiotics (ng/L)			
		Sand	Silt	Clay	OFL	RTM	ETM	SMZ
Bulk Seawater	2.62 µg/ml	Not available			10	6	< LOQ ^b	<LOQ
Bulk Sediment	0.88 %	12.36	59.22	28.42	<LOQ	<LOQ	<LOQ	<LOQ
Suspended particulate matter ^a	1.13 %	1.14	64.21	34.65	Not available			

^a Suspended particulate matter was collected during the running of the flume for a week, before being spiked with antibiotics.

^b The LOQs for OFL, RTM, ETM, and SMZ were 10, 5, 5, and 1 ng/L, and 50, 20, 20, and 10 ng/g in seawater and marine sediment, respectively.

Table 2 The K_d and K_{oc} values of four selected antibiotics

Antibiotics	K_d (L/Kg)		K_{oc}	
	This study (mean)	References	This study	References
OFL	2982	1192~4525 ^a	447300	50056~1104595 ^a
RTM	1420	470 ^b	213000	-
ETM	337	165 ^c	50550	-
SMZ	89	0.23~43.1 ^a	13350	62.2~607 ^a

^a From (Drillia et al. 2005)

^b From (Gobel et al. 2005)

^c From (Jones et al. 2002)

Table 3 The DT₅₀ and DT₉₀ values of the four antibiotics in seawater and sediment


Antibiotics	In the human body (h)		In the flume system (d)		In the quiescent system (d)	
	DT ₅₀	DT ₉₀	DT ₅₀	DT ₉₀	DT ₅₀	DT ₉₀
OFL	5.0~10.0	— ^a	3.4 ^b (21.4) ^c	7.5 (>60)	12.9 (34.0)	>60 (>60)
RTM	8.4 ~15.5	—	6.7 (2.3)	30 (>60)	29 (41.1)	>60 (>60)
ETM	1.4 ~2	—	7.3 (2.1)	>60 (>60)	18 (27.0)	>60 (>60)
SMZ	8.0~12.0	—	14.7 (3.1)	>60 (29)	14.5 (24.3)	>60 (>60)

^a Not available

^b In seawater

^c In sediment

645



646

647 **Fig. 1** The setting of the experimental flume (LEAF)

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673




Fig. 2 Relationship of the lid rotational speed (RS) with the averaged flow velocity ($\langle u \rangle$) and the bed shear stress (τ_b) (adopted from Chan et al. 2006)




Fig. 3 Temporal changes of the four antibiotics in surface water (a), bottom water (b), and sediment (c) of the flume system




Fig. 4 Temporal changes of the four antibiotics in surface water (a) and sediment (b) of the quiescent system




Fig. 5 Concentration profiles of the four antibiotics in sediment layers of the flume system