

The spatial and temporal distribution of heavy metals in sediments of Victoria Harbour, Hong Kong

Chloe Wing-yee Tang^a, Carman Ching-man Ip^a, Gan Zhang^b, Paul, K.S. Shin^c, Pei-yuan Qian^d, and Xiang-dong Li^{a,*}

^aDepartment of Civil & Structural Engineering, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong

^bState Key Laboratory of Organic Geochemistry, Guangzhou Institute of Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China

^cDepartment of Biology and Chemistry, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong

^dDepartment of Biology & Coastal Marine Laboratory, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong

Abstract

Victoria Harbour has received substantial loadings of pollutants from industrial and municipal wastewater discharged since the 1950s. Inputs of contaminants have declined dramatically during the last two decades as a result of better controls at the source and improved wastewater treatment facilities. To assess the spatial and temporal changes of metal contaminants in sediments in Victoria Harbour, core and grab sediments were collected. The central harbour areas were generally contaminated with heavy metals. The spatial distribution of trace metals can probably be attributed to the proximity of major urban and industrial discharge points, and to the effect of tidal flushing in the harbour. In the sediment cores, the highest concentrations of trace metals were observed to have accumulated during the 1950s to 1980s, corresponding with the

* Corresponding author. Tel.: + 852-2766-6041; Fax: + 852-2334-6389.
E-mail address: cexdli@polyu.edu.hk (X.D. Li).

period of rapid urban and industrial development in Hong Kong. From the late 1980s, there has been a major decline in the concentrations of trace metals, due to a reduction in industrial activities and to the enactment of wastewater pollution controls in the territory. The Pb isotopic compositions of the sediments revealed the anthropogenic inputs of Pb to the harbour. The $^{206}\text{Pb}/^{207}\text{Pb}$ ratios varied from 1.154 to 1.197, which were lower than those of background geological materials in Hong Kong ($^{206}\text{Pb}/^{207}\text{Pb}$: 1.201–1.279). The data also indicated that the Pb in the harbour sediments most likely originated from mixed sources, including the leaded gasoline used in the past and other anthropogenic sources.

Keywords: Trace metals; Pb isotopes; Sediment; Historical sources; Victoria Harbour; Hong Kong

1. Introduction

Victoria Harbour is located in the southeastern part of the Pearl River Estuary (PRE) in south China. It is a major port of Hong Kong, and lies between the most urbanized areas of the Kowloon Peninsula and the northern shore of Hong Kong Island. The harbour has strong tidal flushing, whereby seawater flows from the east at Lei Yue Mun Channel to west of the harbour during flood tides, and in the opposite direction during ebb tides. Water in the harbour is further influenced by discharges of freshwater from the PRE in the west (Morton and Wu, 1975). Victoria Harbour has been subjected to significant environmental pollution from increasing urban and industrial development in the last few decades. The major sources of pollution include municipal wastewater, urban runoff, and industrial effluents. In 1995, the harbour received an estimated 1.5

million tonnes of sewage and industrial wastewater every day (Hong Kong Government, 1995). As a consequence, there was a rapid decline in the quality of the water in the harbour, as bacteria, organic materials, and other pollutants seriously affected fish and other marine life by depleting oxygen levels in the water (Yung et al., 1999). However, the input of contaminants to Victoria Harbour has been greatly reduced in recent years, due to enhanced efforts to control the problem at the source, improved wastewater treatment facilities, and increased public awareness (HKEPD, 2004).

Once contaminants are released into the aquatic environment, they may be removed from the water column by interacting with suspended solids and subsequently be deposited as bottom sediments (Croudace and Cundy, 1995). Coastal sediments are considered to be a major sink for, and also a secondary source of pollutants. The fate of contaminants in sediments depends on a number of factors, including the resuspension, bioturbation, and diagenesis of sediments (Bothner et al., 1998; Lee and Cundy et al., 2001). Heavy metals, unlike other pollutants, are not biodegradable and can accumulate in sediments over time. The contamination of metals in sediments can reach a level that is toxic to aquatic life. Metals that have accumulated in sediment profiles can also act as a historical record of human activities in a watershed and provide insight on the accumulation of metal contaminants over time (Finney and Huh, 1989).

Sedimentation rates and sediment chronology in sediment cores are now commonly estimated by the ^{210}Pb dating technique. This dating technique has proven to be successful for determining dates back to about 100 years and is widely used in coastal marine, estuarine, and marsh environments (i.e., Hornberger et al., 1999; Bay et al., 2003; Ip et al., 2004). As a common metal contaminant, stable Pb isotopes have frequently been used to identify different sources of lead in the environment (Sturges and Barrie, 1987; Erel et al., 2001; Zhu et al., 2001). Anthropogenic Pb is derived from

lead sulfide ore deposits and is released into the environment through the combustion of leaded gasoline and as a byproduct of industrial Pb use. In general, the Pb in most ore bodies is less radiogenic and has a lower $^{206}\text{Pb}/^{207}\text{Pb}$ ratio than that found in most natural bedrock. The inherited differences in Pb isotopic signatures between natural and anthropogenic Pb, thus allow Pb isotopic compositions to be used to identify Pb sources and, in some cases, to provide source apportionment for potential sources of Pb (Farmer et al., 1996).

A number of previous research projects have focused on studying trace metals in surface sediments of Victoria Harbour. The sediments in Victoria harbour have been found to be highly contaminated with Cu, Zn, Cr, and Pb due to the discharge of untreated industrial waste and sewage since the 1960s (Yim, 1981; Wong et al., 1995; HKEPD, 2003). Metals in the sediments of the harbour have also been found to induce toxic effects in fish and algal bioassays (Wong et al., 1995; Wong et al., 2000). Over the last two decades, measures aimed at improving the environment, such as the enactment of the Water Pollution Ordinance, the introduction of unleaded petrol, the operation of a chemical waste treatment centre, and the implementation of the Harbour Area Treatment Scheme (HATS) have resulted in a reduction in the amount of metals being discharged into Victoria Harbour (HKEPD, 2000). However, there have been very few studies focusing on the changes in the metal contaminants in sediments due to such a reduction. It is of importance to assess the changes in the contamination of sediments over time, particularly the recent two decades. The present study aims to study the current contamination of trace metals and the historical fluxes of metals in Victoria Harbour, and to evaluate the potential sources of metal pollutants based on the geochemical features and Pb isotopic compositions in the sediments.

2. Methodology

2.1. Sediment sampling

Five sampling sites from east (B1) to west (B5) of Victoria Harbour were selected for collecting core and grab sediments (Figure 1). Grab surface sediments were collected in Victoria Harbour during the four cruise outings in December 2004, February 2005, October 2005, and April 2006, respectively, to provide representative data on the spatial distribution of trace metals in the harbour. Core sediments were collected in October 2004 and used for the study of the temporal distribution of metals in the profiles. These locations have good coverage of the harbour area. Moreover, the sampling locations were chosen with the aim of avoiding areas with known sediment disturbance, such as near shore areas where intensive reclamation activities are taking place and dredged areas. Two grab sediment samples (a and b) were collected within 50 m of each sampling station. The samples of grab sediment were taken with a Van Veen Grab Sampler. A core of sediment was taken at each sampling site using a Kajak gravity corer[®]. The corer was driven into the sediment by gravity, and the sediment core was retained in a PVC tube. The diameter of the PVC coring tube was 70 mm, and about 60–65 cm sediment profiles were obtained. Each core was sliced into thin sections at 2 cm intervals. All of the samples were then stored in polyethylene bags at 4–6°C prior to undergoing laboratory analyses. The physical-chemical parameters, including the salinity, pH, and water depth of the harbour waters, were measured in situ using YSI 6600 Sonde. In the laboratory, the size of the particles in the core samples was analyzed using a laser diffraction particle sizer “Coulter[®] LS100”. The organic carbon was determined in freeze-dried sediments, after acidification with 1M HCl, using a CHNS

Perkin Elmer 2400 elemental analyzer, with a precision of 0.1 µg.

2.2. Trace metal and major element analysis

For the analysis of the concentrations of metals, sediment samples were freeze-dried and then ground in an agate grinder until fine particles were obtained. Sediment samples were analyzed for concentrations of major and trace elements using the nitric and perchloric acid digestion method (Ip et al., 2004). About 0.2 g of the samples of ground sediment were digested with a mixture of 6 ml of nitric acid and 1.5 ml of perchloric acid in a heating block at 50°C for 3 h, 100°C for 1 h, 125°C for 1 h, 150°C for 3 h, 175°C for 2 h and 190°C until completely dry. The residual was then leached with 10 ml of 5% (v/v) nitric acid and heated at 70°C for 1 h. Trace and major elements in the solutions were determined by inductively coupled plasma-atomic emission spectrometry (ICP-AES, Perkin-Elmer Optima 3300DV). All plastic materials and glassware were soaked in 10% nitric acid overnight and rinsed thoroughly with de-ionized water before use. For quality assurance, reagent blanks, sample replicates, and standard reference materials (Buffalo River Sediment–NIST SRM 8704) were used to assess the accuracy and precision of the analysis. The analytical results showed no signs of contamination and the precision and bias were less than 10%. In the standard reference material (NIST SRM 8704), the recovery rates of trace metals were around 80–105%. For Fe and Al, the recovery rates were around 50–60% due to the incomplete digestion of aluminosilicate minerals without hydrofluoric acid (HF).

2.3. Pb isotopic analyses

Selected core samples were used for an analysis of the Pb isotopic composition. Solutions from digested sediment samples were diluted using a 5% high purity nitric acid solution and analysed for stable Pb isotope ratios by inductively coupled plasma-mass spectrometry (ICP-MS, Perkin Elmer ELAN 6100 DRC^{plus}). An instrumental parameter of 150 sweeps/reading was used. The relative standard deviation (RSD) of the ten replicate readings of each sample was <0.5%. An international standard reference material (the SRM 981 Common Pb Isotopic Standard) from the National Institute of Standards and Technology (NIST) was used for calibration and quality control. It was inserted at the beginning of each run, between every five samples, and at the end of each run, respectively. The average measured ratios of $^{204}\text{Pb}/^{207}\text{Pb}$, $^{206}\text{Pb}/^{207}\text{Pb}$, and $^{208}\text{Pb}/^{207}\text{Pb}$ of the standard reference materials were 0.0645 ± 0.0002 , 1.093 ± 0.002 , and 2.368 ± 0.004 , respectively. These results were in good agreement with the certified standard values (0.0646, 1.0933, and 2.3704, respectively).

2.4. ^{210}Pb -dating of sediment cores

The ^{210}Pb radiometric technique was used to estimate the chronology of sediment cores. The ^{210}Pb activities of the samples were determined by measuring the alpha-radioactivity of its granddaughter nuclide ^{210}Po . The ^{210}Po was extracted, purified, and self-plated onto a silver disc at 70–80 °C in 0.5M HCl, and ^{209}Po was used as the yield monitor and tracer in quantification. The alpha-activity was counted by computerized multi-channel alpha spectrometry with gold-silicon surface barrier detectors. The supported ^{210}Pb was determined by the alpha activity of the supporting parent, ^{226}Ra , via the co-precipitation of BaSO_4 . In this study, the constant flux/constant sedimentation rate (CF/CS) model was adopted to calculate the rate of sedimentation

(Appleby and Oldfield, 1983).

2.5. Statistical analysis

A one-way ANOVA and a Student *t*-test were performed to detect whether there were any significant differences in particle sizes and metal concentrations among the sediment samples collected at different sampling periods as well as various sampling locations. A Pearson correlation was performed to identify metal associations in harbour sediments. The critical level for all statistical analyses was carried out at a 95% confidence interval ($p < 0.05$). A factor analysis was also applied to identify the major source(s) of metals in the harbour. It was performed on the concentrations of metals in all sediment cores, and Varimax with Kaiser normalization was used as the rotation method in the analysis. All statistical analyses were performed using SPSS® for Window Release 10.1.

3. Results and Discussion

3.1. Physical-chemical properties of harbour sediments

The depth, salinity, pH of the water, and total organic carbon (TOC) of surface sediments in Victoria Harbour are given in Figure 2. In general, the water depths in the harbour varied from 7 to 19 m, with increasing from west to east. The pH was consistent from site to site, with an average of 8.1, which was slightly alkaline and within the normal range for marine water. The salinity in the eastern part was slightly higher than that in the western part, indicating the influence of oceanic water from the west Pacific

and of the discharge of freshwater from the PRE. The amount of organic carbon in surface sediments ranged from 0.4 to 1.87%, with the highest TOC found in the central part of the harbour and the lowest in the eastern part. A higher amount of TOC in the central part was most likely due to the discharge of untreated urban runoff.

The grain sizes of the sediment cores are shown in Table 1. The sediments displayed significant variations in grain size among different sampling locations ($p < 0.05$). In general, the finer-grained sediments (with 97% $< 63 \mu\text{m}$) were found at B5, whereas coarser sediments (with 52–80% $< 63 \mu\text{m}$) were found at B1 and B2, as the latter locations were affected by a strong tidal flow. It is well known that trace metals tend to be adsorbed on the surfaces of particles and that higher concentrations of metals are usually associated with smaller particles (Daskalakis and O'Connor, 1995; Schiff and Weisberg, 1999). In Victoria Harbour, trace metals, including Cu, Pb, and Zn, showed no significant correlation with the percentage of fine-grained particles in the sediments ($p > 0.05$) (Figure 3). Therefore, the result indicates that anthropogenic inputs were a more significant control on the distribution of trace metals in the harbour sediments than natural geochemical process.

3.2. Spatial distribution of trace metals in harbour sediments

The concentrations of metals in the sediments showed significant differences among the sampling locations, but no major difference during the four sampling periods (ANOVA, $p > 0.05$). The concentrations of Cu, Pb, and Zn in the four sampling periods were averaged and presented in Table 2. The present findings were consistent with a recent study conducted by the Hong Kong Environmental Protection Department (HKEPD, 2004). According to the Student *t*-test analysis, the degree of Cu enrichment

was found to be $B2 \sim B3 > B4 > B5 > B1$, while that of Pb and Zn was $B2 \sim B3 > B4 \sim B5 > B1$. The central part of the harbour had the largest amounts of Cu, Pb, and Zn, with mean concentrations of 173, 44.3, and 171 mg/kg, respectively. The significant spatial variation of the metals could be explained by the different urban settings in the territory. The substantial enrichment of trace metals (Cu, Pb, and Zn) found in the central part (B2 & B3) was probably caused by the proximity of the location to the most heavily urbanized areas. These areas have a high density of population, a high density of traffic, and are close to old industrial districts. In contrast, B1 sediments with less enrichment of metals are located in an area of strong tidal flushing, with a low density of population and less anthropogenic activity.

The results were compared with the baseline values taken from the measurements in sediment cores from Victoria Harbour, which were typically deeper than 5 m (Tanner et al., 2000). There was a significant enrichment of trace metals in the central and western parts of the harbour, particularly for Cu in the central part, which was 20 times higher than the baseline value (Table 2). The high concentrations of Cu in surface sediments also exceeded the Upper Chemical Exceedance Level of Hong Kong, which may pose some hazards to aquatic life. Further study on the chemical form / bioavailability of the Cu present in these sediments is required.

The results obtained from the current study are also compared with those from other harbours in the world (Table 3). The levels of Pb and Zn in surface sediments of Victoria Harbour are comparable to those of moderately polluted harbours with small catchments and a relatively short history of environmental pollution, such as Darwin Harbour (Peerzada and Rohoza, 1989), Manila Bay (Prudente et al., 1994), and Tolo Harbour (Owen and Sandhu, 2000). In contrast, the Cu in Victoria Harbour was within the ranges reported in New Bedford Harbour (Stoffers et al., 1977), Baltimore Harbour

(McGee et al., 1999), and Sydney Harbour (Birch and Taylor, 1999). These are heavily polluted harbours that receive substantial loads of pollutants from river drainages and that have been subjected to pollution for around 100 to 200 years.

3.3. Temporal distribution of trace metals in harbour sediments

As sediments in Victoria Harbour are highly mixed by physical processes, such as tidal currents and/or human disturbances, it was only possible to successfully construct the chronology for sediment core B4 using the ^{210}Pb dating method. The unsupported ^{210}Pb activities of the core decreased linearly as a function of sediment depth in core B4 while cores B2 and B5 showed non-linear patterns of unsupported ^{210}Pb (Figure 4). The linear ^{210}Pb profiles indicated an exponential decline in ^{210}Pb activity with depth, and this allowed the CF/CS model to be used to calculate the rate of sedimentation (Appleby and Oldfield, 1983). The average sedimentation rate estimated by the CF/CS model in core B4 was 1.2 cm/yr. The concentration profiles of Cu, Zn, and Pb at B1 to B5 are shown in Figure 5. The sediment of core B4 showed significant temporal changes in trace metal concentrations, particularly for Cu and Zn, indicating changing inputs of metal to the harbour during the last few decades. The trend for Cu closely followed the trend for Zn in the sediment core, and a significant correlation between these two metals was observed ($p < 0.05$) (Figure 6). This raises the possibility that the two metals probably derived from the same source(s) of industrial emissions and other inputs.

The elevated Cu and Zn values in the sediments were observed in the mid-1960s, and significant increases of both metals occurred in the 1970s. The concentrations Cu and Zn reached peak values at 33 cm in the sediment profile corresponding to 1977,

with maximum concentrations of 91 and 155 mg/kg, respectively. Such a change could be related to the urban and industrial development that was taking place in the territory. Since the 1950s, the manufacturing industry in the territory had grown substantially and remained the largest single contributor to local Gross Domestic Product (GDP) until 1987 (HKGID, 1996). Therefore, intensive industrial activities led to significant inputs of metal in the harbour during the 1950s to the 1980s. During that time, the peak concentration of Cu in the sediments was 8 times higher than the baseline value, and increased to 1.7 times of the Cu concentration in the deep sediments of the 1950s. The enrichment of Cu was most likely related to the large-scale manufacturing of printed circuit boards (PCBs) and electroplating industry in the territory before the 1980s (Yim, 1984; Tanner et al., 2000; HKEPD, 2002).

A decline was observed in metal concentrations in the sediment core from the late 1980s, which could reflect the reduction in industrial activities due to the relocation of industries to mainland China and the setting up of the Chemical Waste Treatment Centre in the territory. The current concentrations of Cu and Zn are much lower than the peak values in the 1970s, the estimated reduction of metals being 45% and 27% for Cu and Zn, respectively.

The trend of the Pb profile was slightly different from that of Cu and Zn at B4 (Figure 5). The deeper sediments had an average Pb concentration of 37.5 mg/kg, higher than the baseline concentration in Hong Kong marine sediment (see Table 3). A rise in the concentrations of Pb to 42–49 mg/kg occurred during the period 1970 to 1982. Thereafter, the concentrations of Pb declined to 38.2 mg/kg, although an abrupt increase to 47.8 mg/kg was observed in 1992. The surface sediments were characterized by declining concentrations of Pb, with 35.7 mg/kg Pb in the topmost sediment. In Hong Kong, a gradual reduction in the amount of Pb permitted in leaded petrol began in the

early 1980s, and leaded petrol was totally banned in 1999. Consequently, the increased enrichment of Pb during the period from 1970 to 1982 in the B4 core can be attributed to the use of leaded gasoline in the territory and to the discharges of industrial wastewater (Yim, 1984; Wong, 1987; Chan et al., 2001).

In general, the distribution of trace metals in Victoria Harbour varied spatially and temporally in the surface and core sediments in the eastern, central, and western parts of the harbour. Spatial variations of trace metals in sediments are likely caused by discharges of sewage, industrial activities, and surface runoff along the coastal areas. The rapid urban and industrial development that has taken place in Hong Kong since the mid-1950s led to a relatively slight increase for Zn and Pb, with a greater increase for Cu in subsurface sediments. Subsequent declines in trace metals were observed to have occurred from the late-1980s, due to a gradual reduction in industrial discharges and to better pollution controls in the territory.

3.4. Potential sources of trace metals in the harbour

To further elucidate the potential sources of trace metals in Victoria Harbour, the associations of trace elements in sediments were studied using factor analysis. The results of the factor analysis of the total concentrations of metals (Al, Fe, Mn, Ca, Co, Cu, Ni, Pb and Zn) in the sediment cores are shown in Table 4.

In this analysis, three major components were extracted. They accounted for 83.7% of the total variance. The first component, which includes Al, Fe, Ca, Co, and Ni, may represent the natural source of the sediments. The second component, including Cu, Pb, and Zn, probably represents anthropogenic sources associated with the discharge of sewage, industrial inputs, and surface runoff in the urban environment. The third

component consists of Mn in weak association with Fe and Ca, and this probably indicating the geochemical process of Mn and Fe oxides in the sediments.

3.5. Pb isotopic composition of harbour sediments

The mean and ranges of the $^{206}\text{Pb}/^{207}\text{Pb}$ ratios in the sediments are listed in Table 5. The range of the Pb isotope ratios varied from 1.154 to 1.197 in these sediments. In general, the $^{206}\text{Pb}/^{207}\text{Pb}$ isotopic ratios of the anthropogenic Pb derived from various Pb ores were usually lower than those of background geological materials (Croudace and Cundy, 1995; Farmer and Eades, 1996). The natural background of $^{206}\text{Pb}/^{207}\text{Pb}$ isotopic ratios in Hong Kong ranged from 1.201 to 1.279 (Ip et al., 2004), while lower values were observed in the harbour sediments. This may provide strong evidence that the Pb in Victoria Harbour derives from anthropogenic sources. Among the sediment cores, B1 was characterized by relatively low concentrations of Pb and high $^{206}\text{Pb}/^{207}\text{Pb}$ ratios, which were close to the natural background ratios. Thus, the eastern part of the harbour is probably less contaminated with anthropogenic Pb. The $^{206}\text{Pb}/^{207}\text{Pb}$ ratios at the B4 core ranged from 1.171 to 1.190, whereas noticeably lower Pb isotope ratios with a corresponding increase in the concentrations of Pb were observed at a depth of 32 cm (Figure 7). This shift occurred in the sediments that had been deposited since the 1970s, giving a strong indication that inputs of anthropogenic Pb increased as a result of rapid urban and industrial development in the territory.

To evaluate the potential anthropogenic sources of Pb in harbour sediments, the $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios of the samples, geological materials in Hong Kong, road dust in Hong Kong, contaminated soils in Hong Kong, and Australian Pb ore are plotted in Figure 8. The $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{207}\text{Pb}$ ratios of the harbour sediments and

the geological materials formed a linear mixing pattern with those of contaminated urban soils and road dust. This probably suggests that the Pb contamination in the sediments is associated with the input of gasoline lead to the harbour, perhaps from the stormwater runoff of surface soils in the urban setting. The wide range of Pb isotopic ratios in harbour sediments can most likely be attributed to the mixed sources of gasoline Pb and other anthropogenic sources in the harbour, e.g. industrial and shipping activities.

4. Conclusions

Trace metals in sediments of Victoria Harbour exhibited considerable spatial and temporal variability as a result of various pollutant loadings over the past few decades. The most contaminated site was found in the central part of the harbour, which is close to urban discharge points and former industrial areas. High concentrations of Cu (up to 460 mg/kg) were found in the subsurface sediments in the central part of the harbour. These can be attributed to the intensive industrial activities that took place in the territory and to the discharge of untreated sewage into the harbour from the mid-1950s to the late-1980s. The anthropogenic source of the Pb was reflected by the lower $^{206}\text{Pb}/^{207}\text{Pb}$ ratios in the harbour sediments. The Pb isotopic signatures showed that the elevated levels of Pb in the harbour were the mixed product of past gasoline Pb and a number of other anthropogenic inputs.

Acknowledgements

This work was supported by the Area of Excellence (AoE) project under Grant No. AoE/P-04/2004 from the University Grant Council of Hong Kong. We thank the

two reviewers on their constructive comments and suggestions which are very helpful in improving the quality of the manuscript.

References

- Appleby, P.G., Oldfield, F., 1983. The assessment of ^{210}Pb data from sites with varying sediment accumulation rates. *Hydrobiologia* 103, 29–35.
- Bay, S.M., Zeng, E.Y., Lorenson, T.D., Tran, K., Alexander, C., 2003. Temporal and spatial distributions of contaminants in sediments of Santa Monica Bay, California. *Marine Environmental Research* 56, 255–276.
- Birch, G., Taylor, S., 1999. Source of heavy metals in sediments of the Port Jackson estuary, Australia. *The Science of the Total Environment* 227, 123–138.
- Bothner, M.H., Buchholtz ten Brink, M., Manheim, F.T., 1998. Metal concentrations in surface sediments of Boston Harbor-changes with time. *Marine Environmental Research* 45, 127–155.
- Chan, L.S., Ng, S.L., Davis, A.M., Yim, W.S., Yeung, C.H., 2001. Magnetic properties and heavy metal contents of contaminated seabed sediments of Penny's Bay, Hong Kong. *Marine Pollution Bulletin* 42, 569–583.
- Croudace, I.W., Cundy, A.B., 1995. Heavy metal and hydrocarbon pollution in recent sediments from Southampton water, Southern England: A geochemical and isotopic study. *Environmental Science and Technology* 29, 1288–1296.
- Daskalakis, K.D., O'Connor, T.P., 1995. Normalization and elemental sediment contamination in the Coastal United States. *Environmental Science and Technology* 29, 470–477.
- Duzgoren-Aydin, N.S., Li, X.D., Wong, S.C., 2004. Lead contamination and isotope signatures in the urban environment of Hong Kong. *Environmental International* 30, 209–217.
- Erel, Y., Dubowski, Y., Halicz, L., Erez, J., Kaufman, A., 2001. Lead concentrations and isotopic ratios in the sediments of the Sea of Galilee. *Environmental Science and Technology* 35, 292–299.
- Farmer, J.G., Eades, L.J., Mackenzie, A.B., Kirika, A., Bailey-Watts, T.E., 1996. Stable lead isotope record of lead pollution in Loch Lomond sediments since 1630 A.D.

- Environmental Science and Technology 30, 3080–3083.
- Finney, B.P., Huh, C.A., 1989. History of metal pollution in the Southern California Bight: an update. Environmental Science and Technology 23, 294–303.
- HKEPD, 1996–2004. Marine Water Quality in Hong Kong. Environmental Protection Department, Hong Kong Government, Hong Kong.
- Hong Kong Government, 1995. The shape of things to come – an overview of the role of harbour reclamation in the future development of Hong Kong. Hong Kong Government: Planning, Environment and Lands Branch, pp. 136.
- HKGID, 1996. Hong Kong's Manufacturing Industries. Hong Kong Government Industry Department, Hong Kong Government, Hong Kong.
- Hornberger, M.L., Luoma, S.N., Green, A., Fuller, C., Anima, R., 1999. Historical trends of metals in the sediments of San Francisco Bay, California. Marine Chemistry 64, 39–55.
- Ip, C.C.M., Li, X.D., Zhang, G., Farmer, J.G., Wai, O.W.H., Li, Y.S., 2004. Over one hundred years of trace metal fluxes in the sediments of the Peal River Estuary, South China. Environmental Pollution 132, 157–172.
- Lee, S.V., Cundy, A.B., 2001. Heavy metal contamination and mixing processes in sediments from the Humber Estuary, Eastern England. Estuarine, Coastal and Shelf Science 53, 619–636.
- McGee, B.L., Fisher, D.J., Yonkos, L.T., Ziegler, G.P., Turley, S., 1999. Assessment of sediment contamination, acute toxicity, and population viability of the estuarine amphipod *Leptocheirus plumulosus* in Baltimore Harbor, Maryland, USA. Environmental Toxicology and Chemistry 18, 2151–2160.
- Morton, B., Wu, S.S., 1975. The hydrology of the coastal waters of Hong Kong. Environmental Research 10, 319–347.
- Owen, R.B., Sandhu, N., 2000. Heavy metal accumulation and anthropogenic impacts on Tolo Harbour, Hong Kong. Marine Pollution Bulletin 40, 174–180.
- Peerzada, N., Rohoza, W., 1989. Some heavy metals in sediments from Darwin Harbour, Australia. Marine Pollution Bulletin 20, 91–92.
- Prudente, M.S., Ichihashi, H., Tatsukawa, R., 1994. Heavy metal concentrations in sediments from Manila Bay, Philippines and inflowing rivers. Environmental Pollution 86, 83–88.
- Schiff, K.C., Weisberg, S.B., 1999. Iron as a reference element for determining trace metal enrichment in Southern California coastal shelf sediments. Marine

- Environmental Research 48, 161–176.
- Stoffers, P., Summerhayes, C., Forstner, U., Patchineelam, S.R., 1977. Copper and other heavy metal contamination in sediments from New Bedford Harbor, Massachusetts: A preliminary note. *Environmental Science and Technology* 11, 819–821.
- Sturges, W.T., Barrie, L.A., 1987. Lead-206/207 isotope ratios in the atmosphere of north-America as tracers of United States and Canadian emissions. *Nature* 329, 144–146.
- Tanner, P.A., Leong, L.S., Pan, S.M., 2000. Contamination of heavy metals in marine sediment cores from Victoria Harbour, Hong Kong. *Marine Pollution Bulletin* 40, 769–779.
- Wong, C.K.C., Yeung, H.Y., Cheung, Y.H., Yung, K.K.L., Wong, M.H., 2000. Ecotoxicological assessment of persistent organic and heavy metal contamination in Hong Kong coastal sediment. *Archives of Environmental Contamination and Toxicology* 38, 486–493.
- Wong, C.S.C., Li, X.D., 2004. Pb contamination and isotopic composition of urban soils in Hong Kong. *The Science of the Total Environment* 319, 185–195.
- Wong, M.H., 1987. A review on lead contamination of Hong Kong's environment. In Hutchinson, T.C. and Meema, K.M., eds., *Lead, Mercury, Cadmium and Arsenic in the Environment*, pp. 217–233. John Wiley & Sons Ltd.
- Wong, Y.S., Tam, N.F.Y., Lau, P.S., Xue, X.Z., 1995. The toxicity of marine sediments in Victoria Harbour, Hong Kong. *Marine Pollution Bulletin* 31, 464–470.
- Yim, W.W.S., 1981. Heavy metal in marine sediments of Hong Kong. *Hong Kong Engineer*, 33–39.
- Yim, W.W.S., 1984. Geochemical mapping of bottom sediments as an aid to marine waste disposal in Hong Kong. *Conservation & Recycling* 7, 309–320.
- Yung, Y.K., Yau, K., Wong, C.K., Chan, K.K., Yeung, I., Kueh, S.W., Broom, M.J., 1999. Some observations on the changes of physico-chemical and biological factors in Victoria Harbour and vicinity, Hong Kong, 1988-1996. *Marine Pollution Bulletin* 39, 315–325.
- Zago, C., Giblin, A.E., Bergamasco, A., 2001. Changes in the metal content of surficial sediments of Boston Harbor since the cessation of sludge discharge. *Marine Environmental Research* 51, 389–415.
- Zhu, B.Q., Chen, Y.W., Peng, J.H., 2001. Lead isotope geochemistry of the urban

environment in the Pearl River Delta. *Applied Geochemistry* 16, 409–417.

Table 1. Grain size distribution in selected samples of sediment from locations B1 to B5 in Victoria Harbour

| Location | Depth (cm) | % fine-grained sediment ($< 63\mu\text{m}$) |
|----------|------------|--|
| B1 | 4–5 | 80.1 |
| | 16–18 | 57.4 |
| | 24–26 | 71.0 |
| | 36–38 | 79.4 |
| | 52–54 | 61.5 |
| B2 | 3–4 | 51.9 |
| | 7–8 | 68.8 |
| | 14–16 | 68.5 |
| | 20–22 | 61.7 |
| B3 | 2–4 | 84.9 |
| | 10–12 | 75.4 |
| | 16–18 | 96.4 |
| | 24–26 | 99.9 |
| B4 | 2–3 | 80.8 |
| | 7–8 | 80.5 |
| | 14–16 | 86.9 |
| | 32–34 | 74.3 |
| B5 | 1–2 | 95.2 |
| | 5–6 | 99.9 |
| | 14–16 | 99.1 |
| | 20–22 | 99.2 |

Table 2. Mean (\pm S.D.) concentrations of Cu, Pb, and Zn (mg/kg, dry weight) in surface sediments of Victoria Harbour collected in December 2004, February 2005, October 2005, and April 2006 (n=8)

| | Cu (mg/kg) | Pb (mg/kg) | Zn (mg/kg) |
|---|------------------------------|------------------------------|-----------------------------|
| B1 | 21.2 \pm 4.5 ^a | 19.6 \pm 3.3 ^a | 55.0 \pm 5.3 ^a |
| B2 | 172 \pm 34.3 ^b | 44.0 \pm 17.1 ^b | 165 \pm 28.9 ^b |
| B3 | 173 \pm 60.5 ^b | 44.6 \pm 6.06 ^b | 176 \pm 27.7 ^b |
| B4 | 51.4 \pm 14.5 ^c | 30.8 \pm 6.89 ^c | 115 \pm 16.3 ^c |
| B5 | 30.2 \pm 8.01 ^d | 30.7 \pm 5.45 ^c | 109 \pm 6.76 ^c |
| LCEL ¹ | 65 | 75 | 200 |
| UCEL ² | 110 | 110 | 270 |
| HKEPD ³ | 17–230 | 24–130 | 47–300 |
| Background values in Hong Kong sediments ⁴ | 11 | 32 | 67 |

The small letters in the columns stand for statistical significance at the 0.05 level with a Student *t*-test among the sampling locations.

¹LCEL Lower Chemical Exceedance Level for assessments of sediment quality in Hong Kong. Biological effects would be observed above this level (HKEPD, 2004).

²UCEL Upper Chemical Exceedance Level for assessments of sediment quality in Hong Kong. Adverse biological effects would be observed above this level (HKEPD, 2004).

³HKEPD Data from an HKEPD sediment monitoring programme conducted in 2003 (HKEPD, 2004).

⁴Tanner et al., 2000

Table 3. Metal concentrations in the sediments of other polluted harbours (mg/kg, dry weight)

| Location | Cu | Pb | Zn |
|--|----------|---------|----------|
| Boston Harbour, U.S.A. ¹ | 7–142 | 18–263 | 39–414 |
| New Bedford Harbour, U.S.A. ² | 117–3136 | 104–616 | 315–1550 |
| Baltimore Harbour, U.S.A. ³ | 9.5–396 | 1.0–349 | 40–2105 |
| Sydney Harbour, Australia ⁴ | 9.3–1053 | 38–3604 | 108–7622 |
| Darwin Harbour, Australia ⁵ | 16–32 | 24–91 | 103–270 |
| Manila Bay, Philippines ⁶ | 32–118 | 6.0–95 | 60–329 |
| Tolo Harbour, Hong Kong ⁷ | 21–84 | 48–144 | 100–270 |
| Victoria Harbour, Hong Kong ⁸ | 16–280 | 21–85 | 52–221 |

¹Zago et al., 2001

²Stoffers et al., 1977

³McGee et al., 1999

⁴Birch and Taylor, 1999

⁵Peerzada and Rohoza, 1989

⁶Prudente et al., 1994

⁷Owen and Sandhu, 2000

⁸The present study

Table 4. Result of the factor analysis on total concentrations of metals in B1 to B5 sediment cores (n=116)

| % of variance | Principle loading factors | | |
|---------------|---------------------------|----------------|----------------|
| | PC 1 37.3 % | PC 2 28.9 % | PC 3 17.6 % |
| Cu | -0.083 | 0.732 | -0.508 |
| Ni | 0.782 | 0.202 | -0.276 |
| Pb | 0.051 | 0.949 | -0.004 |
| Zn | 0.222 | 0.863 | -0.248 |
| Al | 0.870 | 0.351 | -0.019 |
| Fe | 0.842 | -0.203 | <i>0.449</i> |
| Mn | -0.002 | -0.190 | 0.924 |
| Co | 0.933 | -0.188 | 0.181 |
| Ca | -0.594 | -0.371 | <i>0.300</i> |

Table 5. Mean (\pm S.D.) of $^{206}\text{Pb}/^{207}\text{Pb}$ ratios and range of Pb concentrations and $^{206}\text{Pb}/^{207}\text{Pb}$ ratios in sediment cores, B1 to B5, of Victoria Harbour

| | Range of Pb concentration (mg/kg) | Mean $^{206}\text{Pb}/^{207}\text{Pb}$ | Range $^{206}\text{Pb}/^{207}\text{Pb}$ |
|-----------|--------------------------------------|--|---|
| B1 (n=5) | 25.9 – 32.4 | 1.181 ± 0.005 | 1.173 – 1.185 |
| B2 (n=5) | 35.1 – 56.9 | 1.175 ± 0.006 | 1.168 – 1.182 |
| B3 (n=5) | 39.8 – 52.2 | 1.174 ± 0.005 | 1.168 – 1.180 |
| B4 (n=22) | 33.2 – 52.7 | 1.182 ± 0.012 | 1.170 – 1.190 |
| B5 (n=5) | 33.7 – 45.0 | 1.175 ± 0.015 | 1.154 – 1.189 |

Figure Captions

Figure 1. The sampling locations, B1–B5, in Victoria Harbour

Figure 2. Physical parameters of the water (water depth, salinity, and pH) and the total organic carbon (TOC) of surface sediments in Victoria Harbour at different sampling periods

Figure 3. The correlation of Cu, Zn, and Pb with the percentage of fine-grained particles in Victoria Harbour

Figure 4. Results of ^{210}Pb dating in sediment cores B2, B4, and B5 of Victoria Harbour

Figure 5. Temporal distribution of Cu, Zn, and Pb in core sediments, B1 to B5, of Victoria Harbour (Core B4 with the estimated date)

Figure 6. The correlation between Cu and Zn in sediments of core B4

Figure 7. $^{206}\text{Pb}/^{207}\text{Pb}$ ratios and Pb concentrations (mg/kg) in the dated sediment core B4 of Victoria Harbour

Figure 8. Relationship between $^{208}\text{Pb}/^{207}\text{Pb}$ and $^{206}\text{Pb}/^{207}\text{Pb}$ in the sediments of Victoria Harbour and those from known sources in Hong Kong




Figure 1. The sampling locations, B1–B5, in Victoria Harbour




Figure 2. Physical parameters of the water (water depth, salinity, and pH) and the total organic carbon (TOC) of surface sediments in Victoria Harbour at different sampling periods




Figure 3. The correlation of Cu, Zn, and Pb with the percentage of fine-grained particles in Victoria Harbour




Figure 4. Results of ^{210}Pb dating in sediment cores B2, B4, and B5 of Victoria Harbour




Figure 5. Temporal distribution of Cu, Zn, and Pb in core sediments, B1 to B5, of Victoria Harbour (Core B4 with the estimated date)




Figure 6. The correlation between Cu and Zn in the dated sediment core B4




Figure 7. $^{206}\text{Pb}/^{207}\text{Pb}$ ratios and Pb concentrations (mg/kg) in the dated sediment core B4 of Victoria Harbour




Figure 8. Relationship between $^{208}\text{Pb}/^{207}\text{Pb}$ and $^{206}\text{Pb}/^{207}\text{Pb}$ in the sediments of Victoria Harbour and those from known sources in Hong Kong