



## Correction to: Functional Capacity Evaluation in Different Societal Contexts: Results of a Multicountry Study

Jone Ansuategui Echeita<sup>1</sup> · Matthias Bethge<sup>2</sup> · Berry J. van Holland<sup>3</sup> · Douglas P. Gross<sup>4</sup> · Jan Kool<sup>5</sup> · Peter Oesch<sup>5</sup> · Maurizio A. Trippolini<sup>6,7,8</sup> · Elizabeth Chapman<sup>9</sup> · Andy S. K. Cheng<sup>10</sup> · Robert Sellars<sup>11</sup> · Megan Spavins<sup>12</sup> · Marco Streibelt<sup>13</sup> · Peter van der Wurff<sup>14,15</sup> · Michiel F. Reneman<sup>1</sup>

Published online: 26 June 2018  
© The Author(s) 2018

**Correction to: Journal of Occupational Rehabilitation**  
<https://doi.org/10.1007/s10926-018-9782-x>

The original version of this article unfortunately contained a mistake in Table 2. The data under column head “Left hand-grip strength (n = 336)” was erroneously omitted during the production process. The corrected Table 2 is given below.

The original article has been corrected.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The original article can be found online at <https://doi.org/10.1007/s10926-018-9782-x>.

✉ Jone Ansuategui Echeita  
j.ansuategui.echeita@umcg.nl

<sup>1</sup> Department of Rehabilitation Medicine, University Medical Center Groningen, University of Groningen, P.O. Box 30.002, 9750 RA Haren, Groningen, The Netherlands

<sup>2</sup> Institute of Social Medicine and Epidemiology, University of Lübeck, Lübeck, Germany

<sup>3</sup> Institute for Sports Studies, Hanze University of Applied Sciences, Groningen, The Netherlands

<sup>4</sup> Department of Physical Therapy, University of Alberta, Edmonton, AB, Canada

<sup>5</sup> Rehabilitation Centre Valens, Valens, Switzerland

<sup>6</sup> Center for Disability Research, Liberty Mutual Research Institute for Safety, Hopkinton, Boston, USA

<sup>7</sup> PhD in Rehabilitation Sciences Program, Institute for Health Professions, Massachusetts General Hospital (MGH), Charlestown, Boston, USA

<sup>8</sup> Department of Work Rehabilitation, Rehaklinik Bellikon, Suva Care, Bellikon, Switzerland

<sup>9</sup> Toyota Motor Manufacturing Canada, Puslinch, ON, Canada

<sup>10</sup> Ergonomics and Human Performance Laboratory, Department of Rehabilitation Sciences, The Hong Kong Polytechnic University, Hong Kong, China

<sup>11</sup> FCE Systems Ltd, Geraldine, New Zealand

<sup>12</sup> Occupational Therapy Inc., Randburg, South Africa

<sup>13</sup> Department of Rehabilitation, German Federal Pension Insurance, Berlin, Germany

<sup>14</sup> Research & Development, Military Rehabilitation Center Aardenburg, Doorn, The Netherlands

<sup>15</sup> Institute for Human Movement Studies, HU University of Applied Sciences, Utrecht, The Netherlands

**Table 2** Results from multiple multilevel regression analyses with FCE test performances as dependent variables are given; unstandardized regression coefficients and their standard errors (b(SE)), and model's deviance ( $-2 * \text{LogLikelihood}$ ) by the addition of random and fixed effects are shown

	Floor-to-waist lift <sup>a</sup> (n = 294)		Six minute walk <sup>b</sup> (n = 224)		Right handgrip strength <sup>c</sup> (n = 335)		Left handgrip strength <sup>c</sup> (n = 336)	
	b (SE)	Deviance	b (SE)	Deviance	b (SE)	Deviance	b (SE)	Deviance
<i>Null model</i>		2213.26		2878.28		2847.88		2826.82
Random effects		2193.04***		2867.71**		2806.11***		2787.86***
Clinician and measurement country								
Fixed effects <sup>d</sup>		1941.21***		2643.19***		2590.43***		2597.43***
Intercept	-2.01 (9.86)		219.87 (130.89)		-3.67 (18.00)		-18.96 (17.39)	
Age (years)					-0.16 (0.06)		-0.13 (0.05)	
Sex								
Female	-4.97 (1.01)				-11.33 (1.82)		-11.38 (1.77)	
Height (cm)	0.17 (0.05)		1.98 (0.67)		0.34 (0.10)		0.38 (0.09)	
BMI (kg/m <sup>2</sup> )			-3.82 (1.16)					
Affected body area								
Lower extremity					-0.33 (2.01)			
Upper extremity					-8.19 (2.32)			
Neck					-5.94 (2.02)			
Generalized					-5.62 (1.82)			
Other					9.17 (8.30)			
Observed physical effort								
Heavy	7.74 (1.28)							
Maximum	9.04 (1.70)							
Test ended prematurely								
Yes	-4.87 (1.19)		-168.60 (24.50)					
Post-test HR (bpm)			1.99 (0.30)					
Reported pain intensity (NRS)	-0.64 (0.21)		-10.89 (2.81)		-0.77 (0.33)		-0.91 (0.28)	
Reported effort during FCE test (Borg CR-10)	-0.61 (0.20)		-20.49 (2.79)					
Reported social isolation	-0.34 (0.17)							
Reported catastrophizing					-0.61 (0.27)			
Reported disability (PDI)	-0.07 (0.04)							
Physical work demands (DOT)								
Light					0.65 (2.38)		0.93 (2.31)	
Medium					-0.06 (2.24)		0.78 (2.19)	
Heavy					6.77 (2.51)		6.26 (2.43)	
Very heavy					2.66 (2.86)		5.65 (2.77)	
Days off work								
Less than ¼ year			3.57 (22.30)					
¼–½ year			-59.66 (22.75)					
½–1 year			-12.31 (22.18)					
1–2 years			-27.40 (25.08)					
More than 2 years			-89.19 (24.17)					

Reference category: patient's affected body area: low back; observed physical effort by clinicians: light to moderate; patient's physical work demands: sedentary; patient's days off work: no days off

*BMI* Body Mass Index, *HR* heart-rate, *NRS* Numeric Rating Scale, *PDI* Pain Disability Index, *DOT* Dictionary of Occupational Titles, *FCE* Functional Capacity Evaluation

Significance: \* < 0.05; \*\* < 0.01; \*\*\* < 0.001

<sup>a</sup>Measured in kg

<sup>b</sup>Measured in m

<sup>c</sup>Measured in kgF

<sup>d</sup>Each of the fixed effects factors showed a significant improvement of the model at its addition