



Original Article

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.apjtb.org



doi: 10.4103/2221-1691.314053

Impact Factor: 1.90

Total flavonoids from *Saussurea involucrata* attenuate inflammation in lipopolysaccharide-stimulated RAW264.7 macrophages *via* modulating p65, c-Jun, and IRF3 signaling pathways

Li-Shan Yan¹, Li Wang¹, Brian Chi-Yan Cheng², Yu Ding¹, Jing Kong¹, Qing Gao Wang³, Xiu-Qiong Fu⁴, Shuo-Feng Zhang¹, Gan Luo¹✉, Yi Zhang¹✉

¹School of Chinese Materia Medica, Beijing University of Chinese Medicine, Beijing, China

²College of Professional and Continuing Education, Hong Kong Polytechnic University, Hong Kong, China

³The First Affiliated Hospital, Guangxi University of Chinese Medicine, Guangxi, China

⁴Centre for Cancer and Inflammation Research, School of Chinese Medicine, Hong Kong Baptist University, Hong Kong, China

ABSTRACT

Objective: To investigate the anti-inflammatory effects of the total flavonoids from *Saussurea involucrata* on lipopolysaccharides (LPS)-stimulated murine RAW264.7 macrophages and explore its underlying mechanism of action.

Methods: Total flavonoids from *Saussurea involucrata* were extracted using chromatographic column method. Cell viability was determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide assay. The production of nitric oxide was detected by Griess assay and the release of cytokines (IL-10 and TNF- α) and chemokines (MCP-1, MIP-1 α , and CCL5/RANTES) was determined by ELISA to evaluate the anti-inflammatory activity of total flavonoids from *Saussurea involucrata*. Moreover, nuclear translocation of p65, c-Jun, and IRF3 was detected by immunofluorescence microscopy and Western blotting analysis was performed to determine the expression of related proteins.

Results: Total flavonoids extracted from *Saussurea involucrata* were 751.5 mg/g and the content of rutin was 506.5 mg/g. The production of inflammatory mediators including nitric oxide, cytokines, and chemokines was effectively inhibited by total flavonoids from *Saussurea involucrata*. Meanwhile, total flavonoids also suppressed the nuclear translocation of p65, c-Jun, and IRF3 in LPS-stimulated RAW264.7 cells. The LPS-induced expression of iNOS and COX-2 was remarkably reduced by treatment with total flavonoids from *Saussurea involucrata*. Moreover, total flavonoids decreased the expression levels of p-IKK α / β , p-TBK1, p-p38, p-ERK, p-JNK, p-p65, p-c-Jun, and p-IRF3 in LPS-exposed RAW264.7 macrophages.

Conclusions: Total flavonoids from *Saussurea involucrata*

potentially inhibit the secretion of pro-inflammatory mediators, which may be related to inhibition of p65, c-Jun, and IRF3 signaling pathways in LPS-stimulated RAW264.7 cells.

KEYWORDS: *Saussurea involucrata*; Cytokines; Proinflammatory; Macrophages; Flavonoids

1. Introduction

The immune system is our host defense against various adverse conditions, such as tissue injury from xenobiotic insult[1]. However, over-activated or dysregulated immune responses may conversely result in molecular dysregulation, cellular injury, and tissue damage[2]. In spite of the complexity of immune pathogenesis, molecular evidence has indicated that toll-like receptors (TLRs), especially TLR4, play a key role in various infectious diseases, such as sepsis, plague, asthma, and pneumonia[3,4]. Increased activity

✉To whom correspondence may be addressed. E-mail: yizhang714@163.com (Yi Zhang); luna049@126.com (Gan Luo).

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2021 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow. All rights reserved.

How to cite this article: Yan LS, Wang L, Cheng BC, Ding Y, Kong J, Wang QG, et al. Total flavonoids from *Saussurea involucrata* attenuate inflammation in lipopolysaccharide-stimulated RAW264.7 macrophages *via* modulating p65, c-Jun, and IRF3 signaling pathways. Asian Pac J Trop Biomed 2021; 11(6): 273-284.

Article history: Received 2 July 2020; Revision 4 August 2020; Accepted 24 December 2020; Available online 10 May 2021

of TLR4 has been detected in many types of immune cells[4], such as macrophages[5]. Upon ligand binding to external stimuli [e.g. lipopolysaccharide (LPS)], TLR4 interacts with its co-receptor MD2 and adaptors, and subsequently initiates MyD88-dependent and -independent signaling[6]. The subsequent kinase cascades result in the activation of transcription factors, such as nuclear factor κ B (NF- κ B), activator protein 1 (AP-1), and interferon regulatory factor 3 (IRF3), leading to increased production of inflammatory mediators which trigger immune responses. Therefore, blocking NF- κ B, AP-1, and IRF3 signaling pathways emerges as an attractive therapeutic approach for inhibiting inflammatory actions[7].

Because of the chronic nature of some inflammatory diseases, patients may need long-term treatment using drugs like nonsteroidal anti-inflammatory drugs, which may cause serious adverse effects[8]. Seeking a safe and effective anti-inflammatory agent has been attracting intensive interest of researchers. Herbal medicines may provide an ideal source of safe and effective agents for prevention and/or treatment of inflammatory diseases[9]. However, one of the most important obstacles for Chinese medicinal herbs to be included in mainstream medicine is unclear active constituents and mechanism of action. *Saussurea involucrata* (*S. involucrata*) (Xue Lian Hua in Chinese) is a traditional Chinese herbal medicine and has been widely used for the treatment of rheumatoid arthritis, stomachache, and dysmenorrhea for hundreds of years in China[10]. It has been reported that the extracts of *S. involucrata* possessed a wide array of pharmacological actions, including anti-inflammatory, anti-hypoxic, and anti-fatigue activities[11–13]. The major constituents isolated from *S. involucrata* include flavonoids, amino acids, polysaccharides, and sesquiterpene lactone[14–18]. Recent studies reported that as natural polyphenolic compounds, flavonoids could inhibit regulatory enzymes and transcription factors involved in inflammatory actions[19]. Thus, we hypothesized that flavonoids from *S. involucrata* might possess anti-inflammatory properties. To verify our hypothesis, we isolated the total flavonoids from *S. involucrata* and evaluated the anti-inflammatory effects and the underlying mechanisms of total flavonoids from *S. involucrata* in LPS-stimulated RAW264.7 cells.

2. Materials and methods

2.1. Chemicals, reagents, and antibodies

Chlorogenic acid (batch number: Y24J7K16726) and rutin (batch number: Y16M9S61523) were purchased from Shanghai Yuanye Biotechnology Co. Ltd. (Shanghai, China). The purity of each standard was higher than 98%. AB-8 macroporous resin was obtained from the chemical plant of Nankai University (Tianjin, China). Polyamide resin was bought from Sinopharm Chemical Reagent Co. Ltd. (Shanghai, China). 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl

tetrazolium bromide (MTT), LPS (*Escherichia coli* 055:B5), and modified Griess reagent were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Fetal bovine serum was bought from Biological Industries (Beth-Haemek, Israel). Dulbecco's Modified Eagle Medium was obtained from Corning Cellgro (Manassas, VA, USA). Penicillin-streptomycin solution was bought from Caisson labs (Smithfield, UT, USA). Tumor necrosis factor- α (TNF- α), interleukin 10 (IL-10), macrophage inflammatory protein-1 α (MIP-1 α), monocyte chemoattractant protein-1 (MCP-1), and RANTES ELISA kits were purchased from Thermo Fisher Scientific (San Diego, CA). Prostaglandin E₂ (PGE₂) ELISA kit was provided by Enzo Life Sciences (Exeter, UK). I κ B kinase α (IKK α) and SP1 monoclonal antibodies were supplied by Santa Cruz Biotechnology (Santa Cruz, CA, USA). IRF3 and anti-mouse IgG HRP linked antibody were purchased from Abcam (Cambridge, UK). IRF3 (Ser 396) was provided by ABNOVA (Taiwan, China). Phospho (p)-NF- κ B/p65 (Ser536), NF- κ B/p65, p-c-Jun (Ser73), c-Jun, cyclooxygenase-2 (COX-2), inducible nitric oxide synthase (iNOS), p-IRF3 (Ser396), p-I κ B α (Ser32), I κ B α , p-IKK α / β (Ser176/177), extracellular-signal-regulated kinases (ERK), p-ERK (Thr202/Tyr204), c-Jun N-terminal kinase (JNK), p-JNK (Thr183/Tyr185), p-p38 (Thr180/Tyr182), p38, TANK-binding kinase 1 (TBK1), p-TBK1 (Ser172), anti-rabbit IgG HRP linked antibody, and Alexa Fluor 488-conjugated secondary antibody were bought from Cell Signaling Technology (Boston, MA, USA).

2.2. Preparation of ethanol extract of *S. involucrata*

S. involucrata was bought from local market of Xinjiang and authenticated by Lecturer Guang Xi-ren (School of Chinese Materia Medica, Beijing University of Chinese Medicine). Voucher specimens were deposited at the Department of Pharmacology, School of Chinese Medicine, Beijing University of Chinese Medicine. *S. involucrata* was cut into small pieces. Then the pieces (70 g) were homogenized with 700 mL of 70% ethanol (v:v) at room temperature. Subsequently, heating under reflux was performed 3 times and 1 h for each at room temperature. The extraction solvent was slightly boiled across the whole extraction process and the boiling point of the solvent was 80.2 °C. Then combined extracting solution was filtrated, and vacuum-evaporated to remove the solvent. Finally, ethanol extract of *S. involucrata* (17.82 g) was obtained.

2.3. Preparation of flavonoids of *S. involucrata* powder

A chromatographic column (3.5 cm \times 100 cm) wet-packed with AB-8 macroporous resin (chemical plant of Nankai University, Tianjin, China) was used to carry out the enrichment experiment. The ratio of diameter to height was 1:8 and the bed volume (BV) was 270 mL. Briefly, 16 grams of ethanol extract of *S. involucrata* were dissolved by 50 mL of water and the mixture was then loaded on the

chromatographic column. The flow rate was 2 BV/h. Next, 4 BV of water was used to remove the aqueous impurities. The column was then eluted with 6 BV 15% and 40% methanol aqueous solution, respectively. The 40% ethanol eluate was collected and vacuum dried at 60 °C to afford a residue (1.3 g flavonoid enrichment of *S. involucrata*).

The purification was conducted on a glass column (2.6 cm × 50 cm) wet-packed with polyamide resin (Sinopharm Chemical Reagent Co., Ltd, Shanghai, China). The ratio of diameter to height was 1:8 and the BV was 110 mL. One gram of flavonoid enrichment was dissolved in 50 mL of water and loaded onto the column with a flow rate of 2 BV/h. Next, 4 BV of water and 4 BV of 20% ethanol were used to remove impurities. The column was then eluted with 6 BV 50% ethanol. The eluates were collected, merged, and evaporated under reduced pressure using a rotator evaporator and vacuum dried at 60 °C. Finally, 305 mg of flavonoid of *S. involucrata* was obtained. For ultra-performance liquid chromatography (UPLC) analysis, ethanol extract and flavonoid enrichment were diluted by 50% methanol-water solution, and flavonoids of *S. involucrata* were diluted by 70% methanol. All samples were filtered through a millipore membrane filter with an average pore diameter of 0.22 µm, and each sample (5 µL) was injected for UPLC analysis.

2.4. Characterization of the ethanol extract, flavonoid enrichment, and flavonoids of *S. involucrata*

To control the quality of flavonoids of *S. involucrata*, UPLC analysis was conducted using a Waters Acquity UPLC system (Waters Corporation, Milford, MA, USA), consisting of a binary solvent delivery pump, an autosampler, and a photodiode array detector. The chromatographic separation was performed by an Acquity UPLC® BEH C₁₈ column (1.7 µm, 100 mm × 2.1 mm) with a Van Guard™ Pre-Column (1.7 µm, 5 mm × 2.1 mm). The gradient mobile phase consisted of solvent A (acetonitrile) and solvent B (0.2% formic acid in water). The UPLC elution profile was as follows: 0–6 min, 5%–9% A; 6–12 min, 9%–13% A; 12–19 min, 13%–16% A; 19–28 min, 16%–17% A; 28–40 min, 17%–28% A. The flow rate was 0.3 mL/min and the column temperature was set at 40 °C. The chromatograms were monitored with the PDA detector at a wavelength of 340 nm to detect chlorogenic acid and rutin.

2.5. Detection of total flavonoids content

The total flavonoid contents in three samples were determined according to a previous study with some modifications[20]. Each mixture containing 0.2 mL of 5% (w/v) NaNO₂ and 0.1 mL of test compound was incubated for 6 min at 22 °C before adding 0.2 mL of 10% (w/v) Al(NO₃)₃. After additional 6 min incubation, 2 mL of 4% NaOH and 2.7 mL of 70% methanol were added and the absorbance was read at 505 nm. The experiment was performed in

three replicates.

2.6. Cell lines and cell culture

The RAW264.7 cell line was obtained from American Type Culture Collection (Manassas, VA, USA). Cells were cultured in Dulbecco's Modified Eagle Medium supplemented with penicillin/streptomycin antibiotics (1%) and heat-inactivated fetal bovine serum (10%) in a 5% CO₂ incubator at 37 °C. Flavonoids of *S. involucrata* were dissolved in dimethyl sulfoxide to prepare a stock solution and then freshly diluted with cell culture medium to various concentrations.

2.7. MTT assay

RAW264.7 cells were seeded (6×10^3 cells/well) into 96-well plates. Cells were treated with 1 µg/mL lipopolysaccharides (LPS) in the presence or absence of various concentrations of flavonoids of *S. involucrata* (6.25–400 µg/mL) for 24 h at 37 °C. Then, cells were incubated with 10 µL of the MTT solution (5 mg/mL) for 3 h at 37 °C. The medium was carefully aspirated and MTT formazan precipitate was dissolved in 100 µL dimethyl sulfoxide. Then the absorbance was measured at 570 nm using a microplate spectrophotometer (BMG SPECTROstar Nano, Germany). Cell viability in the control group was set as 100%. Six replicates of experimental data were used.

2.8. Griess assay

As nitric oxide (NO) plays important roles in LPS-induced inflammatory responses and is implicated in the pathogenesis of various inflammatory conditions[21,22], we determined the content of NO in the culture medium after LPS treatment by Griess assay. Briefly, RAW264.7 cells were seeded at 2×10^5 cells/mL on 24-well culture plates for 24 h. Cells were treated with 1 µg/mL LPS in the presence or absence of various concentrations of flavonoids of *S. involucrata* (25–200 µg/mL) for 24 h at 37 °C. NO production was determined by measuring the accumulated nitrite (a stable degradation product of NO) in the culture medium with Griess reagent. Absorbance at 540 nm was measured with a NaNO₂ standard curve. NO production was then determined.

2.9. ELISA assays

Macrophages activated by endotoxin initiate the release of pro-inflammatory cytokines and chemokines in the early stage of inflammatory response[5,23]. Hence, we detected several inflammatory cytokines and chemokines in the culture medium after LPS exposure using the corresponding ELISA kits in this study. RAW264.7 cells were seeded (2×10^5 cells/well) into 24-well plates. Cells were treated with 1 µg/mL LPS in the presence or absence of various concentrations of flavonoids of *S. involucrata* (25–

200 µg/mL) for 24 h at 37 °C. Cell-free supernatants were obtained for the determination of PGE₂, TNF-α, IL-10, MCP-1, MIP-1α, and RANTES using ELISA kits according to the manufacturer's instructions. Four replicates of experimental data were used.

2.10. Immunofluorescence staining

The nuclear localization of p65, c-Jun, and IRF3 was detected with Nikon A1R Eclipse Ti confocal microscope (Nikon Corp., Tokyo, Japan) as previously described[24]. Briefly, cells were seeded into chamber slides for 12 h and then treated with flavonoids of *S. involucrata* (100 and 200 µg/mL) for 1 h. After LPS (1 µg/mL) stimulation for 1 h, cells were fixed by 4% paraformaldehyde for 10 min and then permeabilized with Triton X-100 (0.25%) for 30 min at 37 °C. Cells were then blocked for 1 h with 2% bovine serum albumin and incubated with target antibodies including p65, c-Jun, and IRF3 diluted in cold PBS containing 2% bovine serum albumin at 4 °C overnight. Then, the cells were incubated with FITC-labeled anti-rabbit secondary antibody (Cell Signaling Technology, catalog no. 4412; 1:500 dilution) in dark under room temperature for 1 h. After incubation, cells were washed 3 times by cold PBS. 4',6-diamidino-2-phenylindole (DAPI, YESEN, Shanghai, China) was used to stain nucleus right before performing assay.

2.11. Cytosolic and nuclear protein separation

RAW264.7 cells were seeded (5×10^5 cells/well) into 60 mm diameter culture dishes and cultured for 24 h. Cells were pre-treated with flavonoids of *S. involucrata* (100 and 200 µg/mL) for 1 h and LPS for additional 30 min. Cells were collected and nuclear extraction kit (Solarbio, Beijing, China) was used to separate nuclear and cytosolic protein according to the manufacturer's instructions. After that, samples were dissolved in lysis buffer for Western blotting assay.

2.12. Immunoblotting

The quantified cell lysates were assessed by Western blotting analysis as described previously[25]. Briefly, cells (5×10^5) were seeded into 60-mm culture dishes and then treated with flavonoids of *S. involucrata* at concentrations of 100 and 200 µg/mL. LPS (1 µg/mL) was added 1 h after pre-treatment with flavonoids of *S. involucrata* and the cells were incubated at 37 °C for 30 min. Subsequently, cells were collected and lysed with RIPA buffer (Beyotime Biotechnology, Beijing, China). The supernatant protein from each sample was heated with 4 × sodium dodecyl sulfate (SDS) sample buffer at 95 °C for 8 min. An aliquot of 30 µg of protein was subjected to 10% SDS-PAGE and electro-transferred onto polyvinylidene fluoride membrane. After blocking with 5% non-fat milk in TBST, the membrane was incubated with indicated primary

antibodies overnight at 4 °C. Antibodies against the following molecules were used: p65 (Cell Signaling Technology, catalog no. 8242; 1:1 000), p-p65 (Ser536; Cell Signaling Technology, catalog no. 3033; 1:1 000), c-Jun (Cell Signaling Technology, catalog no. 9165; 1:1 000), p-c-Jun (Ser73; Cell Signaling Technology, catalog no. 9164; 1:1 000), COX-2 (Cell Signaling Technology, catalog no. 12282; 1:1 000), iNOS (Cell Signaling Technology, catalog no. 13120; 1:1 000), p-IRF3 (Ser396; Abnova, catalog no. PAB31634; 1:1 000), IκBα (Cell Signaling Technology, catalog no. 4814; 1:1 000), p-IκBα (Ser32; Cell Signaling Technology, catalog no. 2859; 1:1 000), p-IKKα/β (Ser176/177; Cell Signaling Technology, catalog no. 2078; 1:1 000), ERK (Cell Signaling Technology, catalog no. 4695; 1:1 000), p-ERK (Thr202/Tyr204; Cell Signaling Technology, catalog no. 4370; 1:1 000), JNK (Cell Signaling Technology, catalog no. 9252; 1:1 000), p-JNK (Thr183/Tyr185; Cell Signaling Technology, catalog no. 4668; 1:1 000), p38 (Cell Signaling Technology, catalog no. 8690; 1:1 000), p-p38 (Thr180/Tyr182; Cell Signaling Technology, catalog no. 4511; 1:1 000), TBK1 (Cell Signaling Technology, catalog no. 3013; 1:1 000), p-TBK1 (Ser172; Cell Signaling Technology, catalog no. 5483; 1:1 000), IRF3 (Abcam, catalog no. ab68481; 1:1 000), IKKα (Santa Cruz, catalog no. sc-7607; 1:500), β-actin (Cell Signaling Technology, catalog no. 4970; 1:1 000) and SP1 (Santa Cruz, catalog no. sc-420; 1:500) were used as internal control. The membranes were washed three times and incubated with the anti-rabbit IgG secondary antibody (Cell Signaling Technology, catalog no. 7074; 1:2 000) or anti-mouse IgG HRP-linked antibody (Abcam, catalog no. ab6789; 1:2 000) for 1 h. Visualization was performed using Tanon 5200 Multi chemiluminescent imaging system (Tanon Science & Technology Co., Ltd., Shanghai, China) with an enhanced-chemiluminescence substrate, and the blots were analyzed using Image J software [National Institutes of Health (NIH), Bethesda, MD, USA]. Protein levels were corrected with values determined on β-actin blots. Three replicates were used.

2.13. Statistical analysis

All values were expressed as mean ± standard deviation (SD). Data were analyzed by one-way ANOVA using SPSS (version 16.0) statistical analysis program followed by Dunnett's multiple comparisons. $P < 0.05$ was considered statistically significant.

3. Results

3.1. Characterization of ethanol extract, flavonoid enrichment, and flavonoids of *S. involucrata* powders

UPLC chromatogram showed that the contents of chlorogenic acid and rutin were different among the ethanol extract, flavonoid




Figure 1. UPLC chromatograms of chlorogenic acid (peak 1) and rutin (peak 2) in different samples of *Saussurea involucrata* (*S. involucrata*). (A) Chlorogenic acid and rutin; (B) ethanol extract of *S. involucrata*; (C) flavonoid enrichment of *S. involucrata*; (D) flavonoids of *S. involucrata*.

enrichment, and flavonoids of *S. involucrata* powders (Figure 1). The mean contents of chlorogenic acid/rutin in these three samples were 15.4/10.7, 3.2/136.7, and 0/506.5 mg/g, respectively. Moreover, the contents of total flavonoids in the three extracts were 130.7, 646.9, and 751.5 mg/g, respectively.

3.2. Effect of flavonoids of *S. involucrata* on cell viability

Compared with the control group, the cell viability of RAW264.7 cells did not change when up to 400 µg/mL of flavonoids of *S. involucrata* was added with LPS for 24 h (Figure 2). We therefore chose the sub-toxic concentrations of flavonoids of *S. involucrata* (25–200 µg/mL) in the following experiments.

3.3. Effects of flavonoids of *S. involucrata* on the production of NO and PGE₂ and the expression of iNOS and COX-2

Figures 3A and 3B show that the production of NO and PGE₂ was significantly increased in LPS-stimulated RAW264.7 cells. Flavonoids of *S. involucrata* (25–200 µg/mL) concentration-

dependently suppressed the release of NO and PGE₂. In addition, the expressions of iNOS and COX-2 were also suppressed by flavonoids of *S. involucrata* in a concentration-dependent manner (Figure 3C).




Figure 2. Effect of flavonoids of *S. involucrata* (FSI) on cell viability of LPS-stimulated RAW264.7 macrophages. Cells were treated with FSI (0–400 µg/mL) in the presence of lipopolysaccharides (LPS) (1 µg/mL) for 24 h. The cell viability was compared with cells without FSI and LPS stimulation using MTT assay. Data are expressed as mean ± SD of six independent experiments.

3.4. Effects of flavonoids of *S. involucrata* on the production of TNF- α , IL-10, MCP-1, MIP-1 α and RANTES

As shown in Figures 4A and 4E, the release of inflammatory

cytokines, including TNF- α and IL-10 was obviously elevated after LPS stimulation in RAW264.7 cells. Flavonoids of *S. involucrata* inhibited the secretion of these inflammatory cytokines in a concentration-dependent manner. The same trend was observed




Figure 3. Effect of FSI on the production of NO and PGE₂ and the expression of iNOS and COX-2 in LPS-stimulated RAW264.7 macrophages. The production and expression of NO (A) and PGE₂ (B) and enzymes (C) were detected by Griess assays, ELISA, and Western blotting, respectively. Data are expressed as mean \pm SD. Four replicates were used in ELISA assays and three independent experiments in Western blotting. ** $P < 0.01$ vs. the control group; # $P < 0.05$, ## $P < 0.01$ vs. the LPS group. Statistical analysis was performed using one-way ANOVA followed by Dunnett's multiple comparison test. NO: nitric oxide, PGE₂: prostaglandin E₂, iNOS: inducible nitric oxide synthase, COX-2: cyclooxygenase-2.




Figure 4. Effect of FSI on the production of pro-inflammatory cytokines and chemokines in LPS-stimulated RAW264.7 macrophages. The production of cytokines (A and E) and chemokines (B-D) in the culture medium were detected by ELISA assay. Data are expressed as mean \pm SD of four independent experiments. ** $P < 0.01$ vs. the control group; # $P < 0.05$, ## $P < 0.01$ vs. the LPS group. Statistical analysis was performed using one-way ANOVA followed by Dunnett's multiple comparison test. CCL5/RANTES: C-C motif chemokine ligand 5, TNF- α : tumor necrosis factor-alpha, IL-10: interleukin 10, MIP-1 α : macrophage inflammatory protein-1 α , MCP-1: monocyte chemoattractant protein-1.





Figure 5. Effect of FSI on the nuclear localization of p65, c-Jun, and IRF3. The nuclear localization of p65, c-Jun, and IRF3 was detected by immunofluorescence analyses. Bar scale=33 µm.

A



B




Figure 6. Effect of FSI on the expression of p65, c-Jun, and IRF3. The expression levels of nuclear (A) and cytoplasmic (B) proteins of p65, c-Jun, and IRF3 were detected by Western blotting. The nuclear proteins were normalized against SP1 levels, and the cytoplasmic proteins were normalized for β -actin levels. Data are expressed as mean \pm SD of three independent experiments. ** $P < 0.01$ vs. the control group; # $P < 0.05$, ## $P < 0.01$ vs. the LPS group. Statistical analysis was performed using one-way ANOVA followed by Dunnett's multiple comparison test. SP1: specificity protein 1.

in the production of MCP-1, MIP-1 α , and CCL5/RANTES after treatment with flavonoids of *S. involucrata* in LPS stimulated RAW264.7 cells (Figures 4B-D).

3.5. Effect of flavonoids of *S. involucrata* on the nuclear localization of p65, c-Jun, and IRF3

Immunofluorescence was employed to detect the nuclear localization of p65, c-Jun, and IRF3. Figure 5 shows that the LPS exposure promoted the translocation of these transcriptional factors from the cytoplasm to the nucleus in RAW264.7 macrophages. Flavonoids of *S. involucrata* remarkably reduced the nuclear localization of p65, c-Jun, and IRF3, demonstrating potential ability to regulate transcription factor activity in inflammatory response.

3.6. Effects of flavonoids of *S. involucrata* on the expression of p65, c-Jun, and IRF3

We further confirmed the nuclear translocation of these key transcriptional factors using Western blotting. As shown in Figure 6, exposure of RAW264.7 cells to LPS for 30 min significantly elevated the nuclear protein levels of p65, c-Jun, and IRF3. Flavonoids of *S. involucrata* concentration-dependently decreased the nuclear protein levels of these three transcriptional factors. Furthermore, the cytoplasmic protein levels of p65, c-Jun, and IRF3 were decreased in LPS-stimulated cells, whereas no significant changes in cytoplasmic proteins of these transcriptional factors were observed after treatment with flavonoids of *S. involucrata*. Consistent with the data from Figure 5, flavonoids of *S. involucrata* inhibited p65, c-Jun, and IRF3

nuclear translocation, thus reducing their target gene expression.

3.7. Effect of flavonoids of *S. involucrata* on the p65, c-Jun, and IRF3 related signaling pathways

To elucidate the underlying mechanism of the inhibitory effect of flavonoids of *S. involucrata* on the nuclear translocation of these three transcriptional factors, we tested the effect of flavonoids of *S. involucrata* on the related molecular signaling pathways. As shown in the Figure 7, the expression of phosphorylated p65, c-Jun, and IRF3 was significantly elevated after exposure to LPS. Flavonoids of *S. involucrata* suppressed the phosphorylation of p65, c-Jun, and IRF3 in a concentration-dependent manner. Moreover, the phosphorylated MAPKs, including p38, ERK, and JNK were markedly increased after LPS stimulation and these effects were prevented by flavonoids of *S. involucrata* in a concentration-dependent manner. Meanwhile, the phosphorylation of I κ B α , IKK α / β , and TBK1 was significantly upregulated by LPS treatment, which was downregulated by flavonoids of *S. involucrata* (Figure 7). Taken together, these data showed that IKK α / β /NF- κ B, MAPKs/AP-1, and TBK1/IRF3 signaling pathways were involved in the inhibitory effect of flavonoids of *S. involucrata* on the nuclear translocation of p65, c-Jun, and IRF3, thereby reducing the secretion of inflammatory mediators.

4. Discussion

In recent years, traditional Chinese medicine has increased in




Figure 7. Effect of FSI on p65, c-Jun, and IRF3 signaling pathways. (A) Representative bands of proteins; (B) Quantification results. The expression levels of phosphorylated and total IκBα, IKKα/β, TBK1, p38, ERK, JNK, p65, c-Jun, and IRF3 were determined by Western blotting. Data are expressed as mean ± SD of three independent experiments. * $P < 0.05$, ** $P < 0.01$ vs. the control group; # $P < 0.05$, ## $P < 0.01$ vs. the LPS group. Statistical analysis was performed using one-way ANOVA followed by Dunnett's multiple comparison test. TBK1: TANK-binding kinase 1, IRF3: interferon regulatory factor 3, c-JNK: c-Jun N-terminal kinase, ERK: extracellular signal-regulated kinase, p38: p38 mitogen-activated protein kinase, IKKα/β: IκB kinase α/β.

popularity, especially for preventing and curing inflammatory diseases[26,27]. Flavonoids are a large group of polyphenolic compounds found in almost all natural supplements and exhibit strong anti-inflammatory properties *via* modulating a broad range of inflammation-associated signaling pathways[28]. Among these flavonoids, rutin is one of the most popular phenolic compounds known to have a variety of biological activities, including anti-allergic, anti-inflammatory, anti-proliferative, and anti-carcinogenic effects[29]. In our present study, UPLC analysis was performed and rutin was used as a marker to estimate the contents of flavonoid compounds in the ethanol extract, flavonoid enrichment, and flavonoids of *S. involucrata*. The results showed that the contents of rutin and chlorogenic acid (a marker of phenolic acid) were progressively increased and decreased after purification, respectively, indicating the total flavonoid was successfully separated from *S. involucrata*. Given that rutin has been reported as a good anti-inflammatory agent, the anti-inflammatory effect of flavonoids of *S. involucrata* may be related to the presence of rutin. Further studies will be conducted to verify the role of rutin in the anti-inflammatory effects of flavonoids of *S. involucrata* and identify other active components (*e.g.* phenolic acid) responsible for the action.

During the initial phase of the inflammatory response, NO and PGE₂ act as vasodilators to facilitate the influx of neutrophils, macrophages, and mast cells from the bloodstream, leading to swelling and edema at the site of infection or tissue injury[30,31]. It has been implicated that iNOS and COX-2 are the key enzymes of NO and PGE₂, respectively[31,32]. Elevated activities of iNOS and COX-2 promote the production of NO and PGE₂, thereby amplifying inflammatory responses and even contributing to tissue destruction[2]. Our current research showed that flavonoids of *S. involucrata* suppressed the production of NO and PGE₂ as well as their synthesizing enzymes. We also observed that the inhibitory concentrations (25–200 µg/mL) of flavonoids of *S. involucrata* against these inflammatory mediators did not cause cytotoxicity. These results suggested that flavonoids of *S. involucrata* inhibited the NO and PGE₂ production *via* decreasing the activities of iNOS and COX-2 in LPS-stimulated RAW264.7 cells.

Cellular response to LPS is initiated by the interaction of LPS and its receptor complex composed of LBP, CD14, MD2, and TLR4[33]. This activates their down-stream proteins IKK complex, which contains two catalytic subunits (IKK α and IKK β), triggering the degradation of I κ B α by phosphorylation[34]. Subsequently, NF- κ B is phosphorylated and translocates into the nucleus, promoting transcription of pro-inflammatory mediators such as NO, PGE₂, TNF- α , and IL-10[35]. According to previous studies, the extracts of *S. involucrata* ameliorated infiltration of inflammatory cells and reduced the elevation of TNF- α in synovial tissue as well as serum TNF- α , IL-1 β , and IL-6 in arthritic rats[36,37]. Moreover, *S. involucrata* and its main component rutin could decrease the

expression of COX-2 *via* downregulation of NF- κ B activity[38]. Consistent with these findings, in our present study, we found that the production of TNF- α and IL-10 were suppressed by flavonoids of *S. involucrata* in LPS-stimulated RAW264.7 cells. We also observed that the phosphorylation of the upstream intermediary proteins of these molecules, including p65, I κ B α , and IKK α / β was inhibited by flavonoids of *S. involucrata*, revealing that flavonoids of *S. involucrata* effectively inhibited the p65 signaling pathway. Besides, AP-1 and IRF3 also play an important role in the regulation of inflammation[33,39]. Once AP-1 is activated, it translocates into the nucleus and binds to the target DNA, triggering the secretion of pro-inflammatory chemokines, such as MCP-1 and MIP-1 α . This attracts inflammatory cells to enter the tissue during inflammatory response[40]. MAPKs (p38, ERK, and JNK) are serine-threonine kinases that regulate the transcriptional activation of AP-1 and its DNA-binding activity which mediates transcription of a wide array of AP-1 target genes[41]. We found that the production of MCP-1 and MIP-1 α was reduced after treatment with flavonoids of *S. involucrata*. The nuclear-phosphorylated protein levels of the main component of AP-1 (c-Jun) were also inhibited by flavonoids of *S. involucrata*. It also reduced the phosphorylated upstream proteins including p38, ERK, and JNK. That may attribute to the presence of rutin which has been implied as a potent inhibitor of p38 and JNK activity[42]. These results indicated that flavonoids of *S. involucrata* suppressed the MAPKs/AP-1 signaling transduction. In addition, in a TRIF-dependent manner, activated TLR4 promotes the binding of TRIF to TBK1/IKK ϵ , which then initiates the cascade of activation of IRF3 and facilitates the nuclear translocation of IRF3, thereby increasing the production of chemokines, such as RANTES[43]. Our study revealed that the release of RANTES was increased after LPS exposure and decreased by flavonoids of *S. involucrata* in RAW264.7 cells, showing that IRF3 signaling might also be a target of flavonoids of *S. involucrata*. Therefore, the expression levels of the upstream proteins were detected. The results showed that the nuclear translocation of IRF3 was inhibited in LPS-stimulated RAW264.7 cells after treatment with flavonoids of *S. involucrata*. The phosphorylation of IRF3 and TBK1 was also suppressed. These data suggested that inhibition of IRF3 signaling may also contribute to the effects of flavonoids of *S. involucrata* on adaptive immune responses.

In summary, the current study showed that flavonoids of *S. involucrata* downregulated the LPS-induced activities of pro-inflammatory mediators in RAW264.7 cells. Our work also proved that flavonoids of *S. involucrata* suppressed the nuclear translocation of NF- κ B/p65, AP-1/c-Jun, and IRF3 in LPS-treated RAW264.7 cells. The underlying mechanism may be related to inhibition of the related signaling pathways. However, lack of *in vivo* data is a major limitation of this study. Our further investigation will employ inflammatory animal models, such as mouse model for arthritis,

to confirm the anti-inflammatory properties of flavonoids of *S. involucrata*. Taken together, the results may provide a chemical and pharmacological justification for the use of flavonoids of *S. involucrata* in the treatment of patients with inflammatory diseases.

Conflict of interest statement

The authors declare that they have no competing interests.

Funding

This work was supported by the National Natural Science Foundation of China (Grant Number 81803793 and 82003957), the Young Scientist Program by Beijing University of Chinese Medicine, and the Fundamental Research Funds for the Central Universities (Grant Number 2018-JYBZZ-XJSJJ008).

Authors' contributions

YZ contributed to the conceptualization, funding acquisition and project administration. YZ, LSY, LW and YD conducted the biological investigation. GL and JK performed UPLC analysis. GL, LSY, BCC, QGW and SFZ analyzed data. YZ and XQF performed data visualization. YZ and GL drafted the original manuscript. BCC revised the manuscript.

References

- [1] Medzhitov R. Origin and physiological roles of inflammation. *Nature* 2008; **454**(7203): 428-435.
- [2] Fullerton JN, Gilroy DW. Resolution of inflammation: A new therapeutic frontier. *Nat Rev Drug Discov* 2016; **15**(8): 551-567.
- [3] Zhang Y, Guo H, Cheng BC, Su T, Fu XQ, Li T, et al. Dingchuan tang essential oil inhibits the production of inflammatory mediators via suppressing the IRAK/NF- κ B, IRAK/AP-1, and TBK1/IRF3 pathways in lipopolysaccharide-stimulated RAW264.7 cells. *Drug Des Devel Ther* 2018; **12**: 2731-2748.
- [4] Miller SI, Ernst RK, Bader MW. LPS, TLR4 and infectious disease diversity. *Nat Rev Microbiol* 2005; **3**(1): 36-46.
- [5] Hamidzadeh K, Christensen SM, Dalby E, Chandrasekaran P, Mosser DM. Macrophages and the recovery from acute and chronic inflammation. *Annu Rev Physiol* 2017; **79**: 567-592.
- [6] O'Neill LAJ, Bowie AG. The family of five: TIR-domain-containing adaptors in Toll-like receptor signalling. *Nat Rev Immunol* 2007; **7**(5): 353-364.
- [7] Roy A, Srivastava M, Saqib U, Liu D, Faisal SM, Sugathan S, et al. Potential therapeutic targets for inflammation in toll-like receptor 4 (TLR4)-mediated signaling pathways. *Int Immunopharmacol* 2016; **40**: 79-89.
- [8] Perkins JR, Sanak M, Canto G, Blanca M, Cornejogarcia JA. Unravelling adverse reactions to NSAIDs using systems biology. *Trends Pharmacol Sci* 2015; **36**(3): 172-180.
- [9] Schumacher M, Juncker T, Schnekenburger M, Gaascht F, Diederich M. Natural compounds as inflammation inhibitors. *Genes Nutr* 2011; **6**(2): 89-92.
- [10] Gong G, Huang J, Yang Y, Qi B, Han G, Zheng Y, et al. *Saussurea involucrata* Herba (snow lotus): Review of chemical compositions and pharmacological properties. *Front Pharmacol* 2020; **10**: 1549.
- [11] Ma HP, Fan PC, Jing LL, Yao J, He XR, Yang Y, et al. Anti-hypoxic activity at simulated high altitude was isolated in petroleum ether extract of *Saussurea involucrata*. *J Ethnopharmacol* 2011; **137**(3): 1510-1515.
- [12] Yi T, Zhao ZZ, Yu ZL, Chen HB. Comparison of the anti-inflammatory and anti-nociceptive effects of three medicinal plants known as "Snow Lotus" herb in traditional Uighur and Tibetan medicines. *J Ethnopharmacol* 2010; **128**(2): 405-411.
- [13] Chik WI, Zhu L, Fan LL, Yi T, Zhu GY, Gou XJ, et al. *Saussurea involucrata*: A review of the botany, phytochemistry and ethnopharmacology of a rare traditional herbal medicine. *J Ethnopharmacol* 2015; **172**: 44-60.
- [14] Li GL, Sun ZW, Song CH, Xia L, Zheng J, Suo YR, et al. A sensitive fluorescence reagent, 2-(2-(7H-dibenzo(a,g)carbazol-7-yl)-ethoxy) ethyl chloroformate, for amino acids determination in *Saussurea involucrata* and *Artemisia capillaris* Thunb using high-performance liquid chromatography with fluorescence detection and identification with mass spectroscopy/electrospray ionization source. *Biomed Chromatogr* 2011; **25**(6): 689-696.
- [15] Wang XL, Gesang SL, Jiao W, Liao X, Ding LS. Two new sesquiterpenoid glucosides from the aerial parts of *Saussurea involucrata*. *J Integr Plant Biol* 2007; **49**(5): 609-614.
- [16] Xiao W, Li X, Li N, Bolati M, Wang XJ, Jia XG, et al. Sesquiterpene lactones from *Saussurea involucrata*. *Fitoterapia* 2011; **82**(7): 983-987.
- [17] Chen H, Zhao ZZ, Jiang ZH, Cai S, Wang TM. Identification and determination of the major constituents in the traditional uighur medicinal plant *Saussurea involucrata* by LC-DAD-MS. *Chromatographia* 2009; **69**: 537-542.
- [18] Chen QL, Zhu L, Tang YN, Kwan HY, Zhao ZZ, Chen HB, et al. Comparative evaluation of chemical profiles of three representative 'snow lotus' herbs by UPLC-DAD-QTOF-MS combined with principal component and hierarchical cluster analyses. *Drug Test Anal* 2017; **9**(8): 1105-1115.
- [19] Rengasamy KRR, Khan H, Gowrishankar S, Lagoa R, Mahomoodally F, Khan Z, et al. The role of flavonoids in autoimmune diseases:

- Therapeutic updates. *Pharmacol Ther* 2019; **194**: 107-131.
- [20]Deseo M, Elkins A, Rochfort S, Kitchen B. Antioxidant activity and polyphenol composition of sugarcane molasses extract. *Food Chem* 2020; **314**: 126180.
- [21]Sharma JN, Al-Omran A, Parvathy SS. Role of nitric oxide in inflammatory diseases. *Inflammopharmacology* 2007; **15**(6): 252-259.
- [22]Luo G, Cheng BC, Zhao H, Fu XQ, Xie R, Zhang SF, et al. *Schisandra chinensis* lignans suppresses the production of inflammatory mediators regulated by NF- κ B, AP-1, and IRF3 in lipopolysaccharide-stimulated RAW264.7 cells. *Molecules* 2018; **23**(12): 3319.
- [23]Nakanishi G, Rosenberg DW. Multifaceted roles of PGE₂ in inflammation and cancer. *Semin Immunopathol* 2013; **35**(2): 123-137.
- [24]Zhang Y, Yan LS, Ding Y, Cheng BC, Luo G, Kong J, et al. *Edgeworthia gardneri* (Wall.) Meisn. water extract ameliorates palmitate induced insulin resistance by regulating IRS1/GSK3 β /FoxO1 signaling pathway in human HepG2 hepatocytes. *Front Pharmacol* 2020; **10**: 1666.
- [25]Zhang Y, Cheng BC, Xie R, Xu B, Gao XY, Luo G. Re-Du-Ning inhalation solution exerts suppressive effect on the secretion of inflammatory mediators *via* inhibiting IKK α / β /I κ B α /NF- κ B, MAPKs/AP-1, and TBK1/IRF3 signaling pathways in lipopolysaccharide stimulated RAW 264.7 macrophages. *RSC Adv* 2019; **9**(16): 8912-8925.
- [26]Li J, Zhang F, Li J. The immunoregulatory effects of traditional Chinese medicine on treatment of asthma or asthmatic inflammation. *Am J Chin Med* 2015; **43**(6): 1059-1081.
- [27]Ma HD, Deng YR, Tian Z, Lian ZX. Traditional Chinese medicine and immune regulation. *Clin Rev Allergy Immunol* 2013; **44**(3): 229-241.
- [28]Chen L, Teng H, Jia Z, Battino M, Miron A, Yu Z, et al. Intracellular signaling pathways of inflammation modulated by dietary flavonoids: The most recent evidence. *Crit Rev Food Sci Nutr* 2018; **58**(17): 2908-2924.
- [29]Gullón B, Lú-Chau TA, Moreira MT, Lema JM, Eibes G. Rutin: A review on extraction, identification and purification methods, biological activities and approaches to enhance its bioavailability. *Trends Food Sci Technol* 2017; **67**: 220-235.
- [30]Coleman JW. Nitric oxide in immunity and inflammation. *Int Immunopharmacol* 2001; **1**(8): 1397-1406.
- [31]Park JY, Pillinger MH, Abramson SB. Prostaglandin E₂ synthesis and secretion: The role of PGE₂ synthases. *Clin Immunol* 2006; **119**(3): 229-240.
- [32]Aktan F. iNOS-mediated nitric oxide production and its regulation. *Life Sci* 2004; **75**(6): 639-653.
- [33]Akira S, Takeda K. Toll-like receptor signalling. *Nat Rev Immunol* 2004; **4**(7): 499-511.
- [34]Lawrence T. The nuclear factor NF- κ B pathway in inflammation. *Cold Spring Harb Perspect Biol* 2009; **1**(6): a001651.
- [35]Cheng BC, Yu H, Guo H, Su T, Fu XQ, Li T, et al. A herbal formula comprising *Rosae Multiflorae Fructus* and *Lonicerae Japonicae Flos*, attenuates collagen-induced arthritis and inhibits TLR4 signalling in rats. *Sci Rep* 2016; **6**(1): 20042.
- [36]Xu M, Guo Q, Wang S, Wang N, Wei L, Wang J. Anti-rheumatoid arthritic effects of *Saussurea involucrata* on type II collagen-induced arthritis in rats. *Food Funct* 2016; **7**(2): 763-770.
- [37]Han X, Su D, Xian X, Zhou M, Li X, Huang J, et al. Inhibitory effects of *Saussurea involucrata* (Kar. et Kir.) Sch. -Bip. on adjuvant arthritis in rats. *J Ethnopharmacol* 2016; **194**: 228-235.
- [38]Yang YC, Lin HY, Su KY, Chen CH, Yu YL, Lin CC, et al. Rutin, a flavonoid that is a main component of *Saussurea involucrata*, attenuates the senescence effect in *D*-galactose aging mouse model. *Evid Based Complement Alternat Med* 2012; **2012**: 980276.
- [39]Srivastava M, Saqib U, Naim A, Roy A, Liu D, Bhatnagar D, et al. The TLR4–NOS1-AP1 signaling axis regulates macrophage polarization. *Inflamm Res* 2017; **66**(4): 323-334.
- [40]Griffith JW, Sokol CL, Luster AD. Chemokines and chemokine receptors: Positioning cells for host defense and immunity. *Annu Rev Immunol* 2014; **32**: 659-702.
- [41]Arthur JS, Ley SC. Mitogen-activated protein kinases in innate immunity. *Nat Rev Immunol* 2013; **13**(9): 679-692.
- [42]Yeh CH, Yang JJ, Yang ML, Li YC, Kuan YH. Rutin decreases lipopolysaccharide-induced acute lung injury *via* inhibition of oxidative stress and the MAPK-NF- κ B pathway. *Free Radic Biol Med* 2014; **69**: 249-257.
- [43]Fitzgerald KA, McWhirter SM, Faia KL, Rowe DC, Latz E, Golenbock DT, et al. IKKepsilon and TBK1 are essential components of the IRF3 signaling pathway. *Nat Immunol* 2003; **4**(5): 491-496.