

Chao, J., Siu, A. M., Leung, O., Lo, A., Chu, M., Lee, W. K., ... & Chien, C. W. (2019). Chinese version of the recovery self-assessment scale: psychometric evidence from Rasch analysis and reliability estimates. *Journal of Mental Health*, 28(2), 206-212.

This is an Accepted Manuscript of an article published by Taylor & Francis in *Optimization* on 17 Nov 2018 (published online), available at: <http://www.tandfonline.com/10.1080/09638237.2018.1521931>

Chinese Version of the Recovery Self-Assessment Scale: Psychometric Evidence from Rasch Analysis and Reliability Estimates

Abstract

Aim. This study aims to develop a Chinese version of the Recovery Self-Assessment (RSA), which assesses the recovery orientation of hospital-based mental health services.

Methods. We conducted forward and backward translations of the RSA. After making modifications suggested by a team of content experts, the Chinese Recovery Self-Assessment Service User version (CRSA-R) was ready for testing. We recruited 350 people with mental illnesses who regularly attend hospital, day, and outpatient mental health services. The participants completed the CRSA-R, and convergent measures on hope and mental well-being.

Results. The Rasch analysis provides support for five of the six factors, and suggests that the “Life Goal” factor could be further split into two factors of “Life Goals for My Recovery” and “Life Goals Supported by Staff.” We identified three misfit items (items 6, 12, and 17) that could be considered for removal in the future. Both the internal consistency and test-retest reliability are between satisfactory and very good within each subscale, with the exception of the Choice subscale. The seven subscales had low positive correlations with measures of hope and mental well-being, which supported the convergent validity of CRSA-R.

Conclusion. The results supported the factor structure, reliability, and convergent validity of the CRSA-R.

Keywords: Mental health, Recovery, Recovery Self-assessment, Rasch, Psychometric

Introduction

Mental health recovery is a journey of healing, which enables a person with a mental illness to live a fuller and more meaningful life in their community, and make choices to reach their full potential (Whitney & Drake, 2010). Mental health services have faced ongoing criticism that they place too much emphasis on the person's sick role instead of paying more attention to the person's illness experience and recovery process (Slade, 2009). In many modern societies, traditional mental health systems focus on symptom management, since the societies have low expectations for the recovery of people with severe mental illnesses (Maassen, Schrevel, Dedding, Broerse, & Regeer, 2017). The recovery model advocates that persons with mental illnesses can progress beyond the impact of mental illness and develop new meaning and purpose in their lives. Over the past two decades, there has been a paradigm shift within mental health services toward recovery-oriented practice, which has been widely adopted around the world in countries such as the USA, Australia, New Zealand, and the UK (Ramon et al., 2007).

The recovery model advocates that persons with severe mental illnesses have self-determination, autonomy, and choice in their own care and recovery; it is imperative for service providers to adopt a consumer-driven approach that encourages the participatory involvement of service users. Service users are one of the key stakeholders in the planning and evaluation of recovery-oriented services, in addition to service providers, service administrators and caregivers (Ostrow & Adams, 2012). Over the years, practitioners and researchers around the world have developed a number of self-completed questionnaires to evaluate the recovery orientation of mental health services and the outcomes of recovery-oriented services. Among the eleven instruments reviewed by Burgess, Pirkis, Coobs, & Rosen (2011), the Recovery Self-Assessment (RSA) (O'Connell et al., 2005) is one of the four instruments that met the criteria for

1
2
3 a quality assessment instrument of recovery practice: 1) the instrument measures areas that are
4 directly relevant to recovery orientation, 2) it is manageable and easy to administer, 3) it has
5 gone through an appropriate process of development and validation, 4) it includes a consumer
6 perspective, 5) it is applicable to the local context, and 6) it is acceptable to consumers. In
7 addition, the RSA is also one of the most widely used instruments globally (Williams et al.,
8 2012).

9
10
11
12
13
14
15
16
17
18 The present study selected the RSA for conducting ongoing evaluation of recovery-
19 oriented services in Hong Kong for several reasons. First, the RSA is versatile; it has been
20 applied in both hospital and community-based services (Salyers et al., 2007), and could assess
21 both individuals' recovery as well as the recovery-orientation of the service (Burgess et al.,
22 2011). Second, there is a wealth of evidence supporting the psychometric properties of the RSA
23 worldwide. Several countries, such as the USA (Campbell-Orde et al., 2005) and Australia
24 (Burgess et al., 2011), recommended the use of the RSA for regulatory evaluation of recovery
25 services. Third, the RSA Revised version (RSA-R) has four parallel versions – person in
26 recovery, family/significant other/advocate, provider, and administrator versions – which enables
27 a more comprehensive evaluation of recovery-oriented services by various stakeholders
28 (Campbell-Orde et al., 2005; O'Connell et al., 2007). Lastly, the RSA has been translated and
29 applied in many non-English speaking countries and cultural contexts without the need for major
30 modifications (Chiba et al., 2010; Rosenberg et al., 2015; Ye, Pan, Wong, & Bola, 2013).

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49 In fact, two studies have developed a Chinese version of the RSA-R for use in some
50 community mental health settings in Hong Kong. Some initial psychometric properties were
51 collected (Ye et al., 2013), and cross-validation of the RSA-R with other mental health recovery
52 measures and convergent measures (like hope) were conducted. However, it was necessary to
53
54
55
56
57
58
59
60

1
2
3 address several gaps in clinical application and validation research of the Chinese RSA-R. First,
4 in the studies by Ye et al. (2013) and Bola, Chan, Chen, & Ng, 2016), one translated the RSA
5 while the other translated the RSA-R respectively. Both studies attempted to validate the RSA or
6 RSA-R as one of the instruments in a battery of recovery measures for use in community mental
7 health settings. However, both studies did not clearly report on the translation process or the
8 review of content validity and cultural relevance of the RSA or RSA-R for use with Chinese
9 populations. This study aims to conduct forward and backward translation, as well as a detailed
10 expert panel review of quality of translation, content validity, and cultural relevance before pilot
11 test of the RSA-R person-in-recovery version.
12
13
14
15
16
17
18
19
20
21
22

23
24
25 Second, there are uncertainties in the factor structure of the RSA in both the Chinese and
26 English versions, despite many studies that have tried to explore and confirm the factor structure.
27 For the English version, two key studies came up with similar groupings of items under the
28 following five factors: 1) goal/success orientation and hope, 2) reliance on others, 3) personal
29 confidence, 4) no domination by symptoms, and 5) willingness to ask for help (Corrigan et al.,
30 2004; McNaught et al., 2007). A study of the Japanese version of the RSA replicated this five-
31 factor structure (Chiba et al., 2010); however, this factor structure is quite different from the
32 original theoretical design of the instrument (O'Connell et al., 2005). Furthermore, the study on
33 the translated Chinese version by Ye et al. (2013) did not find a stable factor structure, and used
34 the RSA total score for further analyses. In a recent attempt, Barbic et al. (2015) developed a 12-
35 item RSA brief version using the Rasch measurement model. While there is support for the
36 unidimensionality and reliability of this brief version, the authors commented that there is a need
37 to re-visit the categorization of items under the original theory, as well as the selection of items
38 used to form the brief version. In summary, there was a strong need to re-examine the factor
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 structure of the RSA. The current study uses Rasch analysis to evaluate the construct validity of
4 the CRSA-R on the basis that it could examine whether it is appropriate to treat the CRSA-R as a
5 unidimensional scale, and evaluate how items contribute to diversity of opinion among
6 respondents.
7
8
9
10
11
12

13 Third, there is a need to examine the convergent validity and test-reliability of the
14 Chinese version of the RSA (Williams et al., 2012). Two previous RSA studies on the Chinese
15 population provided some initial estimates on the internal consistency and reliability of the total
16 score, and its correlations with quality of life and several recovery measures (Bola et al., 2016;
17 Ye et al., 2013). The present study will also provide estimates of internal consistency and test-
18 retest reliability of the scale and subscales, as well as providing evidence on its validity with
19 convergent measures of mental well-being and hope. With the additional psychometric
20 information from this study, clinicians and researchers would have a better understanding of how
21 ready the Chinese version of RSA is for further application in clinical practice and research.
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 **Method**

37
38
39 This study aims to translate the Service User version of the Recovery Self-Assessment
40 into Chinese (abbreviated CRSA-R) and evaluate the dimensionality of the translated version
41 using the Rasch measurement model approach. We also aim to provide estimates of internal
42 consistency and test-retest reliability, as well as evidence for its convergent validity.
43
44
45
46
47
48

49 **Participants**

50
51 We selectively recruited a sample of persons with mental illness who are attending
52 occupational therapy services in six different settings under the Hong Kong Hospital Authority.
53
54
55
56
57
58
59
60

1
2
3 All participants met the following inclusion criteria: 1) aged above 18 years, 2) diagnosed with a
4 psychiatric disorder, and 3) regularly attends rehabilitation programs in in-patient, outpatient, or
5 day rehabilitation services. Among the participants (N = 350), around half were male (n = 168,
6 48%) and the remainder were female (Table 1). Participants had a mean age of 42.63 years (SD
7 = .66). Almost half (49.1%) of the participants had a high school education, while 15.8% had a
8 tertiary education. The key diagnoses of the participants were schizophrenia, schizotypal and
9 delusional disorder (65.6%), and mood disorders (25.8%). In-patients (44%) and day-patients
10 (40%) made up the main proportions of the sample, while the rest were outpatients (16%). The
11 mean duration of living with psychiatric illness was 13.3 years (SD = .58), with a range of less
12 than 1 year to a maximum of 45 years.
13
14
15
16
17
18
19
20
21
22
23
24
25

26 27 **Instruments**

28
29 **Chinese Recovery Self-Assessment, Service User version (CRSA-R).** After obtaining
30 permission from the original author of RSA, we translated the Recovery Self-Assessment (RSA-
31 R) Personal in Recovery version into Chinese (the CRSA-R). A team of five mental health
32 experts reviewed the first draft of the translated instrument. After some revisions, the second
33 draft was reviewed by 25 mental health professionals and a final version was drafted. We hired a
34 language expert to conduct a backward translation of the CRSA-R into English. We compared
35 the back translated version (in English) with the original English version, and final
36 improvements were made to the translated Chinese version.
37
38
39
40
41
42
43
44
45
46
47

48
49 **Adult Hope Scale (AHS).** The AHS is a 12-item self-completed questionnaire designed
50 to measure a person's level of hope (Snyder et al., 1991). The scale has two subscales: 1) the
51 Agency subscale, which measures the person's goal-directed energy in initiating and sustaining
52 actions, and 2) Pathways, which measures the person's belief in generating activities to pursue
53
54
55
56
57
58
59
60

1
2
3 and reach goals. We anticipated the CRSA-R would have significant positive correlations with
4 the convergent measure of AHS, as we expected a recovery-oriented environment to cultivate
5 higher hope in the service users.
6
7
8
9

10
11 **Chinese Warwick-Edinburgh Mental Well-being Scale (C-WEMWBS).** This seven-
12 item instrument was designed to measure the mental well-being of subjects. Both validation
13 studies of the original English and Chinese versions of the WEMWBS reported a unidimensional
14 factor structure, as well as excellent reliability and construct validity (Brown, et al., 2009; Bass,
15 Dawkin, Muncer, Vigurs, & Bostock, 2015; Ng, et al., 2014). We anticipated mental well-being
16 to be a convergent measure with recovery orientation of mental health services.
17
18
19
20
21
22
23
24
25

26 **Procedure**

27
28 We obtained ethics approval from the Research Ethics Review Committee of the Hospital
29 Authority and The Hong Kong Polytechnic University to conduct this project. We briefed
30 potential participants who fulfilled the selection criteria, covering information on the purpose
31 and procedures of the study, and how they could participate. We then invited them to join, and
32 required that those who agreed to participate in the study signed a consent form. The participants
33 completed a set of questionnaires that included the CRSA-R, AHS, and C-WEMWBS, and
34 provided some basic background information. Two hundred and ninety six (84.6%) out of 350
35 participants agreed to complete the CRSA-R for a second time to provide data to evaluate the
36 test-retest reliability. The mean duration between the first and second completion of the CRSA-R
37 was 13.8 (SD = 7.7) days.
38
39
40
41
42
43
44
45
46
47
48
49
50

51 **Statistical analysis**

52
53
54
55
56
57
58
59
60

1
2
3 Rasch analysis was used to evaluate the construct validity of the CRSA-R, specifically
4
5 for its unidimensionality or the extent to which the CRSA-R items measure a single construct
6
7 defining recovery orientation. Based on item response theory, Rasch analysis is a probabilistic
8
9 model that examines a matrix of person-ability versus item-difficulty on a common scale and
10
11 converts ordinal scores obtained from rating scales into interval-level measures or so-called
12
13 ‘logits’ (Bond & Fox, 2007). The unidimensionality of a scale can be assessed by using the
14
15 residuals derived from Rasch measures that achieve interval-level scaling. Rasch analysis has
16
17 been recently applied to aid in the validation of the recovery outcome in people with mental
18
19 illnesses (Barbic, Kidd, Davidson, McKenzie, & O’Connell, 2015; Hancock, Scanlan, Honey,
20
21 Bundy, & O’Shea, 2015; McGuire, Kean, Bonfils, Presnell, & Salyers, 2014). As the CRSA-R
22
23 uses a five-point Likert scale (*strongly disagree* to *strongly agree*) across all items, the Rasch
24
25 Rating Scale Model (Andrich, 1978) was applied and the WINSTEPS software version 3.73
26
27 (Linacre, 2011) was used to conduct the analysis.
28
29
30
31
32
33

34 To examine the unidimensionality, a Rasch-based principal component analysis (PCA) of
35
36 residuals with item goodness-of-fit analysis was conducted. The unidimensionality of the CRSA-
37
38 R items is supported if the Rasch identified construct (principal component) account for >50% of
39
40 the total variance, and if the size of the first contrast (the largest secondary component after the
41
42 principal component is removed) is less than 2.0 eigenvalue (Raïche, 2005). We would use
43
44 Rasch-based goodness-of-fit statistics to examine how well the CRSA-R items fit with the
45
46 model’s expectations for hierarchical difficulties. The two types of fit statistics, infit and outfit
47
48 are reported using mean square (MnSq) and standardized Z values (Zstd). Infit and outfit with
49
50 MnSq <1.4 in combination with Zstd values of <2.0 are indicators of acceptable model fit (Bond
51
52 & Fox, 2007; Chien & Bond, 2009). Items with MnSq > 1.4 and Zstd > 2 indicate that the item
53
54
55
56
57
58
59
60

1
2
3 potentially misfit (Infit and Outfit MnSq > 1.4 or Infit & Outfit Zstd > 2.0). The above evidence
4 suggested that the C-RSA is not a unidimensional scale, which led to the decision to examine the
5 specific pattern of factor loading (based on the Rasch principal component analysis of residuals)
6
7 under each of the subscales separately.
8
9
10
11
12

13 **Principal Component Analysis (PCA) of Residuals of CRSA-R subscales.**

14
15 Regarding the dimensionality of CRSA-R items under each subscale, the Rasch analysis
16 supports five out of the six subscales (Table 2). The PCA results for the “Life Goal” subscale
17 were not quite satisfactory because the total variance explained by the Rasch measure was 43%,
18 with the first contrast having an Eigenvalue of 2.1, and there were two items (12 and 17)
19 showing as misfit. After iteratively removing these two misfit items, the PCA of residuals results
20 remained unsatisfactory. Therefore, we split the remaining nine items into two sub-domains of
21 the “Life Goal” subscale, based on the factor loading of PCA of residuals. One sub-domain
22 includes items 3, 7, 8, and 9 (labelled “Life Goals for My Recovery”) and the other includes
23 items 16, 18, 28, 31, and 32 (labelled “Life Goals Guided by Staff”). The PCA results of
24 residuals were satisfactory for the two sub-domains and we did not find misfit items in either
25 sub-domain (see Appendix A). Based on this analysis, we suggest dividing the “Life Goals”
26 domain into two unidimensional sub-domains.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 The PCA result of residuals for the “Choice” dimension was acceptable, indicated by a
45 total explained variance of 39.7% and Eigenvalue of 1.6 for the first contrast. Two items (4 and
46 6), showed a slight misfit with the Rasch model’s expectations. After we removed item 6, the
47 PCA results improved (a total explained variance of 43.6%). Thus, we suggest removing item 6
48 in this subscale, and retaining the other four items in this subscale.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 The PCA results of residuals for the rest of the four subscales, including the Individually-
4 Tailored Service (ITS), Diversity of Treatment Options (DTO), Involvement, and Invite
5 dimension, are satisfactory. The total variance explained by the Rasch measure is substantial
6 (46.2–69.3%), and the first contrast has the eigenvalue of < 2.0 . There are no items showing a
7 significant misfit with the Rasch model's expectations. We suggest using these four subscales
8 without further change.
9
10
11
12
13
14
15
16
17

18 In addition, no subscales of the CRSA-R revealed the floor effects (0.3–3.4%). However,
19 obvious ceiling effects were found in all the subscales of the CRSA-R except for the Choice
20 (13.4%) and DTO (14.6%) subscales. Specifically, 32.9% of the participants achieved the
21 maximum scores in the Invite subscale that includes merely two items, and 15.4–19.1% obtained
22 the maximum scores in the rest of the four subscales including the ITS, Involvement, and two
23 new Life Goal subscales.
24
25
26
27
28
29
30
31
32

33 **Reliability**

34
35 From the results of Rasch analysis (Table 2), item reliability of the subscales ranged from
36 0.73 to 0.98 and all the estimates were acceptable to excellent. For person reliability, the
37 estimates were generally lower than item reliability and ranged from 0.48 to 0.75. We also
38 estimated the reliability of the CRSA-R using the classical test model (Table 3), and Cronbach's
39 α is 0.95 for all 30 items. The Cronbach's α of the seven subscales ranged from 0.61 to 0.90,
40 which indicated fair to very good internal consistency of the subscales. Test-retest reliability of
41 the subscales was acceptable to very good, as indicated by ICCs ranging from 0.77 to 0.94.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Convergent Validity

The seven CRSA-R subscales had positive and significant correlations with the Agency (r ranges from .29 to .42) and Pathway subscale of the Adult Hope Scale (r ranges from .27 to .39) (Table 4). The correlations are mostly of low to moderate strength. The CRSA-R subscales also have significant positive correlations with the SWEMWBS, with r ranging from .19 to .36.

Relationship between Demographic Profile and CRSA-R scores

The CRSA-R had low but significant correlations with age, education level, and duration of illness. CRSA-R subscales had significant positive correlations with age, with r ranging from .14 to .27. Six of the seven CRSA-R subscales (excepting the “Invite” subscale) also had significant positive correlations with duration of illness, with r ranging from .14 to .20. In general, this suggests that older age and longer duration of illness is associated with higher CRSA-R ratings by the participants. On the other hand, higher education is associated with lower scores in five of the seven CRSA-R ratings, as the correlations between them are negative and significant (r ranging from .14 to .27). We did not find significant differences in the CRSA-R subscale scores among participants attending in-patient, outpatient, and day patient services. The F-value varies from .12 to 2.59, with p-values ranging from .08 to .89.

Discussion

There are several important observations ensuing from this validation study. First, the results clearly showed that the CRSA-R is not a unidimensional scale, and it would not be meaningful to use the total score of all items to interpret the recovery orientation of the service. This is consistent with the original conceptualization and design of the RSA that recovery is a multidimensional construct (O'Connell, Tondora, Croog, Evans, & Davidson, 2005). Second, the

1
2
3 analysis found that three items (6, 12, and 17) have a misfit with the Rasch model. This indicates
4 that some item responses may be erratic, may belong to different constructs, or are unlikely to
5 differentiate the responses of participants as good as other items within the same subscale. We
6 discussed the proposal to remove these items with expert panel members, but most panel
7 members suggested that the items should be retained as these items represent important aspects
8 of the subscales they measure. Therefore, we suggest that these items should be used with
9 caution under the “Life Goals” or “Choice” subscales.
10
11
12
13
14
15
16
17
18
19

20 Third, the Rasch analysis and principal component analyses of residuals suggested only
21 one major change to the factor structure. We proposed to divide the “Life Goals” subscale into
22 two subscales, which we renamed as “Life Goals for My Recovery” and “Life Goals Supported
23 by Staff.” The “Life Goals for My Recovery” subscale includes four items that are phrased as
24 “staff members encourage, believe, or are confident that the person could recover (item 3, 7),
25 could manage one’s symptoms (item 8), and make choices about daily living (item 9).” For the
26 “Life Goals Supported by Staff” subscale, the items include phrases that indicate “the staff
27 member helps or encourages me to ... (do different things),” like set up life goals (item 16), other
28 activities (item 18), achieves life goals (item 28). The two other items in this subscale are related
29 to the resourcefulness of staff members (item 31) and whether they come from a diverse
30 background (item 32). In sum, we recommend splitting the “Life Goals” subscale into the two
31 subscales of “Life Goals for My Recovery” and “Life Goals Guided by Staff” in future use, and
32 the CRSA-R would have a 7 instead of 6 subscales. This contrasts with a previous exploratory
33 factor analysis of another version of Chinese RSA (Ye et al., 2013), which could not find
34 sufficient evidence in support for a multi-dimensional factor structure of the RSA.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Fourth, we found that most of the subscales of the CRSA-R had satisfactory to good
4 internal consistency and test-retest reliability. The overall Cronbach's α of all items is 0.95,
5
6 which is comparable to estimates from previous local studies on the RSA and RSA-R (Bola et al.,
7
8 2016; Ye et al., 2013). Only the four-item "Choice" subscale had internal consistency
9
10 (Cronbach's $\alpha = .61$) and test-retest reliability ($ICC = .77$) that were lower than common
11
12 standards. Upon examining the "Cronbach's α if item deleted" measure, there appeared to be no
13
14 gain in internal consistency if any one of the four items was removed – item total correlations of
15
16 the four items ranged from .30 to .47. In further application of the CRSA-R, we advise noting
17
18 that the reliability of the Choice subscale is less satisfactory than other subscales.
19
20
21
22
23
24

25 Fifth, there are several observations from the correlation studies between CRSA-R and
26
27 the two convergent measures of hope (AHS) and mental well-being (SWEMWBS). Both
28
29 measures had low and significant correlations with CRSA-R. This is consistent with expectations
30
31 that a recovery-oriented service environment could promote the hope and mental well-being of
32
33 clients. This provides support to the convergent validity of the CRSA-R. In previous local studies
34
35 of the Chinese RSA and RSA-R, researchers focus on exploring relationships among several
36
37 recovery-oriented measures and quality of life measure. (Bola et al., 2016; Ye et al., 2013). The
38
39 current study adds new information on the convergent validity of the CRSA-R.
40
41
42
43

44 Lastly, there are low but significant correlations between CRSA-R scores in demographic
45
46 variables. Age and duration of illness had a positive correlation while education and duration of
47
48 illness had a negative correlation. These results suggest that persons who are older and have had
49
50 a longer duration of illness would be may have a more positive evaluation of the recovery
51
52 orientation of the services. As age and duration of illness had a high correlation, we could not be
53
54 sure if age or duration was the key factor contributing to this positive correlation with CRSA-R
55
56
57
58
59
60

1
2
3 scores. Persons in recovery with a higher level of education could have higher expectations or be
4 more critical in the evaluation of the recovery-oriented services. Few studies documented the
5
6 more critical in the evaluation of the recovery-oriented services. Few studies documented the
7
8 association of CRSA-R scores with the demographic background variables, and in future studies
9
10 could further examine the potential influence of these variables on the person's view of recovery-
11
12 oriented services.
13

14 15 **Study Limitations**

16
17 This study has several limitations. First, the study only covers some of the key aspects of
18
19 a full validation study, including investigations of the factor structure; reliability; and convergent
20
21 validity. Due to limitations on resources and in order for the CRSA-R to be ready for program
22
23 evaluation within a reasonable period, we could only include these key aspects. Second, all
24
25 participants of this study were recruited from hospital care settings. Most participants were in-
26
27 patients or day patients within the Hospital Authority in Hong Kong, while the two existing local
28
29 studies recruited participants from community mental health services. The results of this current
30
31 study may not be directly comparable with previous studies on the Chinese RSA-R. Third, the
32
33 test-retest period is almost two weeks on average. Although the two-week test-retest period is
34
35 commonly used in validation studies, we cannot completely ensure that every participant had a
36
37 uniform test-retest period, since there were practical difficulties such as the attendance of some
38
39 participants to outpatient settings.
40
41
42
43
44
45
46
47

48 **Conclusion**

49
50 This study has provided support to the factor structure, reliability, and convergent validity of the
51
52 CRSA-R. The Rasch analysis suggests that the CRSA-R includes seven unidimensional
53
54 subscales, and we identified three misfit items that could be removed from the scale. The internal
55
56
57
58
59
60

1
2
3 consistency and test-retest reliability are satisfactory to very good for most subscales, except that
4
5 the four-item “Choice” subscale has relatively low internal consistency and reliability. As
6
7 expected, the CRSA-R scores have significant positive correlations with convergent measures of
8
9 hope and mental well-being. Only minor revisions would be needed before the CRSA-R could be
10
11 adopted in the evaluation of recovery-oriented mental health services.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

References

- Andrich, D. (1978). Application of a psychometric rating model to ordered categories which are scored with successive integers. *Applied Psychological Measurement*, 2(4), 581-594.
- Barbic, S. P., Kidd, S. A., Davidson, L., McKenzie, K., & O'Connell, M. J. (2015). Validation of the Brief Version of the Recovery Self-Assessment (RSA-B) Using Rasch Measurement Theory. *Psychiatric Rehabilitation Journal*, 38(4), 349.
- Bass, M., Dawkin, M., Muncer, S., Vigurs, S., & Bostock, J. (2016). Validation of Warwick-Edinburgh Mental Well-being Scale (WEMWBS) in a population of people using Secondary Care Mental Health Services. *Journal of Mental Health*, 25(4), 323-329.
- Bola, J., Chan, T. H. C., Chen, E. H., & Ng, R. (2016). Cross-validating Chinese Language Mental Health Recovery Measures in Hong Kong. *Research on Social Work Practice*, 26(6), 630-640.
- Bond, T. G., & Fox, C. M. (2007). *Applying the Rasch model: Fundamental measurement in the human sciences* (2nd ed.). Mahwah, NJ: Lawrence Erlbaum Associates.
- Burgess, P., Pirkis, J., Coops, T., & Rosen, A. (2011). Assessing the Value of Existing Recovery Measures for Routine Use in Australian Mental Health Services. *Australian and New Zealand Journal of Psychiatry, Early Online*, 1-14,
- Campbell-Orde, T., Chamberlin, J., Carpenter, J., & Leff, H. S. (2005). Measuring the Promise: A Compendium of Recovery Measures, Volume II, The Evaluation Center @ HSRI.
- Chiba, R., Miyamoto, Y., & Kawakami, N. (2010). Reliability and Validity of the Japanese Version of the Recovery Assessment Scale (RAS) for People with Chronic Mental Illness: Scale Development. *International Journal of Nursing Studies*, 47(3), 314-322.
- Chien, C. W., & Bond, T. G. (2009). Measurement properties of fine motor scale of Peabody developmental motor scales-second edition: A Rasch analysis. *American Journal of Physical Medicine and Rehabilitation*, 88(5), 376-386.
- Corrigan, P. W., Salzer, M., Ralph, R. O., Sangster, Y., & Keck, L. (2004). Examining the Factor Structure of the Recovery Assessment Scale. *Schizophrenia Bulletin*, 30(4), 1035-1041.

- 1
2
3 Hancock, N., Scanlan, J. N., Honey, A., Bundy, A. C., & O'Shea, K. (2015). Recovery
4 Assessment Scale - Domains and Stages (RAS-DS): Its feasibility and outcome
5 measurement capacity. *Australian and New Zealand Journal of Psychiatry*, 49(7), 624-633.
6
7
8 Hobart, J. C., & Thompson, A. J. (2001). The five item Barthel index. *Journal of Neurology*
9 *Neurosurgery and Psychiatry*, 71(2), 225-230.
10
11
12 Hogan, M. F. (2003). *Achieving the Promise: Transforming Mental Health Care in America,*
13 *Executive Summary. The President's New Freedom Commission on Mental Health.*
14 Maryland: US Department of Health and Human Services.
15
16
17 Linacre, J. M. (2011). *User's guide to WINSTEPS.* Chicago, IL: MESA Press.
18
19 Maassen, E. F., Schrevel, S. J., Dedding, C. W., Broerse, J. E., & Regeer, B. J. (2017).
20 Comparing patients' perspectives of "good care" in Dutch outpatient psychiatric services
21 with academic perspectives of patient-centred care. *Journal of Mental Health*, 26(1), 84-94.
22
23
24 McGuire, A. B., Kean, J., Bonfils, K., Presnell, J., & Salyers, M. P. (2014). Rasch analysis of the
25 illness management and recovery scale-clinician version. *Journal of Evaluation in Clinical*
26 *Practice*, 20(4), 383-389.
27
28
29 McNaught, M., Caputi, P., Oades, L. G., & Deane, F. P. (2007). Testing the Validity of the
30 Recovery Assessment Scale Using an Australian Sample. *Australian and New Zealand*
31 *Journal of Psychiatry*, 41(5), 450-457.
32
33
34 Ng, S. S., Lo, A. W., Leung, T. K., Chan, F. S., Wong, A. T., Lam, R. W., & Tsang, D. K.
35 (2014). Translation and Validation of the Chinese Version of the Short Warwick-
36 Edinburgh Mental Well-being Scale for Patients with Mental Illness in Hong Kong. *East*
37 *Asian Archives of Psychiatry*, 24(1), 3.
38
39
40 O'Connell, M., Tondora, J., Croog, G., Evans, A., & Davidson, L. (2005). From Rhetoric to
41 Routine: Assessing Perceptions of Recovery-oriented Practices in a State Mental Health
42 and Addiction System. *Psychiatric rehabilitation journal*, 28(4), 378.
43
44
45 Ostrow, L., & Adams, N. (2012). Recovery in the USA: From Politics to Peer Support.
46 *International Review of Psychiatry*, 24(1), 70-78.
47
48
49 Portney, L. G., & Watkins, M. P. (2009). *Foundations of clinical research: Application to*
50 *practice* (3rd ed.). Upper Saddle River, NJ: Pearson/Prentice Hall.
51
52
53 Raïche, G. (2005). Critical eigenvalue sizes in standardized residual principal components
54 analysis (PCA). *Rasch Measurement Transactions*, 22(1), 1162.
55
56
57
58
59
60

- 1
2
3 Ramon, S., Healy, B., & Renouf, N. (2007). Recovery from Mental Illness as an Emergent
4 Concept and Practice in Australia and the UK. *International Journal of Social*
5 *Psychiatry*, 53(2), 108-122.
6
7
8 Rosenberg, D., Svedberg, P., & Schön, U. K. (2015). Establishing a Recovery Orientation in
9 Mental Health Services: Evaluating the Recovery Self-Assessment (RSA) in a Swedish
10 Context. *Psychiatric Rehabilitation Journal*, 38(4), 328.
11
12
13 Salyers, M. P., Tsai, J., & Stultz, T. A. (2007). Measuring Recovery Orientation in a Hospital
14 Setting. *Psychiatric Rehabilitation Journal*, 31, 131-137.
15
16
17 Slade, M. (2009). *Personal Recovery and Mental Illness: A Guide for Mental Health*
18 *Professionals*. Cambridge University Press.
19
20 Snyder, C. R., Harris, C., Anderson, J. R., Holleran, S. A., Irving, L. M., Sigmon, S. T.,
21 Yoshinob, U. L., Gibb, J., Langelle, C., & Harney, P. (1991). The Will and the Ways:
22 Development and Validation of an Individual-differences Measure of Hope. *Journal of*
23 *Personality and Social Psychology*, 60(4), 570.
24
25
26 Stewart-Brown, S., Tennant, A., Tennant, R., Platt, S., Parkinson, J., & Weich, S. (2009).
27 Internal Construct Validity of the Warwick-Edinburgh Mental Well-being Scale
28 (WEMWBS): A Rasch Analysis Using Data from the Scottish Health Education
29 Population Survey. *Health and Quality of Life Outcomes*, 7(1), 15.
30
31
32
33 Whitley, R., & Drake, R. E. (2010). Recovery: A Dimensional Approach. *Psychiatric*
34 *Services*, 61(12), 1248-1250.
35
36
37 Williams, J., Leamy, M., Bird, V., Harding, C., Larsen, J., Le Boutillier, Oades, L., & Slade, M.
38 (2012). Measures of the Recovery Orientation of Mental Health Services: Systematic
39 Review. *Social Psychiatry and Psychiatric Epidemiology*, 47(11), 1827-1835.
40
41
42 Ye, S., Pan, J. Y., Wong, D. F. K., & Bola, J. R. (2013). Cross-validation of Mental Health
43 Recovery Measures in a Hong Kong Chinese Sample. *Research on Social Work*
44 *Practice*, 23(3), 311-325.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Characteristics of Study Participants (N = 350)

Variables	Categories	n (%)
Sex	Male	168 (48.0%)
	Female	182 (52.0%)
Educational Level	No formal education	1 (0.3%)
	Primary P 1-6	42 (12.1%)
	Secondary Form 1-3	79 (22.7%)
	Secondary Form 4-7	171 (49.1%)
	Tertiary education	55 (15.8%)
Diagnosis	Organic, including symptomatic, mental disorders (F00-F09)	2 (0.6%)
	Mental and behavioral disorders due to psychoactive substance use (F10-F19)	15 (4.3%)
	Schizophrenia, schizotypal and delusional disorders (F20-F29)	229 (65.6%)
	Mood or affective disorders (F30-F39)	90 (25.8%)
	Neurotic, stress-related, and somatoform disorder (F40-F48)	6 (1.7%)
	Disorders of adult personality and behavior (F60-F69)	3 (0.9%)
	Mental retardation (F70-F79)	4 (1.1%)
Service attended	In-patient	154 (44.0%)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Variables	Categories	n (%)
	Out-patient	56 (16.0%)
	Day-patient	140 (40.0%)

For Peer Review Only

Table 2. Summary of Results of Rasch Analysis and Principal Component Analysis

Subscales	Items under each subscale	Items suggested for removal	Total % of variance explained (eigenvalue for 1 st contrast)	Person (Item) Reliability
Life Goals for My Recovery	3, 7, 8, 9	12, 17	56.6% (1.5)	0.75 (0.73)
Life Goals Guided by Staff	16, 18, 28, 31, 32		52.0% (1.7)	0.66 (0.89)
Choice	4, 5, 10, 27	6	43.6% (1.6)	0.48 (0.98)
Individually-Tailored Service	11, 13, 19, 30	---	51.3% (1.6)	0.68 (0.90)
Diversity of Treatment Option	14, 15, 20, 21, 26	---	46.2% (1.6)	0.65 (0.94)
Involvement	22, 23, 24, 25, 27, 29	---	53.3% (1.5)	0.74 (0.93)
Invite	1, 2	---	69.3% (0)	0.72 (0.86)

Table 3. Reliability Estimates of the CRSA-R

RSA-R subscales	Number of items	Internal Consistency (Cronbach's α)	Test-retest Reliability (ICC)
Life Goals for My Recovery	4	.82	.77
Life Goals Guided by Staff	5	.87	.91
Choice	4	.61	.77
Individually-Tailored Service	4	.82	.87
Diversity of Treatment Option	5	.81	.88
Involvement	6	.90	.94
Invite	2	.78	.80
All items	30	.95	.98

Table 4. Correlations between CRSA-R and convergent measures or demographic variables

Convergent measures and demographic variables	Life Goals subscales						
	Life Goal for My Recovery	Life Goals Guided by Staff	Choice	Individually -Tailored Service	Diversity of Treatment Option	Involvement	Invite
Adult Hope Scale (AHS)							
Agency Subscale	.43**	.35**	.32**	.32**	.34**	.29**	.34**
Pathway Subscale	.39**	.36**	.31**	.31**	.35**	.28**	.27**
SWEMWBS	.36**	.24**	.22**	.25**	.26**	.19**	.25**
Age	.27**	.22**	.20**	.22**	.21**	.14*	.26**
Education Level	-.20**	-.22**	-.15*	-.19**	-.19**	-.11	-.11
Duration of Illness	.20**	.17*	.15*	.19**	.17**	.14*	.08