

# Convergence of finite element solutions of stochastic partial integro-differential equations driven by white noise

Max Gunzburger · Buyang Li · Jilu Wang

Received: date / Accepted: date

**Abstract** Numerical approximation of a stochastic partial integro-differential equation driven by a space-time white noise is studied by truncating a series representation of the noise, with finite element method for spatial discretization and convolution quadrature for time discretization. Sharp-order convergence of the numerical solutions is proved up to a logarithmic factor. Numerical examples are provided to support the theoretical analysis.

**Keywords:** stochastic PDE, partial integro-differential equation, space-time white noise, finite element method, convolution quadrature, error estimate

## 1 Introduction

For given  $\alpha \in (0, 2)$ , we consider the stochastic partial integro-differential equation (PDE)

$$\begin{cases} \partial_t \psi(x, t) - \Delta \partial_t^{1-\alpha} \psi(x, t) = f(x, t) + \sigma \dot{W}(x, t) & (x, t) \in \mathcal{O} \times \mathbb{R}_+ \\ \partial_t^{1-\alpha} \psi(x, t) = 0 & (x, t) \in \partial \mathcal{O} \times \mathbb{R}_+ \\ \psi(x, 0) = \psi_0(x) & x \in \mathcal{O} \end{cases} \quad (1.1)$$

in a convex polygon/polyhedron  $\mathcal{O} \subset \mathbb{R}^d$ ,  $d \in \{1, 2, 3\}$ , where  $\Delta : H^2(\Omega) \cap H_0^1(\mathcal{O}) \rightarrow L^2(\mathcal{O})$  denotes the Laplacian operator,  $f(x, t)$  a given deterministic source function,  $\psi_0(x)$  a given deterministic initial data,  $\sigma$  a given positive parameter, and  $\dot{W}(x, t)$  a space-time white noise, i.e., the time derivative of a cylindrical Wiener process on  $L^2(\mathcal{O})$  with an underlying probability sample space  $\Omega$ . The Caputo fractional time derivative/integral

---

The research of M. Gunzburger and J. Wang was supported in part by the USA National Science Foundation grant DMS-1315259, by the USA Air Force Office of Scientific Research grant FA9550-15-1-0001, and by a USA Defense Advanced Projects Agency contract administered under the Oak Ridge National Laboratory subcontract 4000145366. The research of B. Li was supported in part by the Hong Kong RGC grant 15300817.

---

Max Gunzburger

Department of Scientific Computing, Florida State University, Tallahassee, FL 32306, USA  
 E-mail: [gunzburg@fsu.edu](mailto:gunzburg@fsu.edu)

Buyang Li

Department of Applied Mathematics, The Hong Kong Polytechnic University, Kowloon, Hong Kong  
 E-mail: [buyang.li@polyu.edu.hk](mailto:buyang.li@polyu.edu.hk)

Jilu Wang

Department of Scientific Computing, Florida State University, Tallahassee, FL 32306, USA  
 E-mail: [jwang13@fsu.edu](mailto:jwang13@fsu.edu)

$\partial_t^{1-\alpha}\psi$  is defined by (cf. [16, pp. 91])

$$\partial_t^{1-\alpha}\psi(x, t) := \begin{cases} \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \frac{\partial\psi(x, s)}{\partial s} ds & \text{if } \alpha \in (0, 1], \\ \frac{1}{\Gamma(\alpha-1)} \int_0^t (t-s)^{\alpha-2} \psi(x, s) ds & \text{if } \alpha \in (1, 2), \end{cases}$$

where  $\Gamma(s) := \int_0^\infty t^{s-1} e^{-t} dt$  denotes the Euler Gamma function. If  $\alpha = 1$ , then (1.1) recovers the standard stochastic parabolic equation.

Problem (1.1) can be used to describe the behavior of complex phenomena in mathematical physics, such as viscoelasticity and heat conduction in materials with memory subject to stochastic noises [6, 17, 23]. For any given initial data  $\psi_0 \in L^2(\mathcal{O})$  and source  $f \in L^1(0, T; L^2(\mathcal{O}))$ , problem (1.1) has a mild solution  $\psi \in C([0, T]; L^2(\Omega; L^2(\mathcal{O})))$ ; see Appendix A.

Many efforts have been made in developing efficient numerical methods with rigorous error analyses for solving (1.1), with or without the stochastic noise. In [22], Lubich et.al. have considered the deterministic version of this problem in the case  $\alpha \in (1, 2)$ . The discretization used convolution quadrature (CQ) based on backward difference methods (BDFs) in time and piecewise linear finite elements in space. The authors have proved optimal-order convergence rate of the numerical scheme for nonsmooth initial data. To achieve higher-order temporal convergence rates, the CQ generated by second-order BDF and Crank-Nicolson methods have been considered in [7] and [15] for solving (1.1) and its equivalent formulation, respectively. Due to the singularity of the solution of fractional evolution PDEs, the standard BDF and Crank-Nicolson CQs need to be corrected at several initial steps to achieve the desired order of convergence. Initial correction of higher-order BDF methods for fractional evolution PDEs has been considered in [14] recently.

Compared with the deterministic problem, the major technical difficulties in the development and analyses of numerical schemes for (1.1) are due to the space-time white-noise forcing, which leads to low regularity of the solution in both time and space. In the case  $\alpha = 1$ , Allen et. al. [1] developed a fully discrete numerical scheme for solving a stochastic parabolic problem, for which the white noise was approximated by piecewise constant random processes and a sharp order of convergence was proved. See also Du and Zhang [9] for some special noises, Shardlow [25] for the space-time white noise discretized by the spectral method, and Yan [28] for a nonlinear stochastic parabolic system with Wiener process discretized by the generalized  $L^2$ -projection operator. In [17], Kovács and Printems developed a CQ based on backward Euler method for the model (1.1) with  $\alpha \in (1, 2)$ , where the  $Q$ -Wiener process was discretized by the generalized  $L^2$ -projection operator. For the space-time white noise case, a strong order of convergence of the numerical solution was proved in one-dimensional spatial domains, i.e.,

$$\mathbb{E} \|\psi(\cdot, t_n) - \psi_n^{(h)}\|_{L^2(\mathcal{O})} = O(\tau^{\frac{1}{2} - \frac{\alpha}{4} - \varepsilon} + h^{\frac{1}{\alpha} - \frac{1}{2} - \varepsilon}) \quad \text{for } \alpha \in (1, 2) \text{ and } d = 1,$$

where  $\varepsilon$  can be arbitrarily small,  $\psi(\cdot, t_n)$  and  $\psi_n^{(h)}$  denote the PDE's mild solution and numerical solution at time  $t_n$ , respectively,  $\tau$  denotes the temporal step size, and  $h$  denotes the spatial mesh size. For  $\alpha \in (0, 2)$ , a sharp order convergence rate  $O(\tau^{\frac{1}{2} - \frac{\alpha d}{4}})$  was proved in [12] for a CQ time discretization of (1.1) in general  $d$ -dimensional spatial domains, with  $d \in \{1, 2, 3\}$ , without the deterministic forcing. We refer the readers to [2, 4, 10] for numerical analysis of other nonlinear physical stochastic equations.

This article is a continuation of [12] in the spatially discrete setting, by truncating a series representation of the space-time white noise and solving the truncated problem by the finite element method. For the resulting fully discrete numerical scheme, we prove

the sharp-order convergence

$$\mathbb{E}\|\psi(\cdot, t_n) - \psi_n^{(h)}\|_{L^2(\mathcal{O})} = \begin{cases} O(\tau^{\frac{1}{2} - \frac{\alpha d}{4}} + \ell_h^{\frac{1}{2}} h^{\frac{1}{\alpha} - \frac{d}{2}}) & \text{if } \alpha \in [\frac{1}{2}, \frac{2}{d}), \\ O(\tau^{\frac{1}{2} - \frac{\alpha d}{4}} + h^{2 - \frac{d}{2}}) & \text{if } \alpha \in (0, \frac{1}{2}), \end{cases} \quad (1.2)$$

up to a logarithmic factor  $\ell_h^{\frac{1}{2}} = (\ln(e + 1/h))^{\frac{1}{2}}$ , in general  $d$ -dimensional spatial domains,  $d \in \{1, 2, 3\}$ . The main contributions of this paper are the following.

- (1) Sharper-order spatial convergence is proved in the case  $\alpha \in (1, 2)$  and  $d = 1$  (up to a logarithmic factor  $\ell_h^{\frac{1}{2}}$ ).
- (2) The error estimates are extended to  $\alpha \in (0, \frac{2}{d})$  and multi-dimensional domains.
- (3) An interesting phenomenon is found: the spatial order of convergence  $\frac{1}{\alpha} - \frac{d}{2}$  increases to  $2 - \frac{d}{2}$  as  $\alpha$  decreases to  $\frac{1}{2}$ , and stays at this order when  $\alpha$  further decreases.
- (4) Less regularity assumption on  $f$ : the error estimates in the literature all rely on certain regularity of  $\frac{\partial f}{\partial t}$  (even for the deterministic problems, cf. [13, Theorem 3.6] and [22, Theorem 3.3]). We relax such conditions to an optimal integrability condition  $f \in L^{\frac{4}{2+\alpha d}, 1}(0, T; L^2(\mathcal{O}))$  to match the convergence rate of the stochastic problem. Consequently, the source  $f$  does not need to be continuous in time.

The rest of this paper is organized as follows. In section 2, we recall some basic preliminary results, introduce the numerical scheme for problem (1.1), and state the main results. Based on an integral representation of the numerical solution and careful analyses of the resolvent operator, the strong convergence rates are proved in section 3 and section 4. Numerical examples are given in section 5 to illustrate the theoretical results.

Throughout this paper, we denote by  $C$ , with or without a subscript/superscript, a generic constant independent of  $n$ ,  $\tau$ , and  $h$ , which could be different at different occurrences.

## 2 The main results

### 2.1 Notations

We denote by  $(\cdot, \cdot)$  and  $\|\cdot\|$  the inner product and norm of  $L^2(\mathcal{O})$ , respectively. The operator norm on  $L^2(\mathcal{O})$  is also denoted by  $\|\cdot\|$  (as it is induced by the norm of  $L^2(\mathcal{O})$ ). Let  $\dot{H}^s(\mathcal{O}) \subset L^2(\mathcal{O})$  denote the Hilbert space induced by the norm

$$\|\varphi\|_{\dot{H}^s(\mathcal{O})} := \sum_{j=1}^{\infty} \lambda_j^{2s} |(\varphi, \phi_j)|^2,$$

where  $\phi_j$ ,  $j = 1, 2, \dots$ , denote the  $L^2$ -norm normalized eigenfunctions of the Laplacian operator  $-\Delta$  corresponding to the eigenvalues  $\lambda_j$ ,  $j = 1, 2, \dots$ , arranged in nondecreasing order. In particular,  $\dot{H}^0(\mathcal{O}) = L^2(\mathcal{O})$ ,  $\dot{H}^1(\mathcal{O}) = H_0^1(\mathcal{O})$  and  $\dot{H}^2(\mathcal{O}) = H^2(\mathcal{O}) \cap H_0^1(\mathcal{O})$ ; see [27]. For  $1 < p < \infty$  we denote by  $L^{p,1}(0, T; L^2(\mathcal{O}))$  the standard Lorentz space of functions defined on  $\mathcal{O} \times (0, T)$  (see [11, section 1.4]), satisfying

$$\sup_{t \in (0, T)} \int_0^t (t-s)^{-\frac{1}{p'}} \|f(\cdot, s)\| ds \leq C \|f\|_{L^{p,1}(0, T; L^2(\mathcal{O}))} \quad \forall f \in L^{p,1}(0, T; L^2(\mathcal{O})), \quad (2.1)$$

where  $p'$  denotes the dual of  $p$ , i.e.,  $\frac{1}{p'} + \frac{1}{p} = 1$ . For  $1 < p < \infty$ , the Lorentz space  $L^{p,1}(0, T; L^2(\mathcal{O}))$  is an intermediate real interpolation space between  $L^1(0, T; L^2(\mathcal{O}))$  and  $L^\infty(0, T; L^2(\mathcal{O}))$  (see [5, Theorem 5.2.1]), satisfying

$$L^q(0, T; L^2(\mathcal{O})) \hookrightarrow L^{p,1}(0, T; L^2(\mathcal{O})) \quad \forall q > p \geq 1.$$

Let  $\{t_n = n\tau\}_{n=0}^N$  denote a uniform partition of the time interval  $[0, T]$ , with a step size  $\tau = T/N$ , and  $u^n = u(x, t_n)$ . If we denote by  $f_\tau$  the following function (piecewise constant in time):

$$f_\tau(\cdot, s) = \frac{1}{\tau} \int_{t_{n-1}}^{t_n} f(\cdot, t) dt \quad \forall s \in (t_{n-1}, t_n], \quad n = 1, 2, \dots, N, \quad (2.2)$$

then it is well known that

$$\begin{aligned} \|f_\tau\|_{L^1(0, T; L^2(\mathcal{O}))} &\leq \|f\|_{L^1(0, T; L^2(\mathcal{O}))} & \forall f \in L^1(0, T; L^2(\mathcal{O})), \\ \|f_\tau\|_{L^\infty(0, T; L^2(\mathcal{O}))} &\leq \|f\|_{L^\infty(0, T; L^2(\mathcal{O}))} & \forall f \in L^\infty(0, T; L^2(\mathcal{O})). \end{aligned}$$

The real interpolation of the last two inequalities yields (see [5, Definition 2.4.1] and [5, Theorem 5.2.1])

$$\|f_\tau\|_{L^{p,1}(0, T; L^2(\mathcal{O}))} \leq C \|f\|_{L^{p,1}(0, T; L^2(\mathcal{O}))} \quad \forall f \in L^{p,1}(0, T; L^2(\mathcal{O})). \quad (2.3)$$

The last inequality will be used in this paper.

For  $\alpha \in (0, 1]$ , we approximate the Caputo fractional time derivative  $\partial_t^{1-\alpha} u(x, t_n) = \partial_t^{1-\alpha}(u(x, t_n) - u(x, 0))$  by the backward Euler CQ (cf. [15, (2.4)] and [20, 21, 22, 24]):

$$\bar{\partial}_\tau^{1-\alpha}(u_n - u_0) := \frac{1}{\tau^{1-\alpha}} \sum_{j=1}^n b_{n-j}(u_j - u_0), \quad n = 1, 2, \dots, N. \quad (2.4)$$

For  $\alpha \in (1, 2)$ , we approximate the Caputo fractional time derivative  $\partial_t^{1-\alpha} u(x, t_n)$  by the CQ without subtracting the initial data (cf. [22, (1.15)]), i.e.,

$$\bar{\partial}_\tau^{1-\alpha} u_n := \frac{1}{\tau^{1-\alpha}} \sum_{j=1}^n b_{n-j} u_j, \quad n = 1, 2, \dots, N. \quad (2.5)$$

In both (2.4) and (2.5), the coefficients  $b_j$ ,  $j = 0, 1, 2, \dots$ , are determined by the power series expansion

$$(1 - \zeta)^{1-\alpha} = \sum_{j=0}^{\infty} b_j \zeta^j \quad \forall |\zeta| < 1, \quad \zeta \in \mathbb{C}.$$

Besides, we define the standard backward Euler difference operator

$$\bar{\partial}_\tau u_n := \frac{u_n - u_{n-1}}{\tau}, \quad n = 1, 2, \dots, N. \quad (2.6)$$

The complex-valued function

$$\delta_\tau(\zeta) = \frac{1 - \zeta}{\tau} \quad \text{for } \zeta \in \mathbb{C} \setminus [1, \infty) \quad (2.7)$$

is called the generating function of the backward Euler difference operator. It plays an important role in the analysis of the CQ. In particular, for any sequence  $\{v_n\}_{n=0}^\infty \in \ell^2(L^2(\mathcal{O}))$  we have

$$\sum_{n=1}^{\infty} (\bar{\partial}_\tau^{1-\alpha} v_n) \zeta^n = \sum_{n=1}^{\infty} \frac{1}{\tau^{1-\alpha}} \sum_{j=1}^n b_{n-j} v_j \zeta^n = (\delta_\tau(\zeta))^{1-\alpha} \sum_{j=1}^{\infty} v_j \zeta^j, \quad \forall |\zeta| < 1. \quad (2.8)$$

Let  $\mathcal{T}_h$  be a quasi-uniform triangulation of the domain  $\mathcal{O}$  into  $d$ -dimensional simplexes  $\pi_h$ ,  $\pi_h \in \mathcal{T}_h$ , with a mesh size  $h$  such that  $0 < h < h_0$  for some constant  $h_0$ . A continuous piecewise linear finite element space  $X_h$  over the triangulation  $\mathcal{T}_h$  is defined by

$$X_h = \{\phi_h \in H_0^1(\mathcal{O}) : \phi_h|_{\pi_h} \text{ is a linear function}, \forall \pi_h \in \mathcal{T}_h\}.$$

Over the finite element space  $X_h$ , we denote the  $L^2$  projection  $P_h : L^2(\mathcal{O}) \rightarrow X_h$  and Ritz projection  $R_h : H_0^1(\mathcal{O}) \rightarrow X_h$  by

$$\begin{aligned} (P_h \varphi, \phi_h) &= (\varphi, \phi_h) & \forall \phi_h \in X_h, \\ (\nabla R_h \varphi, \nabla \phi_h) &= (\nabla \varphi, \nabla \phi_h) & \forall \phi_h \in X_h. \end{aligned}$$

It is well known that the  $L^2$  projection and Ritz projection satisfy the following standard error estimates ([27]):

$$\|P_h \phi\| \leq C \|\phi\| \quad \forall \phi \in L^2(\mathcal{O}), \quad (2.9)$$

$$\|P_h \phi - \phi\| \leq Ch^\gamma \|\phi\|_{\dot{H}^\gamma(\mathcal{O})} \quad \forall \phi \in \dot{H}^\gamma(\mathcal{O}), \quad \gamma \in [0, 2], \quad (2.10)$$

$$\|P_h \phi - R_h \phi\| \leq Ch^2 \|\phi\|_{\dot{H}^2(\mathcal{O})} \quad \forall \phi \in \dot{H}^2(\mathcal{O}). \quad (2.11)$$

Through defining the discrete Laplacian  $\Delta_h : X_h \rightarrow X_h$  by

$$(\Delta_h \varphi_h, \phi_h) = -(\nabla \varphi_h, \nabla \phi_h) \quad \forall \varphi_h, \phi_h \in X_h$$

and using the inverse inequality, the inequality (2.11) implies

$$\|\Delta_h(P_h \phi - R_h \phi)\| \leq Ch^{-2} \|P_h \phi - R_h \phi\| \leq C \|\phi\|_{\dot{H}^2(\mathcal{O})} \quad \forall \phi \in \dot{H}^2(\mathcal{O}). \quad (2.12)$$

Since  $\Delta_h R_h \phi = P_h \Delta \phi$ , it follows that

$$\begin{aligned} \|\Delta_h P_h \phi\| &\leq \|\Delta_h R_h \phi\| + \|\Delta_h(P_h \phi - R_h \phi)\| \\ &= \|P_h \Delta \phi\| + \|\Delta_h(P_h \phi - R_h \phi)\| \\ &\leq C \|\phi\|_{\dot{H}^2(\mathcal{O})} \quad \forall \phi \in \dot{H}^2(\mathcal{O}). \end{aligned} \quad (2.13)$$

The complex interpolation between (2.9) and (2.13) yields

$$\|\Delta_h^\gamma P_h \phi\| \leq C \|\phi\|_{\dot{H}^{2\gamma}(\mathcal{O})} \quad \forall \phi \in \dot{H}^{2\gamma}(\mathcal{O}), \quad \gamma \in [0, 1]. \quad (2.14)$$

Similarly, the complex interpolation between (2.11) and (2.12) yields

$$\|\Delta_h^\gamma(P_h \phi - R_h \phi)\| \leq Ch^{2-2\gamma} \|\phi\|_{\dot{H}^2(\mathcal{O})} \quad \forall \phi \in \dot{H}^2(\mathcal{O}), \quad \gamma \in [0, 1]. \quad (2.15)$$

The estimates (2.9)-(2.15) will be frequently used in this paper.

## 2.2 The numerical scheme and main theorem

Recall that the cylindrical Wiener process on  $L^2(\mathcal{O})$  can be represented as (cf. [8, Proposition 4.7, with  $Q = I$  and  $U_1$  denoting some negative-order Sobolev space])

$$W(x, t) = \sum_{j=1}^{\infty} \phi_j(x) W_j(t)$$

with independent one-dimensional Wiener processes  $W_j(t)$ ,  $j = 1, 2, \dots$ . We approximate the space-time white noise  $\dot{W}(x, t)$  by

$$\bar{\partial}_\tau W^M(x, t_n) = \sum_{j=1}^M \phi_j(x) \bar{\partial}_\tau W_j(t_n)$$

with  $M := [h^{-d}] + 1$ , the largest integer that does not exceed  $h^{-d} + 1$ . Clearly, we have

$$h^{-d} \leq M \leq Ch^{-d} \quad \forall 0 < h < h_0,$$

where the constant  $C$  may depend on  $h_0$ .

With the above notations, we propose the following fully discrete scheme for problem (1.1): find

$$\psi_n^{(h)} \in \begin{cases} \psi_0^{(h)} + X_h & \text{in the case } \alpha \in (0, 1] \\ X_h & \text{in the case } \alpha \in (1, 2) \end{cases} \quad n = 1, 2, \dots, N,$$

with  $\psi_0^{(h)} = P_h \psi_0$ , such that the following equations are satisfied for all  $\phi_h \in X_h$ :

$$\begin{aligned} & (\bar{\partial}_\tau \psi_n^{(h)}, \phi_h) + (\nabla \bar{\partial}_\tau^{1-\alpha} (\psi_n^{(h)} - \psi_0^{(h)}), \nabla \phi_h) \\ &= (f_n, \phi_h) + (\sigma \bar{\partial}_\tau W^M(\cdot, t_n), \phi_h), \end{aligned} \quad \text{if } \alpha \in (0, 1], \quad (2.16)$$

$$\begin{aligned} & (\bar{\partial}_\tau \psi_n^{(h)}, \phi_h) + (\nabla \bar{\partial}_\tau^{1-\alpha} \psi_n^{(h)}, \nabla \phi_h) \\ &= (f_n, \phi_h) + (\sigma \bar{\partial}_\tau W^M(\cdot, t_n), \phi_h), \end{aligned} \quad \text{if } \alpha \in (1, 2), \quad (2.17)$$

where  $f_n$  is the average of  $f$  over the subinterval  $(t_{n-1}, t_n]$ , i.e.,

$$f_n = \frac{1}{\tau} \int_{t_{n-1}}^{t_n} f(\cdot, t) dt.$$

Through the discrete Laplacian  $\Delta_h$ , we can rewrite the fully discrete scheme (2.16)-(2.17) in the following equivalent forms:

$$\begin{aligned} \bar{\partial}_\tau \psi_n^{(h)} - \Delta_h \bar{\partial}_\tau^{1-\alpha} (\psi_n^{(h)} - \psi_0^{(h)}) &= P_h f_n + \sigma P_h \bar{\partial}_\tau W^M(\cdot, t_n), \quad \text{if } \alpha \in (0, 1], \\ \bar{\partial}_\tau \psi_n^{(h)} - \Delta_h \bar{\partial}_\tau^{1-\alpha} \psi_n^{(h)} &= P_h f_n + \sigma P_h \bar{\partial}_\tau W^M(\cdot, t_n), \quad \text{if } \alpha \in (1, 2). \end{aligned} \quad (2.18)$$

Note that  $\psi_0^{(h)} \in L^2(\Omega; X_h)$  and  $P_h f_n + \sigma P_h \bar{\partial}_\tau W^M(\cdot, t_n) \in L^2(\Omega; X_h)$  for  $n = 1, 2, \dots, N$ . If the numerical solutions  $\psi_n^{(h)} \in L^2(\Omega; X_h)$ ,  $n = 0, 1, \dots, m-1$ , then we define

$$g_m^{(h)} := \begin{cases} \tau^{\alpha-1} \sum_{j=1}^{m-1} b_{m-j} \Delta_h (\psi_j^{(h)} - \psi_0^{(h)}) + P_h f_m + \sigma P_h \bar{\partial}_\tau W^M(\cdot, t_m) & \text{if } \alpha \in (0, 1], \\ \tau^{\alpha-1} \sum_{j=1}^{m-1} b_{m-j} \Delta_h \psi_j^{(h)} + P_h f_m + \sigma P_h \bar{\partial}_\tau W^M(\cdot, t_m) & \text{if } \alpha \in (1, 2). \end{cases}$$

Then  $g_m^{(h)} \in L^2(\Omega; X_h)$ , and the numerical solution defined by (2.18) is given by

$$\psi_m^{(h)} = \begin{cases} \psi_0^{(h)} + (\tau^{-1} - \tau^{\alpha-1} b_0 \Delta_h)^{-1} (\tau^{-1} (\psi_{m-1}^{(h)} - \psi_0^{(h)}) + g_m^{(h)}) & \text{if } \alpha \in (0, 1), \\ (\tau^{-1} - \tau^{\alpha-1} b_0 \Delta_h)^{-1} (\tau^{-1} \psi_{m-1}^{(h)} + g_m^{(h)}) & \text{if } \alpha \in (1, 2), \end{cases}$$

which is well defined in  $L^2(\Omega; X_h)$ . By induction, the numerical solutions  $\psi_n^{(h)} \in L^2(\Omega; X_h)$ ,  $n = 1, 2, \dots, N$ , are well defined.

The main result of this paper is the following theorem.

**Theorem 1** *Let  $\alpha \in (0, \frac{2}{d})$  with  $d \in \{1, 2, 3\}$ ,  $f \in L^{\frac{4}{2+\alpha d}, 1}(0, T; L^2(\mathcal{O}))$  and  $\psi_0 \in \dot{H}^\chi(\mathcal{O})$ , with the notation*

$$\chi = \min \left( 2 - \frac{d}{2}, \frac{1}{\alpha} - \frac{d}{2} \right), \quad (2.19)$$

*and assume that the spatial mesh size satisfies  $0 < h < h_0$  for some constant  $h_0$ . Then the numerical solution given by (2.18) converges to the mild solution of (1.1) with sharp order of convergence, i.e.,*

$$\mathbb{E} \|\psi(\cdot, t_n) - \psi_n^{(h)}\|_{L^2(\mathcal{O})} \leq \begin{cases} C(\sigma + \Upsilon(\psi_0, f)) (\tau^{\frac{1}{2} - \frac{\alpha d}{4}} + \ell_h^{\frac{1}{2}} h^{\frac{1}{\alpha} - \frac{d}{2}}) & \text{if } \alpha \in [\frac{1}{2}, \frac{2}{d}), \\ C(\sigma + \Upsilon(\psi_0, f)) (\tau^{\frac{1}{2} - \frac{\alpha d}{4}} + h^{2 - \frac{d}{2}}) & \text{if } \alpha \in (0, \frac{1}{2}), \end{cases}$$

where

$$\Upsilon(\psi_0, f) = \|\psi_0\|_{\dot{H}^\chi(\mathcal{O})} + \|f\|_{L^{\frac{4}{2+\alpha d}, 1}(0, T; L^2(\mathcal{O}))},$$

$\mathbb{E}$  denotes the expectation operator,  $\ell_h = \ln(e + 1/h)$ , the constant  $C$  is independent of  $h, \tau, n, \psi_0$ , and  $f$  (but may depend on  $T$  and  $h_0$ ).

**Proof.** Without loss of generality, we can assume  $\sigma = 1$  in the proof of Theorem 1. The solution of (1.1) can be decomposed into the solution of the deterministic problem

$$\begin{cases} \partial_t v(x, t) - \Delta \partial_t^{1-\alpha} v(x, t) = f(x, t) & (x, t) \in \mathcal{O} \times \mathbb{R}_+ \\ \partial_t^{1-\alpha} v(x, t) = 0 & (x, t) \in \partial\mathcal{O} \times \mathbb{R}_+ \\ v(x, 0) = \psi_0(x) & x \in \mathcal{O} \end{cases} \quad (2.20)$$

plus the solution of the stochastic problem

$$\begin{cases} \partial_t u(x, t) - \Delta \partial_t^{1-\alpha} u(x, t) = \dot{W}(x, t) & (x, t) \in \mathcal{O} \times \mathbb{R}_+ \\ \partial_t^{1-\alpha} u(x, t) = 0 & (x, t) \in \partial\mathcal{O} \times \mathbb{R}_+ \\ u(x, 0) = 0 & x \in \mathcal{O}. \end{cases} \quad (2.21)$$

Similarly, the solution of (2.18) can be decomposed into the solution of the deterministic finite element equation

$$\begin{cases} \bar{\partial}_\tau v_n^{(h)} - \Delta_h \bar{\partial}_\tau^{1-\alpha} v_n^{(h)} - v_0^{(h)} = P_h f_n & \text{if } \alpha \in (0, 1], \\ v_0^{(h)} = P_h \psi_0 \end{cases} \quad (2.22)$$

or

$$\begin{cases} \bar{\partial}_\tau v_n^{(h)} - \Delta_h \bar{\partial}_\tau^{1-\alpha} v_n^{(h)} = P_h f_n & \text{if } \alpha \in (1, 2), \\ v_0^{(h)} = P_h \psi_0 \end{cases} \quad (2.23)$$

plus the solution of the stochastic finite element equation

$$\begin{cases} \bar{\partial}_\tau u_n^{(h)} - \Delta_h \bar{\partial}_\tau^{1-\alpha} u_n^{(h)} = P_h \bar{\partial}_\tau W^M(\cdot, t_n) \\ u_0^{(h)} = 0. \end{cases} \quad (2.24)$$

In the next two sections, we prove Theorem 1 by estimating  $\mathbb{E}\|u(\cdot, t_n) - u_n^{(h)}\|$  and  $\|v(\cdot, t_n) - v_n^{(h)}\|$  separately. In particular, Theorem 1 follows from (3.1) and (4.1) (in (4.1), we have  $\chi = \frac{1}{\alpha} - \frac{d}{2}$  for  $\alpha \in [\frac{1}{2}, \frac{2}{d})$  and  $\chi = 2 - \frac{d}{2}$  for  $\alpha \in (0, \frac{1}{2})$ ).

### 3 Stochastic problem: estimate of $\mathbb{E}\|u(\cdot, t_n) - u_n^{(h)}\|$

In this section, we prove the following error estimate for the solutions of (2.21) and (2.24):

$$\mathbb{E}\|u(\cdot, t_n) - u_n^{(h)}\| \leq \begin{cases} C(\tau^{\frac{1}{2} - \frac{\alpha d}{4}} + \ell_h^{\frac{1}{2}} h^{\frac{1}{\alpha} - \frac{d}{2}}) & \text{if } \alpha \in [\frac{1}{2}, \frac{2}{d}), \\ C(\tau^{\frac{1}{2} - \frac{\alpha d}{4}} + h^{2 - \frac{d}{2}}) & \text{if } \alpha \in (0, \frac{1}{2}). \end{cases} \quad (3.1)$$

To this end, we introduce a time-discrete system of PDEs:

$$\begin{cases} \bar{\partial}_\tau u_n - \Delta \bar{\partial}_\tau^{1-\alpha} u_n = \bar{\partial}_\tau W(\cdot, t_n), & n = 1, 2, \dots, N, \\ u_0 = 0. \end{cases} \quad (3.2)$$

Then (2.24) can be viewed as the spatially finite element discretization of (3.2), and the error can be decomposed into two parts:

$$\mathbb{E}\|u(\cdot, t_n) - u_n^{(h)}\| \leq \mathbb{E}\|u(\cdot, t_n) - u_n\| + \mathbb{E}\|u_n - u_n^{(h)}\|,$$

where the first part on the right-hand side has been estimated in [12] (in [12] we have only considered zero initial condition  $u(\cdot, 0) = 0$ , and in this case the boundary condition  $u = 0$  on  $\partial\Omega$  is equivalent to  $\partial_t^{1-\alpha}u = 0$  on  $\partial\Omega$ ), i.e.,

$$\mathbb{E}\|u(\cdot, t_n) - u_n\| \leq C\tau^{\frac{1}{2} - \frac{\alpha d}{4}} \quad \forall \alpha \in (0, 2/d), \quad d \in \{1, 2, 3\}.$$

It remains to prove the following estimate in the next three subsections:

$$\mathbb{E}\|u_n - u_n^{(h)}\| \leq \begin{cases} C\ell_h^{\frac{1}{2}} h^{\frac{1}{\alpha} - \frac{d}{2}} & \text{if } \alpha \in \left[\frac{1}{2}, \frac{2}{d}\right), \\ Ch^{2 - \frac{d}{2}} & \text{if } \alpha \in \left(0, \frac{1}{2}\right). \end{cases} \quad (3.3)$$

### 3.1 Integral representations

We estimate  $\mathbb{E}\|u_n - u_n^{(h)}\|$  by using integral representations of  $u_n$  and  $u_n^{(h)}$ , respectively. We first introduce some notations:

$$\begin{aligned} \Gamma_{\theta, \kappa} &= \{z \in \mathbb{C} : |z| = \kappa, |\arg z| \leq \theta\} \cup \{z \in \mathbb{C} : z = \rho e^{\pm i\theta}, \rho \geq \kappa\}, \\ \Gamma_{\theta, \kappa}^{(\tau)} &= \left\{z \in \Gamma_{\theta, \kappa} : |\operatorname{Im}(z)| \leq \frac{\pi}{\tau}\right\}, \end{aligned}$$

which are contours on the complex plane, oriented with increasing imaginary parts. On the truncated contour  $\Gamma_{\theta, \kappa}^{(\tau)}$ , the following estimates hold.

**Lemma 1** ([12]) *Let  $\alpha \in (0, \frac{2}{d})$ ,  $\theta \in (\frac{\pi}{2}, \operatorname{arccot}(-\frac{2}{\pi}))$  and  $\kappa = \frac{1}{T}$  be given, with  $\delta_\tau(\zeta)$  defined in (2.7). Then*

$$\delta_\tau(e^{-z\tau}) \in \Sigma_\theta \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)} \quad (3.4)$$

$$C_0|z| \leq |\delta_\tau(e^{-z\tau})| \leq C_1|z| \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)} \quad (3.5)$$

$$|\delta_\tau(e^{-z\tau}) - z| \leq C\tau|z|^2 \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)} \quad (3.6)$$

$$|\delta_\tau(e^{-z\tau})^\alpha - z^\alpha| \leq C\tau|z|^{\alpha+1} \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)}, \quad (3.7)$$

where  $\Sigma_\theta := \{z \in \mathbb{C} \setminus \{0\} : |\arg z| \leq \theta < \pi\}$ , the constants  $C_0, C_1$  and  $C$  are independent of  $\tau$  and  $\kappa$ .

Let  $\bar{\partial}_\tau W$  denote a piecewise constant function in time, defined by

$$\begin{aligned} \bar{\partial}_\tau W(\cdot, t_0) &:= 0 \\ \bar{\partial}_\tau W(\cdot, t) &:= \frac{W(\cdot, t_n) - W(\cdot, t_{n-1})}{\tau} \quad \text{for } t \in (t_{n-1}, t_n], \quad n = 1, 2, \dots, N. \end{aligned}$$

Similarly, we define

$$\begin{aligned} \bar{\partial}_\tau W_j(t_0) &:= 0 \\ \bar{\partial}_\tau W_j(t) &:= \frac{W_j(t_n) - W_j(t_{n-1})}{\tau} \quad \text{for } t \in (t_{n-1}, t_n], \quad n = 1, 2, \dots, N. \end{aligned}$$

Then the following results hold.

**Lemma 2** *Let  $\alpha \in (0, \frac{2}{d})$  and  $\delta_\tau(\zeta)$  be defined as in (2.7) with the parameters  $\kappa$  and  $\theta$  satisfying the conditions of Lemma 1.*



(1) The solution of the time-discrete problem (3.2) can be represented by

$$u_n = \int_0^{t_n} E_\tau(t_n - s) \bar{\partial}_\tau W(\cdot, s) ds = \sum_{j=1}^{\infty} \int_0^{t_n} E_\tau(t_n - s) \phi_j \bar{\partial}_\tau W_j(s) ds, \quad (3.8)$$

where the operator  $E_\tau(t)$  is given by

$$E_\tau(t)\phi := \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt} \frac{z\tau}{e^{z\tau} - 1} \delta_\tau(e^{-z\tau})^{\alpha-1} (\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1} \phi dz \quad (3.9)$$

for  $\phi \in L^2(\mathcal{O})$ .

(2) The solution of the fully discrete problem (2.24) can be represented by

$$u_n^{(h)} = \sum_{j=1}^M \int_0^{t_n} E_\tau^{(h)}(t_n - s) \phi_j \bar{\partial}_\tau W_j(s) ds, \quad (3.10)$$

where the operator  $E_\tau^{(h)}(t)$  is given by

$$E_\tau^{(h)}(t)\phi := \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt} \frac{z\tau}{e^{z\tau} - 1} \delta_\tau(e^{-z\tau})^{\alpha-1} (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1} P_h \phi dz \quad (3.11)$$

for  $\phi \in L^2(\mathcal{O})$ .

The first statement in Lemma 2 has been proved in [12, Proposition 3.1]. The second statement can be proved in the same way, replacing the operator  $\Delta$  by  $\Delta_h$  and  $W(\cdot, t)$  by  $W^M(\cdot, t)$  (this does not affect the proof therein). From Lemma 2, we see that

$$\begin{aligned} u_n - u_n^{(h)} &= \sum_{j=1}^M \int_0^{t_n} (E_\tau(t_n - s) - E_\tau^{(h)}(t_n - s)) \phi_j \bar{\partial}_\tau W_j(s) ds \\ &\quad + \sum_{j=M+1}^{\infty} \int_0^{t_n} E_\tau(t_n - s) \phi_j \bar{\partial}_\tau W_j(s) ds \\ &=: \mathcal{I}_n + \mathcal{J}_n. \end{aligned} \quad (3.12)$$

We present the estimates for  $\mathcal{I}_n$  and  $\mathcal{J}_n$  in subsections 3.2 and 3.3, respectively.

### 3.2 Estimate of $\mathcal{I}_n$

Now, we start to estimate  $\mathcal{I}_n$ , i.e., the error of space discretization. The following lemmas are useful in the estimates of  $\mathcal{I}_n$  and  $\mathcal{J}_n$ .

**Lemma 3** ([18, 19, 26]) *Let  $\mathcal{O}$  denote a bounded domain in  $\mathbb{R}^d$ ,  $d \in \{1, 2, 3\}$ . Suppose  $\lambda_j$  denotes the  $j^{\text{th}}$  eigenvalue of the Dirichlet boundary problem for the Laplacian operator  $-\Delta$  in  $\mathcal{O}$ . Then, we have*

$$C_0^* j^{\frac{2}{d}} \leq \lambda_j \leq C_1^* j^{\frac{2}{d}} \quad (3.13)$$

for all  $j \geq 1$ , where the constants  $C_0^*$  and  $C_1^*$  are independent of  $j$ .

**Proof.** The well-known Weyl's law gives the asymptotic behavior of the eigenvalues of the Laplacian operator (see [19] and [26, pp. 322]):

$$\lim_{j \rightarrow \infty} \frac{\lambda_j}{j^{2/d}} = (2\pi)^2 (B_d |\mathcal{O}|)^{-\frac{2}{d}},$$

where  $B_d$  denotes the volume of the unit  $d$ -ball. The estimate (3.13) follows immediately from the above result.  $\square$

The following lemma is contained in [12, Lemma 3.2].

**Lemma 4** For any  $z \in \Sigma_\varphi$  with  $\varphi \in (0, \pi)$ , we have

$$\left| \frac{1}{z + \lambda_j} \right| \leq \frac{C_\varphi}{|z| + \lambda_j}, \quad (3.14)$$

where  $j = 1, 2, \dots$ .

The following resolvent estimates will be frequently used in this paper.

**Lemma 5** For  $z \in \Sigma_\theta$  (see the definition in Lemma 1), with  $\theta \in (0, \pi)$ , we have the following resolvent estimates:

$$\|(z - \Delta)^{-1}\| + \|(z - \Delta_h)^{-1}\| \leq C|z|^{-1}, \quad (3.15)$$

$$\|\Delta^{1-\gamma}(z - \Delta)^{-1}\| + \|\Delta_h^{1-\gamma}(z - \Delta_h)^{-1}\| \leq C|z|^{-\gamma}, \quad \gamma \in [0, 1]. \quad (3.16)$$

**Proof.** The first inequality is due to the self-adjointness and nonnegativity of the operators  $-\Delta$  and  $-\Delta_h$ . These properties guarantee that  $\Delta$  and  $\Delta_h$  generate a bounded analytic semigroup of angle  $\frac{\pi}{2}$  on  $L^2(\mathcal{O})$  and  $(X_h, \|\cdot\|_{L^2(\mathcal{O})})$ , respectively; see [3, Example 3.7.5 and Theorem 3.7.11].

Recall that  $\lambda_j$  and  $\phi_j$ ,  $j = 1, 2, \dots$ , are the eigenvalues and eigenfunctions of the operator  $-\Delta$  (see Section 2.1). The second inequality is due to the interpolation inequality

$$\begin{aligned} \|\Delta^{1-\gamma}\varphi\| &= \left\| \sum_{j=1}^{\infty} \lambda_j^{1-\gamma}(\phi_j, \varphi)\phi_j \right\| \\ &= \left( \sum_{j=1}^{\infty} \lambda_j^{2-2\gamma} |(\phi_j, \varphi)|^2 \right)^{\frac{1}{2}} = \left( \sum_{j=1}^{\infty} |(\phi_j, \varphi)|^{2\gamma} \lambda_j^{2-2\gamma} |(\phi_j, \varphi)|^{2-2\gamma} \right)^{\frac{1}{2}} \\ &\leq \left( \sum_{j=1}^{\infty} |(\phi_j, \varphi)|^2 \right)^{\frac{\gamma}{2}} \left( \sum_{j=1}^{\infty} \lambda_j^2 |(\phi_j, \varphi)|^2 \right)^{\frac{1-\gamma}{2}} \quad (\text{use Hölder's inequality}) \\ &\leq \|\varphi\|^\gamma \|\Delta\varphi\|^{1-\gamma}. \end{aligned}$$

Substituting  $\varphi = (z - \Delta)^{-1}\phi$  into the inequality above yields

$$\begin{aligned} \|\Delta^{1-\gamma}(z - \Delta)^{-1}\phi\| &\leq \|(z - \Delta)^{-1}\phi\|^\gamma \|\Delta(z - \Delta)^{-1}\phi\|^{1-\gamma} \\ &\leq (C|z|^{-1}\|\phi\|)^\gamma (C\|\phi\|)^{1-\gamma} \\ &\leq C|z|^{-\gamma}\|\phi\|. \end{aligned}$$

This proves the first part of (3.16). The estimate of  $\|\Delta_h^{1-\gamma}(z - \Delta_h)^{-1}\|$  can be proved similarly (by using the eigenvalues and eigenfunctions of  $-\Delta_h$ ).  $\square$

The following lemma is concerned with the difference between the continuous and discrete resolvent operators.

**Lemma 6** Let  $\alpha \in (0, \frac{2}{d})$  and  $\delta_\tau(\zeta)$  be defined as in (2.7) with the parameters  $\kappa$  and  $\theta$  satisfying the conditions of Lemma 1. Then we have

$$\|[(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1} - (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}P_h]\phi_j\| \leq Ch^{2\varepsilon}(|z|^\alpha + \lambda_j)^{-(1-\varepsilon)} \quad (3.17)$$

for all  $\varepsilon \in [0, 1]$  and  $j = 1, 2, \dots, M$ .

**Proof.** First, (3.4) yields that  $\delta_\tau(e^{-z\tau})^\alpha \in \Sigma_{\alpha\theta}$  for  $z \in \Gamma_{\theta, \kappa}^{(\tau)}$ . Consequently, we have (cf. [22, pp. 7])

$$\|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1} - (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}P_h\| \leq Ch^2. \quad (3.18)$$

Second, by Lemma 1 and Lemma 5, there exists a constant  $C$  which depends only on  $\theta$  and  $\alpha$  such that

$$\|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\| \leq C|\delta_\tau(e^{-z\tau})|^{-\alpha} \leq C|z|^{-\alpha} \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)}, \quad (3.19)$$

$$\|(\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}P_h\| \leq \|(\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}\| \|P_h\| \leq C|z|^{-\alpha} \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)}. \quad (3.20)$$

Since  $(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\Delta = (\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\delta_\tau(e^{-z\tau})^\alpha - I$ , it follows that

$$\|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\Delta\| \leq C.$$

Consequently, we have

$$\begin{aligned} \|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\phi_j\| &= \|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\Delta\Delta^{-1}\phi_j\| \\ &\leq \|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\Delta\| \|\Delta^{-1}\phi_j\| \\ &\leq C\lambda_j^{-1}, \end{aligned}$$

which together with  $h^2 \leq C\lambda_M^{-1}$  (since  $M = [h^{-d}] + 1 \sim h^{-d}$  and  $\lambda_M \leq CM^{\frac{2}{d}}$  by Lemma 3) implies

$$\begin{aligned} &\|(\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}P_h\phi_j\| \\ &\leq \|(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1}\phi_j\| + \|[(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1} - (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}]P_h\phi_j\| \\ &\leq C\lambda_j^{-1} + Ch^2 \\ &\leq C\lambda_j^{-1} \end{aligned} \quad (3.21)$$

for  $j = 1, \dots, M$ . Therefore, (3.19)-(3.21) leads to

$$\begin{aligned} \|[ (\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1} - (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}P_h ]\phi_j\| &\leq C \min\{|z|^{-\alpha}, \lambda_j^{-1}\} \\ &\leq C(|z|^\alpha + \lambda_j)^{-1}. \end{aligned} \quad (3.22)$$

Finally, interpolation between (3.18) and (3.22) yields (3.17). This completes the proof of Lemma 6.  $\square$

Now, we turn to the estimate of  $\mathcal{I}_n$ . From Lemma 6, and choosing  $\beta \in (0, 1)$ , it is easy to derive

$$\begin{aligned} &\|(E_\tau(t) - E_\tau^{(h)}(t))\phi_j\|^2 \\ &\leq \left( \int_{\Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt}| |\delta_\tau(e^{-z\tau})|^{\alpha-1} \|[(\delta_\tau(e^{-z\tau})^\alpha - \Delta)^{-1} - (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1}P_h]\phi_j\| |dz| \right)^2 \\ &\leq \left( C \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} |z|^{\alpha-1} (|z|^\alpha + \lambda_j)^{-(1-\varepsilon)} h^{2\varepsilon} |dz| \right)^2 \\ &\leq Ch^{4\varepsilon} \left( \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|dz|}{|z|^\beta} \right) \left( \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|z|^{2\alpha-2+\beta}}{(|z|^\alpha + \lambda_j)^{2-2\varepsilon}} |dz| \right) \\ &\leq Ch^{4\varepsilon} t^{\beta-1} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|z|^{2\alpha-2+\beta}}{(|z|^\alpha + \lambda_j)^{2-2\varepsilon}} |dz|, \end{aligned}$$

where  $|dz|$  denotes the arc length element. By choosing  $\kappa = \frac{1}{t}$  we have

$$\begin{aligned} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|dz|}{|z|^\beta} &\leq \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} e^{-rt|\cos(\theta)|} \frac{dr}{r^\beta} + \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} \frac{d\varphi}{\kappa^{\beta-1}} \\ &\leq Ct^{\beta-1}. \end{aligned} \quad (3.23)$$

Since  $\frac{W_j(t_i) - W_j(t_{i-1})}{\tau}$ ,  $i = 1, \dots, n$ , and  $j = 1, 2, \dots$ , are stochastically independent of each other, it follows that

$$\mathbb{E}\|\mathcal{I}_n\|^2 = \sum_{j=1}^M \mathbb{E} \left\| \sum_{i=1}^n \frac{W_j(t_i) - W_j(t_{i-1})}{\tau} \int_{t_{i-1}}^{t_i} (E_\tau(t_n - s) - E_\tau^{(h)}(t_n - s)) \phi_j ds \right\|^2$$

$$\begin{aligned}
&= \sum_{j=1}^M \sum_{i=1}^n \frac{1}{\tau} \left\| \int_{t_{i-1}}^{t_i} (E_\tau(t_n - s) - E_\tau^{(h)}(t_n - s)) \phi_j ds \right\|^2 \\
&\leq \sum_{j=1}^M \sum_{i=1}^n \int_{t_{i-1}}^{t_i} \|E_\tau(t_n - s) - E_\tau^{(h)}(t_n - s)\| \phi_j\|^2 ds \quad (\text{Cauchy-Schwarz inequality}) \\
&= \sum_{j=1}^M \int_0^{t_n} \|(E_\tau(t) - E_\tau^{(h)}(t)) \phi_j\|^2 dt \\
&\leq Ch^{4\varepsilon} \int_0^{t_n} t^{\beta-1} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \sum_{j=1}^M \frac{|z|^{2\alpha-2+\beta}}{(|z|^\alpha + \lambda_j)^{2-2\varepsilon}} |dz| dt \\
&= Ch^{4\varepsilon} \int_0^{t_n} t^{\beta-1} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} |z|^{2\varepsilon\alpha-2+\beta} \sum_{j=1}^M \left( \frac{|z|^\alpha}{|z|^\alpha + \lambda_j} \right)^{2-2\varepsilon} |dz| dt.
\end{aligned} \tag{3.24}$$

Since  $\left(\frac{|z|^\alpha}{|z|^\alpha + \lambda_j}\right)^{2-2\varepsilon} \leq 1$  for  $\varepsilon \in [0, 1]$ , it follows that

$$\sum_{j=1}^M \left( \frac{|z|^\alpha}{|z|^\alpha + \lambda_j} \right)^{2-2\varepsilon} \leq M \leq Ch^{-d},$$

where the last inequality is due to our choice  $M = [h^{-d}] + 1$ . Thus (3.24) reduces to

$$\begin{aligned}
\mathbb{E}\|\mathcal{I}_n\|^2 &\leq Ch^{4\varepsilon-d} \int_0^{t_n} t^{\beta-1} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} |z|^{2\varepsilon\alpha-2+\beta} |dz| dt \\
&\leq Ch^{4\varepsilon-d} \int_0^{t_n} t^{\beta-1} \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} e^{-rt|\cos(\theta)|} r^{2\varepsilon\alpha-2+\beta} dr dt \\
&\quad + Ch^{4\varepsilon-d} \int_0^{t_n} t^{\beta-1} \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} \kappa^{2\varepsilon\alpha-1+\beta} d\varphi dt \\
&= Ch^{4\varepsilon-d} \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} r^{2\varepsilon\alpha-2+\beta} \int_0^{t_n} t^{\beta-1} e^{-rt|\cos(\theta)|} dt dr \\
&\quad + Ch^{4\varepsilon-d} \int_0^{t_n} t^{\beta-1} \kappa^{2\varepsilon\alpha-1+\beta} \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} d\varphi dt \\
&\leq Ch^{4\varepsilon-d} \left( \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} r^{2\varepsilon\alpha-2} dr + \kappa^{2\varepsilon\alpha-1+\beta} \int_0^{t_n} t^{\beta-1} dt \right) \\
&\leq Ch^{4\varepsilon-d} \left( \frac{1}{1-2\varepsilon\alpha} \kappa^{2\varepsilon\alpha-1} - \frac{1}{1-2\varepsilon\alpha} \left( \frac{\pi}{\tau \sin(\theta)} \right)^{2\varepsilon\alpha-1} + \kappa^{2\varepsilon\alpha-1} (\kappa t_n)^\beta \right).
\end{aligned}$$

For  $\alpha \in [\frac{1}{2}, \frac{d}{2})$ , we choose  $\varepsilon = \frac{1-1/\ell_h}{2\alpha}$  with  $\ell_h = \ln(e+1/h)$  and  $h^{-1/\ell_h} \leq e$  (can recall that  $\kappa = \frac{1}{T}$ ), we have

$$\mathbb{E}\|\mathcal{I}_n\|^2 \leq CT^{\frac{1}{\ell_h}} \ell_h h^{\frac{2}{\alpha}-d}. \tag{3.25}$$

For  $\alpha \in (0, \frac{1}{2})$ , we choose  $\varepsilon = 1$  and get

$$\mathbb{E}\|\mathcal{I}_n\|^2 \leq CT^{1-2\alpha} h^{4-d}. \tag{3.26}$$

### 3.3 Estimate of $\mathcal{J}_n$

In this subsection, we present the estimate of  $\mathcal{J}_n$  in (3.12), completing the proof of (3.1). In view of the definition of  $E_\tau(t)$  in (3.9), by using Lemma 1 and (3.14) to estimate  $|\delta_\tau(e^{-z\tau})^{\alpha-1}|$  and  $|(\delta_\tau(e^{-z\tau})^\alpha + \lambda_j)^{-1}|$ , we obtain

$$\begin{aligned}
\|E_\tau(t)\phi_j\|^2 &= \left| \frac{1}{2\pi i} \int_{\Gamma_{\theta,\kappa}^{(\tau)}} e^{zt} \frac{z\tau}{e^{z\tau} - 1} \delta_\tau(e^{-z\tau})^{\alpha-1} (\delta_\tau(e^{-z\tau})^\alpha + \lambda_j)^{-1} dz \right|^2 \\
&\leq C \left( \int_{\Gamma_{\theta,\kappa}^{(\tau)}} |e^{zt}| \left| \frac{z\tau}{e^{z\tau} - 1} \right| |\delta_\tau(e^{-z\tau})|^{\alpha-1} (|\delta_\tau(e^{-z\tau})|^\alpha + \lambda_j)^{-1} |dz| \right)^2 \\
&\leq C \left( \int_{\Gamma_{\theta,\kappa}^{(\tau)}} |e^{zt}| |z|^{\alpha-1} (|z|^\alpha + \lambda_j)^{-1} |dz| \right)^2 \quad (\text{use (C.1) here}) \\
&\leq C \left( \int_{\Gamma_{\theta,\kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|dz|}{|z|^\beta} |dz| \right) \left( \int_{\Gamma_{\theta,\kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|z|^{2\alpha-2+\beta}}{(|z|^\alpha + \lambda_j)^2} |dz| \right) \\
&\leq Ct^{\beta-1} \int_{\Gamma_{\theta,\kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|z|^{2\alpha-2+\beta}}{(|z|^{\frac{d\alpha}{2}} + j)^{\frac{4}{d}}} |dz|, \tag{3.27}
\end{aligned}$$

where we have used (3.23). In view of (3.12), we have

$$\begin{aligned}
\mathbb{E}\|\mathcal{J}_n\|^2 &\leq \int_0^{t_n} \sum_{j=M+1}^{\infty} \|E_\tau(t)\phi_j\|^2 dt \\
&\leq C \int_0^{t_n} t^{\beta-1} \int_{\Gamma_{\theta,\kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \sum_{j=M+1}^{\infty} \frac{|z|^{2\alpha-2+\beta}}{(|z|^{\frac{d\alpha}{2}} + j)^{\frac{4}{d}}} |dz| dt \\
&\leq C \int_0^{t_n} t^{\beta-1} \int_{\Gamma_{\theta,\kappa}^{(\tau)}} e^{|z|t \cos(\arg(z))} \frac{|z|^{2\alpha-2+\beta}}{(|z|^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} |dz| dt \\
&\leq C \int_0^{t_n} t^{\beta-1} \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} e^{-rt|\cos(\theta)|} \frac{r^{2\alpha-2+\beta}}{(r^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} dr dt \\
&\quad + C \int_0^{t_n} t^{\beta-1} \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} \frac{\kappa^{2\alpha-1+\beta}}{(\kappa^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} d\varphi dt \\
&= C \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} \frac{r^{2\alpha-2+\beta}}{(r^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} \int_0^{t_n} t^{\beta-1} e^{-rt|\cos(\theta)|} dt dr \\
&\quad + C \int_0^{t_n} t^{\beta-1} \frac{\kappa^{2\alpha-1+\beta}}{(\kappa^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} d\varphi dt \\
&\leq C \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} \frac{r^{2\alpha-2}}{(r^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} dr + \frac{C(\kappa t_n)^\beta \kappa^{2\alpha-1}}{(\kappa^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} \\
&\leq C \left( \int_{\kappa}^{\min\{M^{\frac{2}{d\alpha}}, \frac{\pi}{\tau \sin(\theta)}\}} \frac{r^{2\alpha-2}}{M^{\frac{4}{d}-1}} dr + \int_{\min\{M^{\frac{2}{d\alpha}}, \frac{\pi}{\tau \sin(\theta)}\}}^{\frac{\pi}{\tau \sin(\theta)}} \frac{r^{2\alpha-2}}{r^{2\alpha-\frac{d\alpha}{2}}} dr \right) \\
&\quad + \frac{C\kappa^{2\alpha-1}}{(\kappa^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}} \\
&\leq C \left( \int_{\kappa}^{\min\{M^{\frac{2}{d\alpha}}, \frac{\pi}{\tau \sin(\theta)}\}} \frac{r^{2\alpha-2}}{M^{\frac{4}{d}-1}} dr + M^{1-\frac{2}{d\alpha}} \right) + \frac{C\kappa^{2\alpha-1}}{(\kappa^{\frac{d\alpha}{2}} + M)^{\frac{4}{d}-1}}. \tag{3.28}
\end{aligned}$$

We estimate  $\mathbb{E}\|\mathcal{J}_n\|^2$  in three different cases by using  $M = [h^{-d}] + 1 \leq Ch^{-d}$ .

In the case  $\alpha \in (\frac{1}{2}, \frac{2}{d})$ , (3.28) reduces to

$$\begin{aligned} \mathbb{E}\|\mathcal{J}_n\|^2 &\leq C \left( \int_{\kappa}^{M^{\frac{2}{d\alpha}}} \frac{r^{2\alpha-2}}{M^{\frac{4}{d}-1}} dr + M^{1-\frac{2}{d\alpha}} \right) + CM^{1-\frac{4}{d}} \\ &\leq CM^{1-\frac{2}{d\alpha}} \leq Ch^{\frac{2}{\alpha}-d}. \end{aligned} \quad (3.29)$$

In the case  $\alpha \in (0, \frac{1}{2})$ , (3.28) yields

$$\begin{aligned} \mathbb{E}\|\mathcal{J}_n\|^2 &\leq C(T^{1-2\alpha}M^{1-\frac{4}{d}} + M^{1-\frac{2}{d\alpha}}) \leq C(1 + T^{1-2\alpha})M^{1-\frac{4}{d}} \\ &\leq C(1 + T^{1-2\alpha})h^{4-d}. \end{aligned} \quad (3.30)$$

In the case  $\alpha = \frac{1}{2}$ , (3.28) implies

$$\begin{aligned} \mathbb{E}\|\mathcal{J}_n\|^2 &\leq C \left( \int_{\kappa}^{M^{\frac{2}{d\alpha}}} \frac{r^{-1}}{M^{\frac{4}{d}-1}} dr + M^{1-\frac{2}{d\alpha}} \right) + CM^{1-\frac{4}{d}} \leq C(\ln M + \ln T)M^{1-\frac{4}{d}} \\ &\leq C(\ell_h + \ln T)h^{\frac{2}{\alpha}-d}. \end{aligned} \quad (3.31)$$

It is well-known that  $(\mathbb{E}\|u_n - u_n^{(h)}\|)^2 \leq \mathbb{E}\|u_n - u_n^{(h)}\|^2$ , substituting (3.25)-(3.26) and (3.29)-(3.31) into (3.12) yields (3.3), completing the proof of (3.1).

#### 4 Deterministic problem: estimate of $\|v(\cdot, t_n) - v_n^{(h)}\|$

In this section, we estimate the error  $\|v(\cdot, t_n) - v_n^{(h)}\|$  by minimizing the regularity requirement on  $\psi_0$  and  $f$  to match the convergence rate proved in the last section, where  $v$  and  $v_n^{(h)}$  are the solutions of (2.20) and (2.22) or (2.23), respectively. In particular, we prove the following estimate for  $\alpha \in (0, \frac{2}{d})$ :

$$\|v(\cdot, t_n) - v_n^{(h)}\| \leq C(\|\psi_0\|_{\dot{H}^\chi(\mathcal{O})} + \|f\|_{L^{\frac{4}{2+\alpha d}, 1}(0, t_n; L^2(\mathcal{O}))})(\tau^{\frac{1}{2}-\frac{\alpha d}{4}} + h^\chi), \quad (4.1)$$

where  $\chi \in (0, 2 - \frac{d}{2})$  is defined in (2.19). To this end, we introduce the semi-discrete finite element problem

$$\begin{cases} \partial_t v^{(h)} - \Delta_h \partial_t^{1-\alpha} v^{(h)} = P_h f \\ v^{(h)}(\cdot, 0) = P_h \psi_0. \end{cases} \quad (4.2)$$

Then (2.22) or (2.23) can be viewed as the time discretization of (4.2), with the following decomposition:

$$\|v(\cdot, t_n) - v_n^{(h)}\| \leq \|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| + \|v^{(h)}(\cdot, t_n) - v_n^{(h)}\|.$$

We estimate  $\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\|$  and  $\|v^{(h)}(\cdot, t_n) - v_n^{(h)}\|$  in the next two subsections, respectively.

##### 4.1 Spatial discretization: estimate of $\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\|$

In this subsection we estimate  $\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\|$ , where  $v$  and  $v^{(h)}$  are solutions of (2.20) and (4.2), respectively. Next, we consider the following three cases:

Case 1:  $\psi_0 \neq 0$ ,  $f = 0$ ,  $\alpha \in (0, 1] \cap (0, \frac{2}{d})$ .

Case 2:  $\psi_0 \neq 0$ ,  $f = 0$ ,  $\alpha \in (1, \frac{2}{d})$ . (This requires  $d = 1$ , cf. Theorem 1)

Case 3:  $\psi_0 = 0$ ,  $f \neq 0$ .

In Case 1, the solutions of (2.20) and (4.2) are given by  $v(\cdot, t_n) = \psi_0$  and  $v^{(h)}(\cdot, t_n) = P_h \psi_0$ ,  $n = 1, 2, \dots$ , respectively (this follows from setting  $f = 0$  and  $\sigma = 0$  in (A.3) of Appendix A). Consequently, we have

$$\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| = \|\psi_0 - P_h \psi_0\| \leq C \|\psi_0\|_{\dot{H}^\chi(\mathcal{O})} h^\chi,$$

where we have used (2.10) in the last inequality.

In Case 2, we note that the PDE problem (2.20) is equivalent to (multiplying both sides by the operator  $\partial_t^{\alpha-1}$ )

$$\begin{cases} \partial_t^\alpha v(x, t) - \Delta v(x, t) = 0 & (x, t) \in \mathcal{O} \times \mathbb{R}_+ \\ v(x, t) = 0 & (x, t) \in \partial\mathcal{O} \times \mathbb{R}_+ \\ v(x, 0) = \psi_0(x) \text{ and } \partial_t v(x, 0) = 0 & x \in \mathcal{O} \end{cases} \quad (4.3)$$

and the finite element problem (4.2) is equivalent to (for the same reason)

$$\begin{cases} \partial_t^\alpha v^{(h)} - \Delta_h v^{(h)} = 0 \\ v^{(h)}(\cdot, 0) = P_h \psi_0 \text{ and } \partial_t v^{(h)}(\cdot, 0) = 0. \end{cases} \quad (4.4)$$

By Laplace transform,  $v = E(t)\psi_0$  and  $v^{(h)} = E^{(h)}(t)P_h\psi_0$ , where the operators  $E(t) : L^2(\mathcal{O}) \rightarrow L^2(\mathcal{O})$  and  $E^{(h)}(t) : X_h \rightarrow X_h$  are given in Appendix A. An error estimate for (4.3) and (4.4) was presented in [13, Theorems 3.2]:

$$\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| \leq C \|\psi_0\|_{\dot{H}^2(\mathcal{O})} h^2. \quad (4.5)$$

The boundedness of the solution operators  $E(t) : L^2(\mathcal{O}) \rightarrow L^2(\mathcal{O})$  and  $E^{(h)}(t) : X_h \rightarrow X_h$  (see Appendix A, (A.4) and (A.6)) implies

$$\begin{aligned} \|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| &= \|E(t_n)\psi_0 - E^{(h)}(t_n)P_h\psi_0\| \\ &\leq \|E(t_n)\psi_0\| + \|E^{(h)}(t_n)P_h\psi_0\| \\ &\leq C \|\psi_0\|. \end{aligned} \quad (4.6)$$

Then the interpolation between (4.5) and (4.6) yields

$$\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| \leq C \|\psi_0\|_{\dot{H}^\chi(\mathcal{O})} h^\chi. \quad (4.7)$$

In Case 3, we have

$$\begin{aligned} P_h v(\cdot, t_n) - v^{(h)}(\cdot, t_n) &= P_h \left( v(\cdot, t_n) - v^{(h)}(\cdot, t_n) \right) \\ &= P_h \left( \int_0^{t_n} E(t_n - s) f(\cdot, s) ds - \int_0^{t_n} E^{(h)}(t_n - s) P_h f(\cdot, s) ds \right) \\ &= P_h \int_0^{t_n} E(t_n - s) (f(\cdot, s) - P_h f(\cdot, s)) ds \\ &\quad + P_h \int_0^{t_n} (E(t_n - s) - E^{(h)}(t_n - s) P_h) P_h f(\cdot, s) ds \\ &=: \mathcal{I}^{(h)}(t_n) + \mathcal{J}^{(h)}(t_n), \end{aligned} \quad (4.8)$$

where

$$\begin{aligned} \mathcal{I}^{(h)}(t_n) &= \int_0^{t_n} \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{z(t_n-s)} z^{\alpha-1} P_h (z^\alpha - \Delta)^{-1} (f(\cdot, s) - P_h f(\cdot, s)) dz ds \\ &= \int_0^{t_n} \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{z(t_n-s)} z^{\alpha-1} P_h w_z(\cdot, s) dz ds \end{aligned} \quad (4.9)$$

with

$$w_z(\cdot, s) = (z^\alpha - \Delta)^{-1}(f(\cdot, s) - P_h f(\cdot, s)), \quad (4.10)$$

which satisfies

$$\begin{aligned} \|w_z(\cdot, s)\|_{\dot{H}^2(\mathcal{O})} &= \|\Delta^{-1}\Delta(z^\alpha - \Delta)^{-1}(f(\cdot, s) - P_h f(\cdot, s))\|_{\dot{H}^2(\mathcal{O})} \\ &\leq C\|\Delta(z^\alpha - \Delta)^{-1}(f(\cdot, s) - P_h f(\cdot, s))\| \\ &\leq C\|f(\cdot, s) - P_h f(\cdot, s)\|. \end{aligned} \quad (4.11)$$

By using the Ritz projection operator  $R_h$ , equation (4.10) implies

$$\begin{aligned} ((z^\alpha - \Delta_h)P_h w_z + \Delta_h(P_h w_z - R_h w_z), \phi_h) &= z^\alpha(w_z, \phi_h) + (\nabla R_h w_z, \nabla \phi_h) \\ &= z^\alpha(w_z, \phi_h) + (\nabla w_z, \nabla \phi_h) \\ &= ((z^\alpha - \Delta)w_z, \phi_h) \\ &= (f - P_h f, \phi_h) = 0 \quad \forall \phi_h \in X_h, \end{aligned}$$

i.e.,  $P_h w_z = -(z^\alpha - \Delta_h)^{-1}\Delta_h(P_h w_z - R_h w_z)$ . Consequently, we obtain

$$\begin{aligned} |z|^{\gamma\alpha}\|P_h w_z(\cdot, s)\| &= \||z|^{\gamma\alpha}(z^\alpha - \Delta_h)^{-\gamma}[\Delta_h(z^\alpha - \Delta_h)^{-1}]^{1-\gamma}\Delta_h^\gamma(P_h w_z(\cdot, s) - R_h w_z(\cdot, s))\| \\ &\leq C\|\Delta_h^\gamma(P_h w_z(\cdot, s) - R_h w_z(\cdot, s))\| \\ &\leq C\|w_z(\cdot, s)\|_{\dot{H}^2(\mathcal{O})}h^{2-2\gamma} \quad (\text{here we use (2.15)}) \\ &\leq C\|f(\cdot, s) - P_h f(\cdot, s)\|h^{2-2\gamma} \quad (\text{here we use (4.11)}) \\ &\leq C\|f(\cdot, s)\|h^{2-2\gamma}. \end{aligned}$$

By choosing  $\gamma = 1 - \frac{1}{2}\chi$  so that  $2 - 2\gamma = \chi$ , the inequality (4.9) reduces to

$$\begin{aligned} \|\mathcal{I}^{(h)}(t_n)\| &\leq C \int_0^{t_n} \int_{\Gamma_{\theta, \kappa}} |e^{z(t_n-s)}| |z|^{(1-\gamma)\alpha-1} |z|^{\gamma\alpha} \|P_h w(\cdot, s)\| |dz| ds \\ &\leq C \int_0^{t_n} \int_{\Gamma_{\theta, \kappa}} |e^{z(t_n-s)}| |z|^{\frac{1}{2}\chi\alpha-1} \|f(\cdot, s)\| h^\chi |dz| ds \\ &\leq Ch^\chi \int_0^{t_n} (t_n - s)^{-\frac{1}{2}\chi\alpha} \|f(\cdot, s)\| ds. \end{aligned} \quad (4.12)$$

Furthermore, applying the similar analysis in Lemma 6, we have

$$\begin{aligned} \|(z^\alpha - \Delta)^{-1} - (z^\alpha - \Delta_h)^{-1}P_h\| &\leq Ch^2, \quad (\text{cf. [22, pp. 7]}) \\ \|(z^\alpha - \Delta)^{-1} - (z^\alpha - \Delta_h)^{-1}P_h\| &\leq C|z|^{-\alpha}. \quad (\text{resolvent estimate, see (3.15)}) \end{aligned}$$

The interpolation of the last two inequalities yields

$$\|(z^\alpha - \Delta)^{-1} - (z^\alpha - \Delta_h)^{-1}P_h\| \leq C|z|^{-\gamma\alpha}h^{2-2\gamma} \quad \forall \gamma \in [0, 1].$$

Again, by choosing  $\gamma = 1 - \frac{1}{2}\chi$  (so that  $2 - 2\gamma = \chi$ ), we have

$$\begin{aligned} \|\mathcal{J}^{(h)}(t_n)\| &= \left\| P_h \int_0^{t_n} \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{z(t_n-s)} z^{\alpha-1} \left( (z^\alpha - \Delta)^{-1} - (z^\alpha - \Delta_h)^{-1}P_h \right) P_h f(\cdot, s) dz ds \right\| \\ &\leq \int_0^{t_n} \int_{\Gamma_{\theta, \kappa}} |e^{z(t_n-s)}| |z|^{\alpha-1} \|(z^\alpha - \Delta)^{-1} - (z^\alpha - \Delta_h)^{-1}P_h\| \|P_h f(\cdot, s)\| |dz| ds \\ &\leq C \int_0^{t_n} \int_{\Gamma_{\theta, \kappa}} |e^{z(t_n-s)}| |z|^{\alpha-1} |z|^{-(1-\frac{1}{2}\chi)\alpha} h^\chi \|f(\cdot, s)\| |dz| ds \end{aligned}$$



$$\begin{aligned}
&\leq Ch^\chi \int_0^{t_n} \left( \int_{\Gamma_{\theta,\kappa}} |e^{z(t_n-s)}| |z|^{\frac{1}{2}\chi\alpha-1} |dz| \right) \|f(\cdot, s)\| ds \\
&\leq Ch^\chi \int_0^{t_n} (t_n - s)^{-\frac{1}{2}\chi\alpha} \|f(\cdot, s)\| ds.
\end{aligned} \tag{4.13}$$

Substituting (4.12) and (4.13) into (4.8) yields

$$\|P_h v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| \leq Ch^\chi \int_0^{t_n} (t_n - s)^{-\frac{1}{2}\chi\alpha} \|f(\cdot, s)\| ds. \tag{4.14}$$

Furthermore, we can see that

$$\begin{aligned}
&\|v(\cdot, t_n) - P_h v(\cdot, t_n)\| \\
&\leq Ch^\chi \|v(\cdot, t_n)\|_{\dot{H}^\chi(\mathcal{O})} \\
&\leq Ch^\chi \|\Delta^{\frac{1}{2}\chi} v(\cdot, t_n)\| \\
&= Ch^\chi \left\| \int_0^{t_n} \frac{1}{2\pi i} \int_{\Gamma_{\theta,\kappa}} e^{z(t_n-s)} z^{\alpha-1} \Delta^{\frac{1}{2}\chi} (z^\alpha - \Delta)^{-1} f(\cdot, s) dz ds \right\| \\
&\leq Ch^\chi \int_0^{t_n} \int_{\Gamma_{\theta,\kappa}} |e^{z(t_n-s)}| |z|^{\alpha-1} \|\Delta^{\frac{1}{2}\chi} (z^\alpha - \Delta)^{-1}\| \|f(\cdot, s)\| |dz| ds \\
&\leq Ch^\chi \int_0^{t_n} \int_{\Gamma_{\theta,\kappa}} |e^{z(t_n-s)}| |z|^{\frac{1}{2}\chi\alpha-1} \|f(\cdot, s)\| |dz| ds \quad (\text{here we use (3.16)}) \\
&\leq Ch^\chi \int_0^{t_n} (t_n - s)^{-\frac{1}{2}\chi\alpha} \|f(\cdot, s)\| ds.
\end{aligned} \tag{4.15}$$

The estimates (4.14) and (4.15) imply

$$\begin{aligned}
\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| &\leq Ch^\chi \int_0^{t_n} (t_n - s)^{-\frac{1}{2}\chi\alpha} \|f(\cdot, s)\| ds \\
&\leq Ch^\chi \|f\|_{L^{\frac{2}{2-\chi\alpha},1}(0,T;L^2(\mathcal{O}))}, \quad (\text{here we use (2.1)})
\end{aligned}$$

completing the proof in Case 3.

The combination of Cases 1, 2 and 3 yields

$$\begin{aligned}
\|v(\cdot, t_n) - v^{(h)}(\cdot, t_n)\| &\leq C(\|\psi_0\|_{\dot{H}^\chi(\mathcal{O})} + \|f\|_{L^{\frac{2}{2-\chi\alpha},1}(0,T;L^2(\mathcal{O}))}) h^\chi \\
&\leq C(\|\psi_0\|_{\dot{H}^\chi(\mathcal{O})} + \|f\|_{L^{\frac{4}{2+\alpha d},1}(0,T;L^2(\mathcal{O}))}) h^\chi,
\end{aligned} \tag{4.16}$$

where we have used the fact  $\frac{2}{2-\chi\alpha} \leq \frac{4}{2+\alpha d}$  in the last inequality.

## 4.2 Temporal discretization: estimate of $\|v^{(h)}(\cdot, t_n) - v_n^{(h)}\|$

To estimate  $\|v^{(h)}(\cdot, t_n) - v_n^{(h)}\|$ , we consider the following three cases:

Case 1':  $\psi_0 \neq 0$ ,  $f = 0$ ,  $\alpha \in (0, 1] \cap (0, \frac{2}{d})$ .

Case 2':  $\psi_0 \neq 0$ ,  $f = 0$ ,  $\alpha \in (1, \frac{2}{d})$ . (This requires  $d = 1$ , cf. Theorem 1)

Case 3':  $\psi_0 = 0$ ,  $f \neq 0$ .

In Case 1', it is straightforward to verify that the solutions of (4.2) and (2.22) are given by  $v^{(h)}(\cdot, t_n) = v_n^{(h)} = P_h \psi_0$ ,  $n = 1, 2, \dots$ . Consequently, we have

$$v^{(h)}(\cdot, t_n) - v_n^{(h)} = 0. \tag{4.17}$$

In Case 2', by using the Laplace transform, we can derive the following error representation (see Appendix B):

$$\begin{aligned}
v^{(h)}(\cdot, t_n) - v_n^{(h)} &= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt_n} z^{-1} (z^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz \\
&\quad - \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt_n} e^{-z\tau} \delta_\tau (e^{-z\tau})^{-1} (\delta_\tau (e^{-z\tau})^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz \\
&= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt_n} D_h^{(1)}(z) \Delta_h^\gamma P_h \psi_0 dz \\
&\quad + \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa} \setminus \Gamma_{\theta, \kappa}^{(\tau)}} e^{zt_n} z^{-1} \Delta_h^{1-\gamma} (z^\alpha - \Delta_h)^{-1} \Delta_h^\gamma P_h \psi_0 dz \\
&=: \mathcal{I}_n^{(h)} + \mathcal{J}_n^{(h)}, \tag{4.18}
\end{aligned}$$

where  $\gamma \in [0, 1]$  and

$$D_h^{(1)}(z) = z^{-1} \Delta_h^{1-\gamma} (z^\alpha - \Delta_h)^{-1} - e^{-z\tau} \delta_\tau (e^{-z\tau})^{-1} \Delta_h^{1-\gamma} (\delta_\tau (e^{-z\tau})^\alpha - \Delta_h)^{-1}.$$

Lemma 1 and Lemma 5 imply

$$\begin{aligned}
\|D_h^{(1)}(z)\| &\leq C|z|^{-1} (\|\Delta_h^{1-\gamma} (z^\alpha - \Delta_h)^{-1}\| + \|\Delta_h^{1-\gamma} (\delta_\tau (e^{-z\tau})^\alpha - \Delta_h)^{-1}\|) \\
&\leq C|z|^{-\gamma\alpha-1} \quad (\text{here we use (3.16)})
\end{aligned}$$

and

$$\begin{aligned}
\|D_h^{(1)}(z)\| &\leq C|z|^{-1} \|\Delta_h^{1-\gamma} [(z^\alpha - \Delta_h)^{-1} - (\delta_\tau (e^{-z\tau})^\alpha - \Delta_h)^{-1}]\| \\
&\quad + C|z|^{-1} \|e^{-z\tau} \delta_\tau (e^{-z\tau})^{-1}\| \|\Delta_h^{1-\gamma} (\delta_\tau (e^{-z\tau})^\alpha - \Delta_h)^{-1}\| \\
&\leq C|z|^{-1} \|z^\alpha - \delta_\tau (e^{-z\tau})^\alpha\| \|\Delta_h^{1-\gamma} (\delta_\tau (e^{-z\tau})^\alpha - \Delta_h)^{-1}\| \|(z^\alpha - \Delta_h)^{-1}\| \\
&\quad + C|(\delta_\tau (e^{-z\tau}) - z) + (1 - e^{-z\tau})z| |z|^{-1} |\delta_\tau (e^{-z\tau})|^{-1} |z|^{-\gamma\alpha} \quad (\text{use (3.16)}) \\
&\leq C|z|^{-\gamma\alpha\tau}.
\end{aligned}$$

The last two inequalities further imply

$$\|D_h^{(1)}(z)\| \leq C|z|^{-\gamma\alpha-1+\eta\tau\eta} \quad \forall \eta \in [0, 1].$$

Then, we have

$$\begin{aligned}
\|\mathcal{I}_n^{(h)}\| &\leq C \int_{\Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt_n}| \|D_h^{(1)}(z)\| \|\Delta_h^\gamma P_h \psi_0\| |dz| \\
&\leq C \int_{\Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt_n}| |z|^{-\gamma\alpha-1+\eta\tau\eta} \|\Delta_h^\gamma P_h \psi_0\| |dz| \\
&\leq C\tau^\eta \|\Delta_h^\gamma P_h \psi_0\| \left( \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} e^{rt_n \cos(\theta)} r^{-\gamma\alpha-1+\eta} dr + \int_{-\theta}^{\theta} e^{\kappa t_n \cos(\varphi)} \kappa^{-\gamma\alpha+\eta} d\varphi \right) \\
&\leq C\tau^\eta \|\Delta_h^\gamma P_h \psi_0\| \left( t_n^{\gamma\alpha-\eta} \int_{\kappa t_n}^{\frac{t_n \pi}{\tau \sin(\theta)}} e^{-s|\cos(\theta)|} s^{-\gamma\alpha-1+\eta} ds + \kappa^{-\gamma\alpha+\eta} \right) \\
&\leq C\tau^{\gamma\alpha} (1 + \kappa^{-\gamma\alpha+\eta}) \|\Delta_h^\gamma P_h \psi_0\| \quad (\text{this requires } \eta > \gamma\alpha) \\
&\leq C\tau^{\gamma\alpha} \|\psi_0\|_{\dot{H}^{2\gamma}(\mathcal{O})}, \tag{4.19}
\end{aligned}$$

where we have used (2.14) in the last inequality. Similarly, by using Lemma 5 we have

$$\|\mathcal{J}_n^{(h)}\| \leq C \int_{\Gamma_{\theta, \kappa} \setminus \Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt_n}| |z|^{-1} \|\Delta_h^{1-\gamma} (z^\alpha - \Delta_h)^{-1}\| \|\Delta_h^\gamma P_h \psi_0\| |dz|$$

$$\begin{aligned}
&\leq C \|\Delta_h^\gamma P_h \psi_0\| \int_{\frac{\pi}{\tau \sin(\theta)}}^{\infty} e^{-rt_n |\cos(\theta)|} r^{-1-\gamma\alpha} dr \quad (\text{here we use (3.16)}) \\
&\leq C \tau^{1+\gamma\alpha} \|\Delta_h^\gamma P_h \psi_0\| \int_{\frac{\pi}{\tau \sin(\theta)}}^{\infty} e^{-rt_n |\cos(\theta)|} dr \\
&\leq C \tau^{\gamma\alpha} \|\Delta_h^\gamma P_h \psi_0\| \\
&\leq C \tau^{\gamma\alpha} \|\psi_0\|_{\dot{H}^{2\gamma}(\mathcal{O})}, \tag{4.20}
\end{aligned}$$

where we have used (2.14) again in the last inequality. By choosing  $\gamma = \frac{1}{2\alpha} - \frac{d}{4} \in (0, 1)$  and  $\eta > \frac{1}{2} - \frac{\alpha d}{4} \in (0, 1)$  (recall that  $1 < \alpha < \frac{d}{2}$ ), substituting (4.19) and (4.20) into (4.18) yields

$$\|v^{(h)}(\cdot, t_n) - v_n^{(h)}\| \leq C \tau^{\frac{1}{2} - \frac{\alpha d}{4}} \|\psi_0\|_{\dot{H}^{\frac{1}{\alpha} - \frac{d}{2}}(\mathcal{O})} \leq C \tau^{\frac{1}{2} - \frac{\alpha d}{4}} \|\psi_0\|_{\dot{H}^\infty(\mathcal{O})}, \tag{4.21}$$

where we have noted that  $\frac{1}{\alpha} - \frac{d}{2} \leq \chi$  for  $\alpha > 1$ .

In Case 3', with  $\alpha \in (0, 2)$ , we have

$$v^{(h)}(\cdot, t_n) = \int_0^{t_n} E^{(h)}(t_n - s) P_h f(\cdot, s) ds, \tag{4.22}$$

$$v_n^{(h)} = \int_0^{t_n} E_\tau^{(h)}(t_n - s) P_h f_\tau(\cdot, s) ds, \tag{4.23}$$

where  $E_\tau^{(h)}(t)$  is given by (3.11) and  $f_\tau$  is the piecewise constant (in time) function defined in (2.2). The difference of the two expressions (4.22) and (4.23) yields

$$\begin{aligned}
v^{(h)}(\cdot, t_n) - v_n^{(h)} &= \int_0^{t_n} [E^{(h)}(t_n - s) - E_\tau^{(h)}(t_n - s)] P_h f_\tau(\cdot, s) ds \\
&\quad + \int_0^{t_n} E^{(h)}(t_n - s) [P_h f(\cdot, s) - P_h f_\tau(\cdot, s)] ds \\
&=: \mathcal{K}_n^{(h)} + \mathcal{L}_n^{(h)}. \tag{4.24}
\end{aligned}$$

In the following, we estimate  $\mathcal{K}_n^{(h)}$  and  $\mathcal{L}_n^{(h)}$  separately.

By the inverse Laplace transform rule  $\mathcal{L}^{-1}(\widehat{f}\widehat{g})(t) = \int_0^t \mathcal{L}^{-1}(\widehat{f})(t-s) \mathcal{L}^{-1}(\widehat{g})(s) ds$ , we have

$$\begin{aligned}
\mathcal{K}_n^{(h)} &= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt_n} z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} P_h \widehat{f}_\tau(\cdot, z) dz \\
&\quad - \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt_n} \frac{z\tau}{e^{z\tau} - 1} \delta_\tau(e^{-z\tau})^{\alpha-1} (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1} P_h \widehat{f}_\tau(\cdot, z) dz \\
&= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa} \setminus \Gamma_{\theta, \kappa}^{(\tau)}} e^{zt_n} \widehat{D}_h^{(2)}(z) P_h \widehat{f}_\tau(\cdot, z) dz - \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt_n} \widehat{D}_h^{(3)}(z) P_h \widehat{f}_\tau(\cdot, z) dz \\
&= \int_0^{t_n} D_h^{(2)}(t_n - s) P_h f_\tau(\cdot, s) ds + \int_0^{t_n} D_h^{(3)}(t_n - s) P_h f_\tau(\cdot, s) ds, \tag{4.25}
\end{aligned}$$

where

$$\begin{aligned}
\widehat{D}_h^{(2)}(z) &= z^{\alpha-1} (z^\alpha - \Delta_h)^{-1}, \\
\widehat{D}_h^{(3)}(z) &= z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} - \frac{z\tau}{e^{z\tau} - 1} \delta_\tau(e^{-z\tau})^{\alpha-1} (\delta_\tau(e^{-z\tau})^\alpha - \Delta_h)^{-1},
\end{aligned}$$

and

$$D_h^{(2)}(t)\phi = \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa} \setminus \Gamma_{\theta, \kappa}^{(\tau)}} e^{zt} \widehat{D}_h^{(2)}(z) \phi dz, \quad D_h^{(3)}(t)\phi = \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{zt} \widehat{D}_h^{(3)}(z) \phi dz.$$

Using the similar method as introduced in [12, Lemma 3.4] (as well as the inequalities (3.15) in Lemma 5 and (C.1) in Appendix C), it is easy to see that

$$\|\widehat{D}_h^{(2)}(z)\| + \|\widehat{D}_h^{(3)}(z)\| \leq C|z|^{-1} \quad \text{and} \quad \|\widehat{D}_h^{(3)}(z)\| \leq C\tau. \quad (4.26)$$

The last two inequalities further imply

$$\|\widehat{D}_h^{(3)}(z)\| \leq C|z|^{-\theta}\tau^{1-\theta} \quad \forall \theta \in [0, 1]. \quad (4.27)$$

Consequently,

$$\begin{aligned} \|D_h^{(2)}(t)\| &\leq C \int_{\Gamma_{\theta, \kappa} \setminus \Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt}| \|\widehat{D}_h^{(2)}(z)\| |dz| \\ &\leq C \int_{\frac{\pi}{\tau \sin(\theta)}}^{\infty} e^{rt \cos(\theta)} r^{-1} dr \\ &\leq C\tau \int_{\frac{\pi}{\tau \sin(\theta)}}^{\infty} e^{rt \cos(\theta)} dr \\ &\leq Ct^{-1}\tau \int_{\frac{t\pi}{\tau \sin(\theta)}}^{\infty} e^{s \cos(\theta)} ds \\ &\leq Ct^{-1}\tau \\ &\leq Ct^{-1/q}\tau^{1/q} \quad \forall q > 1, \quad t \geq \tau, \end{aligned}$$

and

$$\begin{aligned} \|D_h^{(2)}(t)\| &\leq C \int_{\Gamma_{\theta, \kappa} \setminus \Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt}| \|\widehat{D}_h^{(2)}(z)\| |dz| \\ &\leq C \int_{\frac{\pi}{\tau \sin(\theta)}}^{\infty} e^{rt \cos(\theta)} r^{-1} dr \\ &\leq C \int_{\frac{t\pi}{\tau \sin(\theta)}}^{\infty} e^{s \cos(\theta)} s^{-1} ds \\ &\leq C \int_1^{\infty} e^{s \cos(\theta)} s^{-1} ds + C \int_{\frac{t\pi}{\tau \sin(\theta)}}^1 e^{s \cos(\theta)} s^{-1} ds \\ &\leq C + C \ln(t^{-1}\tau) \\ &\leq Ct^{-1/q}\tau^{1/q} \quad \forall q > 1, \quad t \in (0, \tau). \end{aligned}$$

A combination of the last two estimates gives the following estimate of  $\|D_h^{(2)}(t)\|$ :

$$\|D_h^{(2)}(t)\| \leq Ct^{-1/q}\tau^{1/q} \quad \forall q > 1, \quad t \geq 0. \quad (4.28)$$

Similarly, we have

$$\begin{aligned} \|D_h^{(3)}(t)\| &\leq C \int_{\Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt}| \|\widehat{D}_h^{(3)}(z)\| |dz| \\ &\leq C\tau^{1/q} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} |e^{zt}| |z|^{-(1-1/q)} |dz| \quad (\text{set } \theta = 1 - 1/q \text{ in (4.27)}) \\ &\leq C\tau^{1/q} \int_{\kappa}^{\frac{\pi}{\tau \sin(\theta)}} e^{rt \cos(\theta)} r^{-(1-1/q)} dr + C\tau^{1/q} \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} \kappa^{1/q} d\varphi \\ &\leq Ct^{-1/q}\tau^{1/q} \int_{\kappa t}^{\frac{t\pi}{\tau \sin(\theta)}} e^{s \cos(\theta)} ds + C\kappa^{1/q}\tau^{1/q} e^{\kappa T} \\ &\leq Ct^{-1/q}\tau^{1/q} + CT^{-1/q}\tau^{1/q} \\ &\leq Ct^{-1/q}\tau^{1/q} \quad \forall q > 1, \quad t \in (0, T]. \end{aligned} \quad (4.29)$$

Substituting (4.28) and (4.29) into (4.25) yields

$$\begin{aligned}
\|\mathcal{K}_n^{(h)}\| &= \left\| \int_0^{t_n} D^{(2)}(t_n - s) P_h f_\tau(\cdot, s) ds + \int_0^{t_n} D^{(3)}(t_n - s) P_h f_\tau(\cdot, s) ds \right\| \\
&\leq C\tau^{1/q} \int_0^{t_n} (t_n - s)^{-1/q} \|P_h f_\tau(\cdot, s)\| ds \\
&\leq C\tau^{1/q} \|f_\tau\|_{L^{q',1}(0,t_n;L^2(\mathcal{O}))} \\
&\leq C\tau^{1/q} \|f\|_{L^{q',1}(0,t_n;L^2(\mathcal{O}))},
\end{aligned} \tag{4.30}$$

where the last inequality follows from (2.3). This completes the estimate of the first term in (4.24).

The second term in (4.24) can be estimated as follows:

$$\begin{aligned}
\|\mathcal{L}_n^{(h)}\| &= \left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} E^{(h)}(t_n - s) [P_h f(\cdot, s) - P_h f_\tau(\cdot, s)] ds \right\| \\
&= \left\| \sum_{j=1}^n \int_{t_{j-1}}^{t_j} (E^{(h)}(t_n - s) - E^{(h)}(t_n - t_{j-1})) [P_h f(\cdot, s) - P_h f_\tau(\cdot, s)] ds \right\| \\
&\leq \sum_{j=1}^n \int_{t_{j-1}}^{t_j} \|E^{(h)}(t_n - s) - E^{(h)}(t_n - t_{j-1})\| \|P_h f(\cdot, s) - P_h f_\tau(\cdot, s)\| ds \\
&\leq \sum_{j=1}^n \int_{t_{j-1}}^{t_j} C\tau^{1/q} (t_n - s)^{-1/q} \|P_h f(\cdot, s) - P_h f_\tau(\cdot, s)\| ds \\
&\leq C\tau^{1/q} \|f - f_\tau\|_{L^{q',1}(0,t_n;L^2(\mathcal{O}))} \\
&\leq C\tau^{1/q} \|f\|_{L^{q',1}(0,t_n;L^2(\mathcal{O}))},
\end{aligned} \tag{4.31}$$

where we have used the following identity to obtain the second equality in (4.31):

$$\begin{aligned}
&\int_{t_{j-1}}^{t_j} E^{(h)}(t_n - t_{j-1}) P_h f_\tau(\cdot, s) ds \\
&= \frac{1}{2\pi i} \int_{t_{j-1}}^{t_j} \int_{\Gamma_{\theta,\kappa}} e^{z(t_n - t_j)} z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} P_h f_\tau(\cdot, s) dz ds \quad (\text{use (A.5) here}) \\
&= \frac{1}{2\pi i} \int_{t_{j-1}}^{t_j} \int_{\Gamma_{\theta,\kappa}} e^{z(t_n - t_j)} z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} \frac{1}{\tau} \int_{t_{j-1}}^{t_j} P_h f(\cdot, \xi) d\xi ds \quad (\text{use (2.2) here}) \\
&= \frac{1}{2\pi i} \int_{t_{j-1}}^{t_j} \int_{\Gamma_{\theta,\kappa}} e^{z(t_n - t_j)} z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} P_h f(\cdot, \xi) d\xi \\
&= \int_{t_{j-1}}^{t_j} E^{(h)}(t_n - t_{j-1}) P_h f(\cdot, s) ds,
\end{aligned}$$

and we have used the following estimate to obtain the second inequality in (4.31): for  $q > 1$ ,

$$\begin{aligned}
&\|E^{(h)}(t_n - s) - E^{(h)}(t_n - t_{j-1})\| \\
&\leq C \int_{\Gamma_{\theta,\kappa}} |e^{z(t_n - s)} - e^{z(t_n - t_{j-1})}| |z|^{\alpha-1} \|(z^\alpha - \Delta_h)^{-1}\| |dz| \\
&\leq C \int_{\Gamma_{\theta,\kappa}} |e^{z(t_n - s)}| |1 - e^{z(s - t_{j-1})}| |z|^{-1} |dz| \quad (\text{here we use (3.15)}) \\
&\leq C \int_{\Gamma_{\theta,\kappa}} |e^{z(t_n - s)}| \tau^{1/q} |z|^{-(1-1/q)} |dz| \quad (\text{here we use (C.2)})
\end{aligned}$$

$$\leq C\tau^{1/q}(t_n - s)^{-1/q}. \quad (4.32)$$

By choosing  $q > 1$  to satisfy  $\frac{1}{q} = \frac{1}{2} - \frac{\alpha d}{4}$ , we have  $q' = \frac{4}{2+\alpha d}$ . Then substituting (4.30) and (4.31) into (4.24) yields

$$\|v_n^{(h)} - v^{(h)}(\cdot, t_n)\| \leq C\tau^{\frac{1}{2} - \frac{\alpha d}{4}} \|f\|_{L^{\frac{4}{2+\alpha d}, 1}(0, t_n; L^2(\mathcal{O}))}, \quad (4.33)$$

completing the proof of Case 3.

By combining (4.17), (4.21) and (4.33) (the results of Cases 1, 2 and 3), we obtain

$$\|v_n^{(h)} - v^{(h)}(\cdot, t_n)\| \leq C(\|\psi_0\|_{\dot{H}^\alpha(\mathcal{O})} + \|f\|_{L^{\frac{4}{2+\alpha d}, 1}(0, t_n; L^2(\mathcal{O}))})\tau^{\frac{1}{2} - \frac{\alpha d}{4}}. \quad (4.34)$$

The estimates (4.16) and (4.34) imply (4.1), completing the proof of Theorem 1.  $\square$

## 5 Numerical examples

In this section, we present numerical examples to illustrate the theoretical analyses.

We consider the one-dimensional stochastic partial integro-differential equation (1.1) for  $0 \leq x \leq 1$ ,  $0 < t \leq 1$ , with homogeneous Dirichlet boundary condition and initial condition  $\psi_0(x) = x(1-x)$ . Here, we let  $\sigma = 1$  in (1.1) and

$$f(x, t) = \begin{cases} 1 & \text{for } 0 \leq x \leq \frac{1}{2}, \\ -1 & \text{for } \frac{1}{2} < x \leq 1. \end{cases}$$

The problem (1.1) is discretized by using the scheme (2.16)-(2.17).

To investigate the convergence rate in space, we first solve the problem (1.1) by taking the mesh size  $h_k = 1/M_k = 2^{-k}$ ,  $k = 2, 3, 4, 5$ , and using a sufficiently small time step  $\tau = 2^{-14}$  so that the temporal discretization error is relatively negligible. Then, the error

$$E(h_k) = \frac{1}{I} \sum_{i=1}^I \|\psi_N^{(h_k)}(\omega_i) - \psi_N^{(h_{k-1})}(\omega_i)\|$$

is computed for  $k = 3, 4, 5$ , by using  $I = 10000$  independent realizations for each spatial mesh size. By Theorem 1, the error  $E(h_k)$  is expected to have the convergence rate  $O(h^{\frac{1}{\alpha} - \frac{1}{2}})$  for  $\alpha \in [\frac{1}{2}, 2)$ , and  $O(h^{\frac{3}{2}})$  for  $\alpha \in (0, \frac{1}{2})$  in one-dimensional spatial domain. The numerical results are presented in Table 1 and consistent with the theoretical analyses that the spatial order of convergence increases as  $\alpha$  decreases and stays at the order  $\frac{3}{2}$  when  $\alpha$  further decreases.

Secondly, we solve the problem (1.1) by using the time step  $\tau_k = 2^{-k}$ ,  $k = 6, 7, 8, 9$ . In order to focus on the temporal discretization error, a sufficiently small spatial mesh size  $h = 1/M = 2^{-10}$  is used such that the spatial discretization error can be relatively negligible. Similarly, we consider  $I = 10000$  independent realizations for each time step and compute the error  $E(\tau_k)$  by

$$E(\tau_k) = \frac{1}{I} \sum_{i=1}^I \|\psi_{N, \tau_k}^{(h)}(\omega_i) - \psi_{N, \tau_{k-1}}^{(h)}(\omega_i)\|$$

for  $k = 7, 8, 9$ , which is expected to have the convergence rate  $O(\tau^{\frac{1}{2} - \frac{\alpha}{4}})$  by Theorem 1. Clearly, the results in Table 2 illustrate the sharp convergence rate.

**Table 1**  $E(h_k)$  and convergence rates in space

$\alpha \backslash h_k$	$2^{-3}$	$2^{-4}$	$2^{-5}$	order
$\alpha = 0.25$	1.1669e-02	3.9124e-03	1.3519e-03	1.555 (1.500)
$\alpha = 0.75$	2.4353e-02	1.2987e-02	6.6322e-03	0.938 (0.833)
$\alpha = 1.25$	8.3694e-02	6.7186e-02	5.4196e-02	0.314 (0.300)

**Table 2**  $E(\tau_k)$  and convergence rates in time

$\alpha \backslash \tau_k$	$2^{-7}$	$2^{-8}$	$2^{-9}$	order
$\alpha = 0.25$	2.2103e-03	1.7275e-03	1.3454e-03	0.359 (0.4375)
$\alpha = 0.75$	1.5613e-02	1.2621e-02	1.0177e-02	0.309 (0.3125)
$\alpha = 1.25$	5.0056e-02	4.4012e-02	3.8869e-02	0.183 (0.1875)

### A Mild solution of (1.1)

In the case  $\alpha \in (1, 2)$ , the boundary condition  $\partial_t^{1-\alpha} \psi = 0$  is equivalent to  $\psi = 0$  on  $\partial\Omega$  (this can be checked by taking Laplace transform in time). Similarly, in the case  $\alpha \in (0, 1]$ , the boundary condition  $\partial_t^{1-\alpha} \psi = 0$  is equivalent to  $\psi - \psi_0 = 0$  on  $\partial\Omega \times [0, \infty)$ , where  $\psi_0 = \psi(\cdot, 0)$  is the initial value in (1.1).

In the case  $\sigma = 0$ , the solution of the corresponding deterministic problem of (1.1) can be expressed by (via Laplace transform, cf. [22, (3.11) and line 4 of page 12] in the case  $\alpha \in (1, 2)$ )

$$\psi(\cdot, t) = \begin{cases} \psi_0 + \int_0^t E(t-s)f(\cdot, s)ds & \text{if } \alpha \in (0, 1], \\ E(t)\psi_0 + \int_0^t E(t-s)f(\cdot, s)ds & \text{if } \alpha \in (1, 2), \end{cases} \quad (\text{A.1})$$

where the operator  $E(t) : L^2(\mathcal{O}) \rightarrow L^2(\mathcal{O})$  is given by

$$E(t)\phi := \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{\alpha-1} (z^\alpha - \Delta)^{-1} \phi dz \quad \forall \phi \in L^2(\mathcal{O}) \quad (\text{A.2})$$

with integration over a contour  $\Gamma_{\theta, \kappa}$  on the complex plane.

Correspondingly, the mild solution of the stochastic problem (1.1) is defined as (cf. [17, Proposition 2.7] and [23])

$$\psi(\cdot, t) = \begin{cases} \psi_0 + \int_0^t E(t-s)f(\cdot, s)ds + \sigma \int_0^t E(t-s)dW(\cdot, s) & \text{if } \alpha \in (0, 1], \\ E(t)\psi_0 + \int_0^t E(t-s)f(\cdot, s)ds + \sigma \int_0^t E(t-s)dW(\cdot, s) & \text{if } \alpha \in (1, 2). \end{cases} \quad (\text{A.3})$$

For any given initial data  $\psi_0 \in L^2(\mathcal{O})$  and source  $f \in L^1(0, T; L^2(\mathcal{O}))$ , the expression (A.3) defines a mild solution  $\psi \in C([0, T]; L^2(\Omega; L^2(\mathcal{O})))$ . In the case  $\psi_0 = f = 0$  and  $\sigma \neq 0$ , a simple proof of this result can be found in [12, Appendix]; in the case  $\sigma = 0$  ( $\psi_0$  and  $f$  may not be zero), the result is a consequence of the boundedness of the operator  $E(t) : L^2(\mathcal{O}) \rightarrow L^2(\mathcal{O})$ , i.e.,

$$\begin{aligned} \|E(t)v\| &\leq C \int_{\Gamma_{\theta, \kappa}} |e^{zt}| |z|^{\alpha-1} \|(z^\alpha - \Delta)^{-1}v\| |dz| \\ &\leq C \|v\| \int_{\kappa}^{\infty} e^{-rt} |\cos(\theta)| r^{-1} dr + C \|v\| \int_{-\theta}^{\theta} e^{\kappa t \cos(\varphi)} d\varphi \\ &\leq C \|v\| \quad \forall v \in L^2(\mathcal{O}). \end{aligned} \quad (\text{A.4})$$

Similarly, the discrete operator  $E^{(h)}(t) : X_h \rightarrow X_h$  defined by

$$E^{(h)}(t)\phi := \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} \phi dz \quad \forall \phi \in X_h, \quad (\text{A.5})$$

is also bounded on the finite element subspace  $X_h$ , i.e.,

$$\|E^{(h)}(t)v\| \leq C \|v\| \quad \forall v \in X_h, \quad (\text{A.6})$$

where the constant  $C$  is independent of the mesh size  $h$ .

## B Representation of the discrete solutions

For  $f = 0$  we prove the following representation of the solutions of (4.2) and (2.23):

$$v^{(h)}(\cdot, t_n) = P_h \psi_0 + \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt_n} z^{-1} (z^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz, \quad (\text{B.1})$$

$$v_n^{(h)} = P_h \psi_0 + \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}^{(\tau)}} e^{tnz} e^{-z\tau} \delta(e^{-z\tau})^{-1} (\delta(e^{-z\tau})^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz, \quad (\text{B.2})$$

which are used in (4.18) in estimating the error of temporal discretization.

In fact, (B.1) is a consequence of (A.3): replacing  $E(t)$  by  $E^{(h)}(t)$  and substituting  $\phi = P_h \psi_0$  yield

$$\begin{aligned} v^{(h)}(\cdot, t_n) &= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{\alpha-1} (z^\alpha - \Delta_h)^{-1} P_h \psi_0 dz \\ &= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{-1} (z^\alpha - \Delta_h + \Delta_h) (z^\alpha - \Delta_h)^{-1} P_h \psi_0 dz \\ &= \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{-1} P_h \psi_0 dz + \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{-1} (z^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz \\ &= P_h \psi_0 + \frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{-1} (z^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz, \end{aligned}$$

where we have used the identity  $\frac{1}{2\pi i} \int_{\Gamma_{\theta, \kappa}} e^{zt} z^{-1} dz = 1$  (i.e., the inverse Laplace transform of  $z^{-1}$  is 1).

It remains to prove (B.2). To this end, we rewrite (2.23) as

$$\bar{\partial}_\tau(v_n^{(h)} - P_h \psi_0) - \Delta_h \bar{\partial}_\tau^{1-\alpha}(v_n^{(h)} - P_h \psi_0) = \Delta_h \bar{\partial}_\tau^{1-\alpha}(P_h \psi_0)_n, \quad (\text{B.3})$$

where  $\bar{\partial}_\tau^{1-\alpha}(P_h \psi_0)_n := \frac{1}{\tau^{1-\alpha}} \sum_{j=1}^n b_{n-j} P_h \psi_0$ . Since we are only interested in the solutions  $v_n^{(h)}$ ,  $n = 1, \dots, N$ , we define

$$\tilde{v}_n^{(h)} = \begin{cases} v_n^{(h)} & 1 \leq n \leq N, \\ P_h \psi_0 & n \geq N+1, \end{cases}$$

which satisfies the equation

$$\bar{\partial}_\tau(\tilde{v}_n^{(h)} - P_h \psi_0) - \Delta_h \bar{\partial}_\tau^{1-\alpha}(\tilde{v}_n^{(h)} - P_h \psi_0) = \Delta_h \bar{\partial}_\tau^{1-\alpha}(P_h \psi_0)_n + g_n, \quad (\text{B.4})$$

with  $g_n = 0$  for  $1 \leq n \leq N$ . The right-hand side of (B.4) differs from (B.3) only for  $n \geq N+1$ , that

$$\begin{aligned} \|g_n\| &\leq \|\Delta_h \bar{\partial}_\tau^{1-\alpha}(\tilde{v}_n^{(h)} - P_h \psi_0)\| + \|\Delta_h \bar{\partial}_\tau^{1-\alpha}(P_h \psi_0)_n\| \\ &\leq \frac{1}{\tau^{1-\alpha}} \sum_{j=1}^N |b_{n-j}| \|\tilde{v}_j^{(h)} - P_h \psi_0\| + \frac{1}{\tau^{1-\alpha}} \sum_{j=1}^n |b_{n-j}| \|P_h \psi_0\| \\ &\leq C\tau^{\alpha-1} \left( \sum_{j=1}^N |b_{n-j}| + \sum_{j=1}^n |b_{n-j}| \right) \\ &\leq C\tau^{\alpha-1} n^{\alpha-1}, \end{aligned}$$

as  $n \rightarrow \infty$ . Thus  $\sum_{n=N+1}^\infty g_n \zeta^n$  is an analytic function of  $\zeta$  for  $|\zeta| < 1$ .

By (2.8), summing up (B.4) times  $\zeta^n$  for  $n = 1, 2, \dots$ , yields

$$\left( \frac{1-\zeta}{\tau} - \left( \frac{1-\zeta}{\tau} \right)^{1-\alpha} \Delta_h \right) \sum_{n=1}^\infty (\tilde{v}_n^{(h)} - P_h \psi_0) \zeta^n = \Delta_h \left( \frac{1-\zeta}{\tau} \right)^{1-\alpha} \frac{\zeta}{1-\zeta} P_h \psi_0 + \sum_{n=N+1}^\infty g_n \zeta^n,$$

which implies

$$\begin{aligned} \sum_{n=1}^\infty (\tilde{v}_n^{(h)} - P_h \psi_0) \zeta^n &= \left( \frac{1-\zeta}{\tau} \right)^{-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \Delta_h P_h \psi_0 \frac{\zeta}{\tau} \\ &\quad + \left( \frac{1-\zeta}{\tau} \right)^{\alpha-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \sum_{n=N+1}^\infty g_n \zeta^n. \end{aligned}$$



For  $\kappa > 0$  and  $\varrho_\kappa = e^{-(\kappa+1)\tau} \in (0, 1)$ , the Cauchy integral formula implies that

$$\begin{aligned}
& \tilde{v}_n^{(h)} - P_h \psi_0 \\
&= \frac{1}{2\pi i} \int_{|\zeta|=\varrho_\kappa} \zeta^{-n-1} \sum_{n=1}^{\infty} (v_n^{(h)} - P_h \psi_0) \zeta^n d\zeta \\
&= \frac{1}{2\pi i} \int_{|\zeta|=\varrho_\kappa} \zeta^{-n} \left( \frac{1-\zeta}{\tau} \right)^{-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \frac{1}{\tau} \Delta_h P_h \psi_0 d\zeta \\
&\quad + \frac{1}{2\pi i} \int_{|\zeta|=\varrho_\kappa} \left( \frac{1-\zeta}{\tau} \right)^{\alpha-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \sum_{m=N+1}^{\infty} g_m \zeta^{m-n-1} d\zeta.
\end{aligned} \tag{B.5}$$

For  $1 \leq n \leq N$  the function  $\left( \frac{1-\zeta}{\tau} \right)^{\alpha-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \sum_{m=N+1}^{\infty} g_m \zeta^{m-n-1}$  is analytic in  $|\zeta| < 1$ . Consequently, Cauchy's integral theorem implies

$$\frac{1}{2\pi i} \int_{|\zeta|=\varrho_\kappa} \left( \frac{1-\zeta}{\tau} \right)^{\alpha-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \sum_{m=N+1}^{\infty} g_m \zeta^{m-n-1} d\zeta = 0.$$

Substituting this identity into (B.5) yields, for  $1 \leq n \leq N$ ,

$$\begin{aligned}
& \tilde{v}_n^{(h)} - P_h \psi_0 \\
&= v_n^{(h)} - P_h \psi_0 \\
&= \frac{1}{2\pi i} \int_{|\zeta|=\varrho_\kappa} \zeta^{-n} \left( \frac{1-\zeta}{\tau} \right)^{-1} \left( \left( \frac{1-\zeta}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \frac{1}{\tau} \Delta_h P_h \psi_0 d\zeta \\
&= \frac{1}{2\pi i} \int_{\Gamma^\tau} e^{tnz} e^{-z\tau} \left( \frac{1-e^{-\tau z}}{\tau} \right)^{-1} \left( \left( \frac{1-e^{-\tau z}}{\tau} \right)^\alpha - \Delta_h \right)^{-1} \Delta_h P_h \psi_0 dz \\
&= \frac{1}{2\pi i} \int_{\Gamma^\tau} e^{tnz} e^{-z\tau} \delta(e^{-z\tau})^{-1} (\delta(e^{-z\tau})^\alpha - \Delta_h)^{-1} \Delta_h P_h \psi_0 dz,
\end{aligned} \tag{B.6}$$

where we have used the change of variable  $\zeta = e^{-z\tau}$ , which converts the path of integration to the contour

$$\Gamma^\tau = \{z = \kappa + 1 + iy : y \in \mathbb{R} \text{ and } |y| \leq \pi/\tau\}.$$

The angle condition (3.4) and [3, Theorem 3.7.11] imply that the integrand on the right-hand side of (B.6) is analytic in the region

$$\Sigma_{\theta, \kappa}^\tau = \left\{ z \in \mathbb{C} : |\arg(z)| \leq \theta, |z| \geq \kappa, |\operatorname{Im}(z)| \leq \frac{\pi}{\tau}, \operatorname{Re}(z) \leq \kappa + 1 \right\},$$

enclosed by the four paths  $\Gamma^\tau$ ,  $\Gamma_{\theta, \kappa}^{(\tau)}$  and  $\mathbb{R} \pm i\pi/\tau$ , where  $\Gamma_{\theta, \kappa}^{(\tau)} = \{z \in \Gamma_{\theta, \kappa} : |\operatorname{Im}(z)| \leq \frac{\pi}{\tau}\}$ . Then Cauchy's theorem allows us to deform the integration path from  $\Gamma^\tau$  to  $\Gamma_{\theta, \kappa}^{(\tau)}$  in the integral (B.6) (the integrals on  $\mathbb{R} \pm i\pi/\tau$  cancels each other). This yields the desired representation (B.2).

## C Some inequalities

In this appendix, we prove the following two inequalities:

$$C_0^\# |z|^\tau \leq |1 - e^{z\tau}| \leq C_1^\# |z|^\tau, \quad \forall z \in \Gamma_{\theta, \kappa}^{(\tau)}, \tag{C.1}$$

$$|1 - e^{z\tau}| \leq C |z|^{1/q} \tau^{1/q}, \quad \forall z \in \Gamma_{\theta, \kappa}, \quad 1 \leq q \leq \infty, \tag{C.2}$$

which have been used in (3.27), (4.26) and (4.32).

**Proof of (C.1).** Note that

$$\begin{aligned}
\Gamma_{\theta, \kappa}^{(\tau)} &= \{z \in \mathbb{C} : |z| = \kappa, |\arg z| \leq \theta\} \cup \left\{ z \in \mathbb{C} : z = \rho e^{\pm i\theta}, \rho \geq \kappa, |\operatorname{Im}(z)| \leq \frac{\pi}{\tau} \right\} \\
&=: \Gamma_{\theta, \kappa}^{(\tau), 1} \cup \Gamma_{\theta, \kappa}^{(\tau), 2}.
\end{aligned} \tag{C.3}$$

For  $z \in \Gamma_{\theta, \kappa}^{(\tau)}$  we have  $|z|^\tau \leq \pi/\sin(\theta)$ . Since  $|z|^\tau$  is bounded, the following Taylor expansion holds:

$$1 - e^{z\tau} = -z\tau + O(|z|^2 \tau^2), \tag{C.4}$$

which implies

$$|1 - e^{z\tau}| \leq C_1^+ |z|\tau, \quad \text{if } z \in \Gamma_{\theta, \kappa}^{(\tau)}.$$

This proves the right-half inequality of (C.1).

From (C.4) we also see that there exists a small constant  $\gamma$  such that

$$C_0^+ |z|\tau \leq |1 - e^{z\tau}|, \quad \text{if } z \in \Gamma_{\theta, \kappa}^{(\tau)}, \quad |z|\tau < \gamma. \quad (\text{C.5})$$

If  $|z|\tau \geq \gamma$ , then the following inequality holds for  $\theta$  satisfying the condition of Lemma 1:

$$\gamma \leq |z|\tau \leq \frac{\pi}{\sin(\theta)} \leq \pi \sqrt{1 + 4/\pi^2} \leq \frac{3}{2}\pi.$$

Since the function  $g(w) := |1 - e^w|$  is not zero for  $\gamma \leq |w| \leq \frac{3}{2}\pi$ , the function  $g(w)$  must have a positive minimum value  $\varpi$  for  $\gamma \leq |w| \leq \frac{3}{2}\pi$ , i.e.,  $g(w) \geq \varpi$ . Consequently, we have

$$\frac{\sin(\theta)}{\pi} |z|\tau \leq \varpi \leq |1 - e^{z\tau}|, \quad \text{if } z \in \Gamma_{\theta, \kappa}^{(\tau)}, \quad |z|\tau \geq \gamma, \quad (\text{C.6})$$

where we have used the inequality  $\frac{\sin(\theta)}{\pi} |z|\tau \leq 1$  in the last inequality. Combining (C.5) and (C.6) yields (C.1).  $\square$

**Proof of (C.2).** If  $z \in \Gamma_{\theta, \kappa}$  and  $|z|\tau \leq \pi/\sin(\theta)$ , then  $z \in \Gamma_{\theta, \kappa}^{(\tau)}$ . In this case, (C.1) implies

$$\begin{aligned} |1 - e^{z\tau}| &\leq C |z|\tau & \forall z \in \Gamma_{\theta, \kappa}, \quad |z|\tau \leq \pi/\sin(\theta), \\ |1 - e^{z\tau}| &\leq C & \forall z \in \Gamma_{\theta, \kappa}, \quad |z|\tau \leq \pi/\sin(\theta). \end{aligned}$$

The combination of the two inequalities above yields

$$|1 - e^{z\tau}| \leq C |z|^{1/q} \tau^{1/q} \quad \forall z \in \Gamma_{\theta, \kappa}, \quad |z|\tau \leq \pi/\sin(\theta). \quad (\text{C.7})$$

If  $z \in \Gamma_{\theta, \kappa}$  and  $|z|\tau \geq \pi/\sin(\theta)$ , then

$$|e^{z\tau}| = e^{-|z|\tau \cos(\theta)} \leq e^{-\pi/\tan(\theta)},$$

which implies

$$|1 - e^{z\tau}| \leq 1 + e^{-\pi/\tan(\theta)} \leq 2 \leq 2 \left( \frac{\sin(\theta)}{\pi} \right)^{\frac{1}{q}} |z|^{1/q} \tau^{1/q} \quad \forall z \in \Gamma_{\theta, \kappa}, \quad |z|\tau \geq \pi/\sin(\theta). \quad (\text{C.8})$$

Combining (C.7) and (C.8) yields (C.2).  $\square$

## References

1. E. J. Allen, S. J. Novosel, and Z. Zhang. Finite element and difference approximation of some linear stochastic partial differential equations. *Stochastics and Stochastics Reports*, 64, 1998.
2. R. Anton, D. Cohen, S. Larsson, and X. Wang. Full discretization of semilinear stochastic wave equations driven by multiplicative noise. *SIAM J. Numer. Anal.*, 54(2):1093–1119, 2016.
3. W. Arendt, C. J. Batty, M. Hieber, and F. Neubrander. *Vector-valued Laplace Transforms and Cauchy Problems, Second Edition*. Birkhäuser, second edition, 2011.
4. L. Bañas, Z. Brzeźnik, and A. Prohl. Computational studies for the stochastic Landau–Lifshitz–Gilbert equation. *SIAM J. Sci. Comput.*, 35(1):B62–B81, 2013.
5. J. Bergh and J. Löfström. *Interpolation Spaces: An Introduction*. Springer-Verlag Berlin Heidelberg, 1976.
6. P. Clément and G. Da Prato. Some results on stochastic convolutions arising in Volterra equations perturbed by noise. *Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl.*, 7(3):147–153, 1996.
7. E. Cuesta, C. Lubich, and C. Palencia. Convolution quadrature time discretization of fractional diffusion-wave equations. *Math. Comp.*, 75(254):673–696, 2006.
8. G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*. Encyclopedia of Mathematics and its Applications. Cambridge University Press, second edition, 2014.
9. Q. Du and T. Zhang. Numerical approximation of some linear stochastic partial differential equations driven by special additive noises. *SIAM J. Numer. Anal.*, 40(4):1421–1445, 2002.
10. X. Feng, Y. Li, and A. Prohl. Finite element approximations of the stochastic mean curvature flow of planar curves of graphs. *Stoch. PDE: Anal. Comp.*, 2(1):54–83, 2014.
11. L. Grafakos. *Classical Fourier Analysis*. Graduate Texts in Mathematics. Springer New York, 2008.

12. M. Gunzburger, B. Li, and J. Wang. Sharp convergence rates of time discretization for stochastic time-fractional PDEs subject to additive space-time white noise. *Math. Comp.*, DOI: 10.1090/mcom/3397.
13. B. Jin, R. Lazarov, and Z. Zhou. Two fully discrete schemes for fractional diffusion and diffusion-wave equations with nonsmooth data. *SIAM J. Sci. Comput.*, 38(1):A146–A170, 2016.
14. B. Jin, B. Li, and Z. Zhou. Correction of high-order BDF convolution quadrature for fractional evolution equations. *SIAM J. Sci. Comput.*, 39(6):A3129–A3152., 2017.
15. B. Jin, B. Li, and Z. Zhou. An analysis of the Crank-Nicolson method for subdiffusion. *IMA J. Numer. Anal.*, 38(1):518–541, 2018.
16. A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo. *Theory and Application of Fractional Differential Equations*. Elsevier Science B.V., Amsterdam, 2006.
17. M. Kovács and J. Printems. Strong order of convergence of a fully discrete approximation of linear stochastic Volterra type evolution equation. *Math. Comp.*, 83(289):2325–2346, 2014.
18. A. Laptev. Dirichlet and Neumann eigenvalue problems on domains in Euclidean spaces. *J. Functional Analysis*, 151:531–545, 1997.
19. P. Li and S. Yau. On the Schrödinger equation and the eigenvalue problem. *Commun. Math. Phys.*, 88(3):309–318, 1983.
20. C. Lubich. Convolution quadrature and discretized operational calculus. I. *Numer. Math.*, 52(2):129–145, 1988.
21. C. Lubich. Convolution quadrature and discretized operational calculus. II. *Numer. Math.*, 52(4):413–425, 1988.
22. C. Lubich, I. H. Sloan, and V. Thomée. Nonsmooth data error estimates for approximations of an evolution equation with a positive-type memory term. *Math. Comp.*, 65(213):1–17, 1996.
23. J. B. Mijena and E. Nane. Space-time fractional stochastic partial differential equations. *Stochastic Process. Appl.*, 125:3301–3326, 2015.
24. J. M. Sanz-Serna. A numerical method for a partial integro-differential equation. *SIAM J. Numer. Anal.*, 25(2):319–327, 1988.
25. T. Shardlow. Numerical methods for stochastic parabolic PDEs. *Numer. Funct. Anal. And Optimiz.*, 20:121–145, 1999.
26. W. A. Strauss. *Partial differential equations: an introduction*. John Wiley & Sons, Inc., 2008.
27. V. Thomée. *Galerkin Finite Element Methods for Parabolic Problems*. Springer-Verlag, Berlin, 2nd edition, 2006.
28. Y. Yan. Galerkin finite element methods for stochastic parabolic partial differential equations. *SIAM J. Numer. Anal.*, 43(4):1363–1384, 2005.