



Research article

Global existence and stability of three species predator-prey system with prey-taxis

Gurusamy Arumugam *

Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Hong Kong S.A.R., China

* **Correspondence:** Email: gurusamy.arumugam@polyu.edu.hk.

Abstract: In this paper, we study the following initial-boundary value problem of a three species predator-prey system with prey-taxis which describes the indirect prey interactions through a shared predator, i.e.,

$$\begin{cases} u_t = d\Delta u + u(1 - u) - \frac{a_1uw}{1 + a_2u + a_3v}, & \text{in } \Omega, t > 0, \\ v_t = \eta d\Delta v + rv(1 - v) - \frac{a_4vw}{1 + a_2u + a_3v}, & \text{in } \Omega, t > 0, \\ w_t = \nabla \cdot (\nabla w - \chi_1 w \nabla u - \chi_2 w \nabla v) - \mu w + \frac{a_5uw}{1 + a_2u + a_3v} + \frac{a_6vw}{1 + a_2u + a_3v}, & \text{in } \Omega, t > 0, \end{cases}$$

under homogeneous Neumann boundary conditions in a bounded domain $\Omega \subset \mathbb{R}^n (n \geq 1)$ with smooth boundary, where the parameters $d, \eta, r, \mu, \chi_1, \chi_2, a_i > 0, i = 1, \dots, 6$. We first establish the global existence and uniform-in-time boundedness of solutions in any dimensional bounded domain under certain conditions. Moreover, we prove the global stability of the prey-only state and coexistence steady state by using Lyapunov functionals and LaSalle’s invariance principle.

Keywords: Predator-prey system; prey-taxis; local existence; global existence; global stabilization

1. Introduction

Predator-prey models were developed to describe the dynamics of interactions between prey and predator species. The relationship between prey and predator has been explored in recent years due to its importance in ecology. In addition to the differential operators in the predator-prey system, predators also move toward the higher prey density, which is so-called the prey-taxis, and it plays an important role in pest control and biological control [1–4]. The first predator-prey model with prey-taxis was

derived by Kareiva and Odell [5] to describe the predator aggregation phenomenon:

$$\begin{cases} u_t = \nabla \cdot (d(w)\nabla u) - \nabla \cdot (u\chi(w)\nabla w) + F(u, w), \\ w_t = d\Delta w + G(u, w), \end{cases} \quad (1.1)$$

where $u = u(x, t)$ and $w = w(x, t)$ denote the predator and prey densities, respectively, and the term $\nabla \cdot (d(w)\nabla u)$ denotes the diffusion of predators with diffusion coefficient $d(w)$. The term $-\nabla \cdot (u\chi(w)\nabla w)$ represents the prey-taxis with $\chi(w)$ as prey-taxis coefficient. The parameter $d > 0$ is the diffusion coefficient of prey. The typical form of the functions $F(u, w) = au f(w) + h(u)$ and $G(u, w) = g(w) - bu f(w)$, where $f(w)$ represents the functional response, for numerous functional response functions (see [6–8]) and the parameters $a, b \in \mathbb{R}$ describe the inter-specific interactions between predator-preys. The intra-specific interactions of predators and prey are described by the functions $h(u)$ and $g(w)$, respectively. The results related to variants of the above prey-taxis system have been studied by many authors, as one can refer to [9–18], and nonlinear prey-taxis [19–24]. Moreover, the predator-prey system with prey-taxis and liquid surroundings was considered in [25], and proved global existence and large time behavior of solutions by using L^p estimates and Lyapunov functionals, respectively.

In this paper, we consider a PDE model of indirect interactions between two prey species and a shared predator with homogeneous Neumann boundary conditions:

$$\begin{cases} u_t = d_1\Delta u + \alpha_1 u \left(1 - \frac{u}{K_u}\right) - \frac{c_1 u w}{1 + c_1 T_1 u + c_2 T_2 v}, & \text{in } \Omega, t > 0, \\ v_t = d_2\Delta v + \alpha_2 v \left(1 - \frac{v}{K_v}\right) - \frac{c_2 v w}{1 + c_1 T_1 u + c_2 T_2 v}, & \text{in } \Omega, t > 0, \\ w_t = \nabla \cdot (d_3 \nabla w - \chi_1 w \nabla u - \chi_2 w \nabla v) - d w \\ + \frac{c_1 \gamma_1 u w}{1 + c_1 T_1 u + c_2 T_2 v} + \frac{c_2 \gamma_2 v w}{1 + c_1 T_1 u + c_2 T_2 v}, & \text{in } \Omega, t > 0, \\ \partial_\nu u = \partial_\nu v = \partial_\nu w = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x) \text{ and } w(x, 0) = w_0(x), & x \in \Omega. \end{cases} \quad (1.2)$$

Where, $\Omega \subset \mathbb{R}^n (n \geq 1)$ is a bounded domain with smooth boundary $\partial\Omega$ and $\frac{\partial}{\partial \nu}$ represents the derivative with respect to outer normal of $\partial\Omega$, u is the native prey, v denotes the invasive prey and w is the predator species. The parameters d_1, d_2 denote the random diffusion rates of prey, and d_3 and d_4 denote the random diffusion rate of predators and the chemical concentration, respectively. K_u and K_v are the carrying capacities for these prey species. The constants α_1 and α_2 are intrinsic growth rate parameters. The parameters T_1 and T_2 usually represent the handling time required for catching and consuming a unit of prey type u and v , respectively. The constants c_1 and c_2 are capture rates per unit prey density while the predator is searching. In particular, c_1 is the capture rate of prey u and c_2 is the capture rate of prey v . In addition, d is an intrinsic growth (death) rate for the predator and η is a self-limiting or crowding coefficient for the predator. κ is the production rate of chemical signal per individual prey u and ξ is the decay rate of the chemical signal. The positive constants γ_1 and γ_2 denote the transformation rates of the predator.

The terms $-\nabla \cdot (\chi_1 w \nabla u)$ and $-\nabla \cdot (\chi_2 w \nabla v)$ denote the tendency of predators moves towards the high density of prey. The parameters χ_1 and χ_2 are the prey-taxis coefficients. The functions $\frac{c_1 u}{1 + c_1 T_1 u + c_2 T_2 v}$ and $\frac{c_2 v}{1 + c_1 T_1 u + c_2 T_2 v}$ these represent Holling type II functional responses for two preys that are consumed in a single habitat, so that handling one prey reduces the time available to capture the other.

Let $\tilde{u} = \frac{u}{K_u}$, $\tilde{v} = \frac{v}{K_v}$, $\tilde{w} = dw$, $d = \frac{d_1}{d_3}$, $\eta = \frac{d_2}{d_1}$, $L = \sqrt{\frac{d_3}{a_1}}$, $T = \frac{L^2}{d_3} = \frac{1}{a_1}$, $\tilde{t} = \frac{t}{T}$, $\tilde{x} = \frac{x}{L}$, $\tilde{y} = \frac{y}{L}$, $r = \alpha_2 T a_1 = c_1 T d$, $a_2 = c_1 T_1 K_u$, $a_3 = c_2 T_2 K_v$, $a_4 = c_2 d T$, $a_5 = c_1 \gamma_1 K_u d$, $a_6 = c_2 \gamma_2 K_v d$. Then, substituting these parameters into system (1.2) and dropping the tilde notation, we get a nondimensional system as follows:

$$\begin{cases} u_t = d\Delta u + u(1-u) - \frac{a_1 u w}{1 + a_2 u + a_3 v}, & \text{in } \Omega, t > 0, \\ v_t = \eta d\Delta v + rv(1-v) - \frac{a_4 v w}{1 + a_2 u + a_3 v}, & \text{in } \Omega, t > 0, \\ w_t = \nabla \cdot (\nabla w - \chi_1 w \nabla u - \chi_2 w \nabla v) - \mu w + \frac{a_5 u w}{1 + a_2 u + a_3 v} + \frac{a_6 v w}{1 + a_2 u + a_3 v}, & \text{in } \Omega, t > 0, \\ \partial_\nu u = \partial_\nu v = \partial_\nu w = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x) \text{ and } w(x, 0) = w_0(x), & x \in \Omega. \end{cases} \quad (1.3)$$

Let us recall some existing works on three species predator-prey systems with prey-taxis. Very recently, Haskel and Bell [26] proved the existence of positive classical solutions for the two-prey one-predator system with prey competitions and prey taxis. In addition, they also established the pattern formation by using bifurcation analysis. Further, they also studied the bifurcation analysis of two competing prey with one shared predator model by using the theories of Crandall-Rabinowitz and Hopf bifurcation in [27]. The steady-state bifurcation analysis of the two-prey one-predator model with two prey taxis was studied by Xu et al. [28]. Jin et al. [29] considered the three-species food chain model in a two-dimensional bounded domain, and they also proved the global existence of classical solutions and global stability of constant steady states. The traveling wave solutions for a nonlocal dispersal predator-prey system with one predator and two prey was studied in [30]. Amorim et al. [31] studied the boundedness and global well-posedness of the spatio-temporal evolution of two competitive prey, and one predator model with the intra-specific competition. The global existence and boundedness of classical solutions for the two-predators and one-prey with competition in a bounded domain with Neumann boundary conditions were proved by Min et al. in [32]. Recently, the global existence of weak solutions to the two-prey one-predator system with prey-taxis, and competition between prey was proved in any dimension in [33]. For the similar mathematical structure of (1.3), we refer to [34–36]. Throughout this paper, we assume the system parameters are positive. To the best of author's knowledge, there is no article that discusses the well-posedness of the considered system (1.3). The main purpose of this article is to discuss the global dynamics of the system (1.3) in any dimension ($n \geq 1$). In particular, we first prove the global existence of a classical solution in all dimensions, and then we investigate the global stability of steady states. Our main result regarding the global existence of classical solutions with uniform-in-time bound is stated below.

Theorem 1.1. (Global existence) *Let $\Omega \subset \mathbb{R}^n$, $n \geq 1$ be a bounded domain with smooth boundary and let $d, \eta > 0$, $h_1, h_2 > 1$, $k \geq 2$, $a_i > 0$, $i = 1, \dots, 6$, $\mu > 0$, $K_0 = \max\{1, \|u_0\|_{L^\infty}\}$ and $K_1 = \max\{1, \|v_0\|_{L^\infty}\}$. For any $(u_0, v_0, w_0) \in [W^{1,p}(\Omega)]^3$ with $p > n$ and $u_0, v_0, w_0 \geq 0$ ($\neq 0$), if $d > \max\{5kK_0, \frac{5kK_1}{\eta}\}$ and χ_1 satisfies*

$$\chi_1 \leq \min \left\{ \frac{d}{5kK_0(d+1)}, \frac{d}{5kK_0} - 1, \frac{d}{5kK_0(d+1)\sqrt{h_2}}, \frac{2(\eta d + 1)\sqrt{dK_1}}{K_0(d+1)\sqrt{5k\eta h_2}} \right\} \quad (1.4)$$

and χ_2 satisfies

$$\chi_2 \leq \min \left\{ \frac{\eta d}{5kK_1 \sqrt{h_1}(\eta d + 1)}, \frac{2\eta(d+1) \sqrt{dK_0}}{\sqrt{5kh_1}(\eta d + 1)K_1}, \frac{\eta d}{5kK_1(\eta d + 1)}, \frac{\eta d}{5kK_1} - 1 \right\}, \quad (1.5)$$

then there exists a unique global classical solution $(u, v, w) \in [C^0(\bar{\Omega} \times [0, \infty)) \cap C^{2,1}(\bar{\Omega} \times (0, \infty))]^3$ solving the problem (1.3). Moreover, the solution satisfies $u, v, w > 0$ for all $t > 0$ and

$$\|u(x, t)\|_{L^\infty} + \|v(x, t)\|_{L^\infty} + \|w(x, t)\|_{L^\infty} \leq C \text{ for all } t > 0,$$

where $C > 0$ is a constant independent of t .

Next, we shall study the large time behaviour of the constant steady states (u_s, v_s, w_s) of the system (1.3) solving the following system

$$\begin{cases} u_s \left[1 - u_s - \frac{a_1 w_s}{1 + a_2 u_s + a_3 v_s} \right] = 0, \\ v_s \left[r(1 - v_s) - \frac{a_4 w_s}{1 + a_2 u_s + a_3 v_s} \right] = 0, \\ w_s \left[\frac{a_5 u_s}{1 + a_2 u_s + a_3 v_s} + \frac{a_6 v_s}{1 + a_2 u_s + a_3 v_s} - \mu \right] = 0. \end{cases}$$

If we solve the above system, we will find the following steady states

$$(u_s, v_s, w_s) = \begin{cases} (0, 0, 0) \text{ or } (1, 0, 0) \text{ or } (0, 1, 0) \text{ or } (1, 1, 0), & \text{if } \mu > \frac{a_5 + a_6}{1 + a_2 + a_3}, a_4 < \frac{a_1 r(a_5 + a_3 a_5 - a_2 a_6)}{a_3 a_5 - a_6 - a_2 a_6}, \\ (0, 0, 0) \text{ or } (1, 0, 0) \text{ or } (0, 1, 0) \text{ or } (1, 1, 0) \text{ or } E_*^1, & \text{if } \mu > \frac{a_5 + a_6}{1 + a_2 + a_3}, a_4 < \frac{a_1 r(a_5 + a_3 a_5 - a_2 a_6)}{a_3 a_5 - a_6 - a_2 a_6}, \\ & \mu < \frac{a_5}{1 + a_2}, \\ (0, 0, 0) \text{ or } (1, 0, 0), (0, 1, 0), (1, 1, 0), E_*^2, & \text{if } \mu > \frac{a_5 + a_6}{1 + a_2 + a_3}, a_4 < \frac{a_1 r(a_5 + a_3 a_5 - a_2 a_6)}{a_3 a_5 - a_6 - a_2 a_6}, \\ & \mu < \frac{a_6}{1 + a_3}, \\ (0, 0, 0) \text{ or } (1, 0, 0) \text{ or } (0, 1, 0) \text{ or } (1, 1, 0) \text{ or } E_*^1 \text{ or } E_*^2 \text{ or } E_* = (u_*, v_*, w_*) & \\ & \mu < \frac{a_5 + a_6}{1 + a_2 + a_3}, a_4 > \frac{a_1 r(a_5 + a_3 a_5 - a_2 a_6)}{a_3 a_5 - a_6 - a_2 a_6}, \end{cases} \quad (1.6)$$

where

$$\begin{aligned} E_*^1 &= \left(\frac{\mu}{a_5 - a_2 \mu}, 0, \frac{a_5(a_5 - (1 + a_2)\mu)}{a_1(a_5 - a_2 \mu)^2} \right), \\ E_*^2 &= \left(0, \frac{\mu}{a_6 - a_3 \mu}, \frac{a_6 r(a_6 - (1 + a_3)\mu)}{a_4(a_6 - a_3 \mu)^2} \right) \\ u_* &= \frac{a_4(a_6 - a_3 \mu) + a_1 r(-a_6 + \mu + a_3 \mu)}{a_1 r(a_5 - a_2 \mu) + a_4(a_6 - a_3 \mu)} \\ v_* &= \frac{a_1 r(a_5 - a_2 \mu) + a_4(-a_5 + \mu + a_2 \mu)}{a_1 r(a_5 - a_2 \mu) + a_4(a_6 - a_3 \mu)} \end{aligned}$$

$$w_* = \frac{-r[(1+a_2)a_4a_6 + a_1(a_5 - a_2a_6)r + a_3(-a_4a_5 + a_1a_5r)](-a_5 - a_6 + (1+a_2+a_3)\mu)}{(a_1r(a_5 - a_2\mu) + a_4(a_6 - a_3\mu))^2}$$

and $(0, 0, 0)$ is the extinction steady state, $(1, 0, 0)$ is the prey u only steady state, $(0, 1, 0)$ is the prey v only steady state. E_*^1 and E_*^2 denote the semi-coexistence steady state. Finally, E_* denotes the coexistence steady state. Next, we shall explore the following question: which of the above seven homogeneous steady states will be asymptotically stable? As we know that the global stability of the cross-diffusion system is difficult and many techniques are not available, we try to use the Lyapunov functionals to prove the global stability of the homogeneous steady states under some conditions. Next, we state our stability results as in the following theorem:

Theorem 1.2. (Global stability) *Assume the conditions in Theorem 1.1 hold. Let (u, v, w) be the solution of (1.3) obtained in Theorem 1.1 and let $K_0 = \max\{1, \|u_0\|_{L^\infty}\}$, $K_1 = \max\{1, \|v_0\|_{L^\infty}\}$, $\Gamma_1 = \frac{a_5 + (a_3a_5 - a_2a_6)v_*}{a_1(1+a_2u_*+a_3v_*)}$ and $\Gamma_2 = \frac{a_5 + (a_3a_5 - a_2a_6)u_*}{a_1(1+a_2u_*+a_3v_*)}$. Then the following results hold true.*

- If $\mu > \frac{a_5 + a_6}{1 + a_2 + a_3}$, $a_4 \leq \frac{a_1r(a_5 + a_3a_5 - a_2a_6)}{a_3a_5 - a_6 - a_2a_6}$, then the steady state $(1, 1, 0)$ is globally asymptotically stable.
- If $\mu < \frac{a_5 + a_6}{1 + a_2 + a_3}$, $a_4 > \frac{a_1r(a_5 + a_3a_5 - a_2a_6)}{a_3a_5 - a_6 - a_2a_6}$, the steady state E_* defined by (1.6) is globally asymptotically stable provided

$$\frac{\Gamma_1(2a_2 + a_3)}{2} + \frac{\Gamma_2a_2r}{2} < \Gamma_1 + \frac{\Gamma_1(2a_2 + a_3)u_*}{2} + \frac{\Gamma_2a_2rv_*}{2}, \quad (1.7)$$

$$\frac{\Gamma_2r(2a_3 + a_2)}{2} + \frac{\Gamma_1a_3}{2} < r\Gamma_2 + \frac{\Gamma_2r(2a_3 + a_2)v_*}{2} + \frac{\Gamma_1a_3u_*}{2}, \quad (1.8)$$

$$4d\Gamma_1\Gamma_2\eta u_* v_* > \Gamma_1\chi_2^2 u_* w_* \|v\|_{L^\infty}^2 + \chi_1^2 \eta \Gamma_2 \|u\|_{L^\infty}^2 v_* w_*. \quad (1.9)$$

where $\|u\|_{L^\infty}$ and $\|v\|_{L^\infty}$ depends on d, η, a_1, a_2, a_3 but independent of χ_1, χ_2 .

The paper is organized as follows: In section 2, we first present some preliminary results, and then we state and prove the local existence of solutions of (1.3). Section 3 deals with the existence of globally bounded classical solutions as stated in Theorem 1.1. In Section 4, we establish the global stability results as stated in Theorem 1.2 by using the Lyapunov functionals and LaSalle's invariance principle.

2. Preliminaries and local existence

In what follows, we shall abbreviate $\int_\Omega f dx$ as $\int_\Omega f$ for simplicity. In this section, we first prove the local existence of classical solutions to the system (1.3) in any dimension $\Omega \subset \mathbb{R}^n$, $n \geq 1$ using the Amann's approach (cf. [37, 38]). We use the following Gagliardo-Nirenberg interpolation inequality in the sequel.

Lemma 2.1. ([39]) *There exists a constant $C_4 > 0$, such that for all $u \in W^{1,q}(\Omega)$,*

$$\|u\|_{L^p} \leq C_4 \|u\|_{W^{1,q}}^a \|u\|_{L^m}^{1-a}, \quad (2.1)$$

where $p, q \geq 1$ which satisfies $p(n-q) < nq$, $m \in (0, p)$ with $a = \frac{\frac{n}{m} - \frac{n}{p}}{\frac{n}{m} + 1 - \frac{n}{q}} \in (0, 1)$.

Lemma 2.2. ([39]) *There exists a constant $C_5 > 0$, such that for all $u \in W_{1,q}(\Omega)$, we have*

$$\|u\|_{W^{1,p}} \leq C_5 (\|\nabla u\|_{L^p} + \|u\|_{L^q}), \quad (2.2)$$

where $p > 1$ and $q > 0$.

Lemma 2.3. ([40]) Let $T > 0, \tau \in (0, T), \sigma \geq 0, a > 0, b \geq 0$, and suppose that $f : [0, T) \rightarrow [0, \infty)$ is absolutely continuous, and satisfies

$$f'(t) + af^{1+\sigma}(t) \leq h(t), \quad t \in \mathbb{R}, \quad (2.3)$$

where $h \geq 0, h(t) \in L^1_{loc}([0, T))$ and

$$\int_{t-\tau}^t h(s)ds \leq b, \quad \text{for all } t \in [\tau, T). \quad (2.4)$$

Then,

$$\sup_{t \in (0, T)} f(t) + a \sup_{t \in (\tau, T)} \int_{t-\tau}^t f^{1+\sigma}(s)ds \leq b + 2 \max\{f(0) + b + a\tau, \frac{b}{a\tau} + 1 + 2b + 2a\tau\}. \quad (2.5)$$

Lemma 2.4. ([39]) Assume that $m \in \{0, 1\}, p \in [1, \infty]$ and $q \in (1, \infty)$. Then there exists some positive constant C_1 , such that

$$\|\phi\|_{W^{m,p}} \leq C_1 \|(A+1)^\theta \phi\|_{L^q}, \quad (2.6)$$

for any $\phi \in D((A+1)^\theta)$, where $\theta \in (0, 1)$ satisfies

$$m - \frac{n}{p} < 2\theta - \frac{n}{q}.$$

If in addition $q \geq p$, then there exist constants $C_2 > 0$ and $\gamma > 0$, such that for any $\phi \in L^p(\Omega)$,

$$\|(A+1)^\theta e^{-t(A+1)} \phi\|_{L^q} \leq C_2 t^{-\theta - \frac{n}{2}(\frac{1}{p} - \frac{1}{q})} e^{-\mu t} \|\phi\|_{L^p}, \quad (2.7)$$

where the semigroup $\{e^{-t(A+1)}\}_{t \geq 0}$ maps $L^p(\Omega)$ into $D((A+1)^\theta)$. Moreover, for any $p \in (0, \infty)$ and $\epsilon > 0$, there exist constants $C_3 > 0$ and $\gamma > 0$, such that

$$\|(A+1)^\theta e^{-tA} \nabla \cdot \phi\|_{L^p} \leq C_3 t^{-\theta - \frac{1}{2} - \epsilon} e^{-\gamma t} \|\phi\|_{L^p} \quad (2.8)$$

this is valid for all \mathbb{R}^n -valued $\phi \in L^p(\Omega)$.

Theorem 2.1. (Local existence) Let the assumptions in Theorem 1.1 hold. Then, there exists a $T_{max} \in (0, \infty]$, such that the problem (1.3) has a unique classical solution

$$(u, v, w) \in \left(C^0(\overline{\Omega} \times [0, T_{max})) \cap C^{2,1}(\overline{\Omega} \times (0, T_{max})) \right)^3,$$

which satisfies $(u, v, w) > 0$ for all $t > 0$. Further,

$$\text{if } T_{max} < \infty, \text{ then } \limsup_{t \nearrow T_{max}} (\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{L^\infty} + \|w(\cdot, t)\|_{L^\infty}) = \infty. \quad (2.9)$$

Proof. Denote $\psi = (u, v, w)$. Then, the problem (1.3) can be written as

$$\begin{cases} \psi_t = \nabla \cdot (A(\psi) \nabla \psi) + F(\psi), & x \in \Omega, t > 0, \\ \partial_\nu \psi = 0, & x \in \partial\Omega, t > 0, \\ \psi(\cdot, 0) = (u_0, v_0, w_0), & x \in \Omega, \end{cases} \quad (2.10)$$

where

$$A(\psi) = \begin{bmatrix} d & 0 & 0 \\ 0 & \eta d & 0 \\ -\chi_1 w & -\chi_2 w & 1 \end{bmatrix} \text{ and } F(\psi) = \begin{bmatrix} \frac{-a_1 u w}{1 + a_2 u + a_3 v} \\ \frac{-a_4 v w}{1 + a_2 u + a_3 v} \\ \frac{a_5 u w^2 + a_6 v w}{1 + a_2 u + a_3 v} \end{bmatrix}. \quad (2.11)$$

Since the eigenvalues of $A(\psi)$ are positive, the system (1.3) is normally parabolic (cf. [37, 38]). Then, the application of Theorem 7.3 and Corollary 9.3 in [37] yields a $T_{max} > 0$, such that system (1.3) admits a unique solution $(u, v, w) \in [C^0(\overline{\Omega} \times [0, T_{max})) \cap C^{2,1}(\overline{\Omega} \times (0, T_{max}))]^3$. Next, we show the nonnegativity of (u, v, w) by using the maximum principle. To do so, we need to rewrite the third equation of the system (1.3) as follows:

$$\begin{cases} w_t = \Delta w + p_1(x, t) \cdot \nabla w + p_2(x, t)w = 0, & x \in \Omega, t \in (0, T_{max}), \\ \partial_\nu w = 0, & x \in \Omega, t \in (0, T_{max}), \\ w(x, 0) = w_0 \geq 0 \text{ in } x \in \Omega, \end{cases} \quad (2.12)$$

where $p_1(x, t) = \chi_1 \nabla u + \chi_2 \nabla v$ and $p_2(x, t) = \chi_1 \Delta u + \chi_2 \Delta v - \frac{a_5 u + a_6 v}{1 + a_2 u + a_3 v}$. Hence, we apply the maximum principle for parabolic equation with Neumann boundary condition to (2.12) and we get $w \geq 0$ for all $(x, t) \in \Omega \times (0, T_{max})$. In addition, we also obtain $w > 0$ by strong maximum principle since the initial function $w_0 \not\equiv 0$. In the same way, we can obtain that $u, v > 0$ for all $(x, t) \in \Omega \times (0, T_{max})$. Because $A(\psi)$ is lower triangular, (2.9) follows from Theorem 5.2 in [41]. \square

Lemma 2.5. *The solution (u, v, w) of the system (1.3) satisfies*

$$0 < u(x, t) \leq K_0 := \max\{\|u_0\|_{L^\infty(\Omega)}, 1\}, \quad \limsup_{t \rightarrow \infty} u(x, t) \leq 1, \quad (2.13)$$

$$0 < v(x, t) \leq K_1 := \max\{\|v_0\|_{L^\infty(\Omega)}, 1\}, \quad \limsup_{t \rightarrow \infty} v(x, t) \leq 1, \quad (2.14)$$

$$\|w(x, t)\|_{L^1(\Omega)} \leq K_2 := \frac{\delta}{a_1 a_4}, \quad (2.15)$$

where $K_0, K_1, K_2, \delta = \max\{a_4 a_5 \|u_0\|_{L^1} + a_1 a_6 \|v_0\|_{L^1} + a_1 a_4 \|w_0\|_{L^1}, \frac{c_2}{c_1}\}$ are positive constants independent of t .

Proof. The proof is similar to Lemma 2.2 in [21] but for reader's convenience, we provide the proof here. We have already proved that the solution (u, v, w) of the system (1.3) is non-negative. Using this fact, we have

$$\begin{cases} u_t - d\Delta u = u(1 - u) - \frac{a_1 u w}{1 + a_2 u + a_3 v} \leq u(1 - u), & x \in \Omega, t > 0, \\ \partial_\nu u = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), & x \in \Omega. \end{cases} \quad (2.16)$$

Let $u^*(t)$ be a solution to the following ODE problem

$$\begin{cases} \frac{du^*}{dt} = u^*(1 - u^*), & t > 0, \\ u^*(0) = \|u_0\|_{L^\infty}. \end{cases} \quad (2.17)$$

The solution of the above ODE satisfies $u^*(t) \leq K_0 = \max\{\|u_0\|_{L^\infty}, 1\}$, and in addition, $u^*(t)$ is a super solution of the following PDE problem

$$\begin{cases} U_t - d\Delta U = U(1 - U) & x \in \Omega, t > 0, \\ \partial_\nu U = 0, & x \in \partial\Omega, t > 0, \\ U(x, 0) = u_0(x), & x \in \Omega. \end{cases} \quad (2.18)$$

Hence, we have $U(x, t) \leq u^*(t)$ for all $(x, t) \in \overline{\Omega} \times (0, \infty)$. Using the strong maximum principle to the problem (2.18), we obtain $0 < U(x, t) \leq u^*(t)$ for all $(x, t) \in \overline{\Omega} \times (0, \infty)$. From (2.16)–(2.18), and using the comparison principle, we conclude that

$$0 < u(x, t) \leq U(x, t) \leq u^*(t) \leq K_0, \text{ for all } (x, t) \in \overline{\Omega} \times (0, \infty), \quad (2.19)$$

which yields (2.13). Similarly, we can also prove (2.14).

Multiplying the first, second and third equations of (1.3) by a_4a_5 , a_1a_6 and a_1a_4 , respectively, and adding the resulting equations and then integrating it over Ω , we get

$$\begin{aligned} \frac{d}{dt} \left(\int_{\Omega} a_4a_5u + a_1a_6v + a_1a_4w \right) &= \int_{\Omega} a_4a_5u(1 - u) + a_1a_6rv(1 - v) - a_1a_4\mu w \\ &= \int_{\Omega} a_4a_5u - \int_{\Omega} a_4a_5u^2 + a_1a_6rv - a_1a_6rv^2 - a_1a_4\mu w. \end{aligned} \quad (2.20)$$

Using Cauchy-Schwartz inequality, one has

$$\left(\int_{\Omega} u \right)^2 \leq \left(\int_{\Omega} u^2 \right) |\Omega|$$

which implies

$$- \int_{\Omega} u^2 \leq -\frac{1}{|\Omega|} \left(\int_{\Omega} u \right)^2. \quad (2.21)$$

Using Young's inequality ($-a^2 - b^2 \leq -2ab, a, b > 0$) yields

$$- \int_{\Omega} u^2 \leq -2 \int_{\Omega} u + |\Omega|. \quad (2.22)$$

Substituting (2.22) into (2.20), we have that

$$\begin{aligned}
\frac{d}{dt} \left(\int_{\Omega} a_4 a_5 u + a_1 a_6 v + a_1 a_4 w \right) &\leq \int_{\Omega} a_4 a_5 u - 2 \int_{\Omega} a_4 a_5 u + a_4 a_5 |\Omega| + \int_{\Omega} a_1 a_6 r v \\
&\quad - 2 a_1 a_6 r \int_{\Omega} v + a_1 a_6 |\Omega| - a_1 a_4 \mu \int_{\Omega} w \\
&\leq - \int_{\Omega} a_4 a_5 u - \int_{\Omega} a_1 a_6 r v - \int_{\Omega} a_1 a_4 \mu w + a_4 a_5 |\Omega| + a_1 a_6 |\Omega|. \quad (2.23)
\end{aligned}$$

Set $y(t) = \int_{\Omega} a_4 a_5 u + a_1 a_6 v + a_1 a_4 w$ and choose $c_1 = \min\{1, r, \mu\}$ then (2.23) can be written as

$$y'(t) + c_1 y(t) \leq c_2, \quad (2.24)$$

where $c_2 = (a_4 a_5 + a_1 a_6) |\Omega|$ and which yields (2.15) with the help of Gronwall's inequality. We further have from (2.17) that $\lim_{t \rightarrow \infty} \sup u(x, t) \leq 1$. The proof of Lemma 2.5 is complete. \square

3. Global existence and boundedness

In this subsection, we prove the global existence and boundedness of solutions. In order to prove the global existence, we first derive a uniform bound for w in L^{n+1} by using a weight function argument and the proof is inspired from [23], which also concerns the predator-prey taxis with a single prey population. It is worth mentioning that the method was initially developed in [42].

Lemma 3.1. *Assume that χ_1 and χ_2 satisfy (1.4) and (1.5), respectively, and let (u, v, w) be the solution of (1.3). Then, there exists a positive constant $c_0 > 0$, such that*

$$\|w(\cdot, t)\|_{L^{n+1}(\Omega)} \leq c_0 \text{ for } t \in (0, T_{max}). \quad (3.1)$$

Proof. Let us define the constants and weight functions

$$k := n + 1, \beta_1 := \sqrt{\frac{(k-1)d}{10k}} \frac{1}{(d+1)K_0}, \beta_2 := \sqrt{\frac{(k-1)\eta d}{10k}} \frac{1}{(\eta d + 1)K_1}, \quad (3.2)$$

$$1 \leq \varphi_1(u) \leq e^{(\beta_1 K_0)^2} := h_1 > 1, 0 \leq u \leq K_0 \text{ and } 1 \leq \varphi_2(v) \leq e^{(\beta_2 K_1)^2} := h_2 > 1, 0 \leq v \leq K_1. \quad (3.3)$$

Now, by using the weight function and the first and second equation of (1.3), we obtain

$$\begin{aligned}
\frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_1(u) &= \int_{\Omega} w^{k-1} \varphi_1(u) w_t + \frac{1}{k} \int_{\Omega} w^k \varphi_1'(u) u_t \\
&= \int_{\Omega} w^{k-1} \varphi_1(u) \Delta w - \chi_1 \int_{\Omega} w^{k-1} \varphi_1(u) \nabla \cdot (w \nabla u) - \chi_2 \int_{\Omega} w^{k-1} \varphi_1(u) \nabla \cdot (w \nabla v) \\
&\quad + \int_{\Omega} w^k \varphi_1(u) \frac{a_5 u + a_6 v}{1 + a_2 u + a_3 v} - \mu \int_{\Omega} w^k \varphi_1(u) + \frac{1}{k} \int_{\Omega} w^k \varphi_1'(u) \Delta u \\
&\quad + \frac{1}{k} \int_{\Omega} w^k \varphi_1'(u) u(1-u) - \frac{1}{k} \int_{\Omega} w^{k+1} \varphi_1'(u) \frac{a_1 u}{1 + a_2 u + a_3 v} \\
&\leq - (k-1) \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \chi_1 (k-1) \int_{\Omega} w^{k-1} \nabla w \cdot \nabla u \varphi_1(u)
\end{aligned} \quad (3.4)$$

$$\begin{aligned}
& + \chi_1 \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2 + \chi_2 (k-1) \int_{\Omega} w^{k-1} \varphi_1(u) \nabla w \cdot \nabla v + \chi_2 \int_{\Omega} w^k \varphi_1'(u) \nabla u \cdot \nabla v \\
& + C \int_{\Omega} w^k \varphi_1(u) - d \int_{\Omega} w^{k-1} \varphi_1'(u) \nabla w \cdot \nabla u - \frac{d}{k} \int_{\Omega} w^k \varphi_1''(u) |\nabla u|^2 \\
& + \frac{1}{k} \int_{\Omega} w^k u \varphi_1'(u) + \frac{2\beta_1^2}{k} \int_{\Omega} w^k \varphi_1(u) u^2,
\end{aligned} \tag{3.5}$$

where we used the boundedness of the functional response, $C > 0$. The above inequality can be written as

$$\begin{aligned}
& \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_1(u) + (k-1) \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \frac{d}{k} \int_{\Omega} w^k \varphi_1''(u) |\nabla u|^2 \\
& \leq - (d+1) \int_{\Omega} w^{k-1} \varphi_1'(u) \nabla u \cdot \nabla w + \chi_1 (k-1) \int_{\Omega} w^{k-1} \nabla w \cdot \nabla u \varphi_1(u) + \chi_1 \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2 \\
& + \chi_2 (k-1) \int_{\Omega} w^{k-1} \varphi_1(u) \nabla w \cdot \nabla v + \chi_2 \int_{\Omega} w^k \varphi_1'(u) \nabla u \cdot \nabla v + C \int_{\Omega} w^k \varphi_1(u) \\
& + \frac{2\beta_1^2}{k} \int_{\Omega} w^k \varphi_1(u) u^2.
\end{aligned} \tag{3.6}$$

By using Young's inequality, we get

$$\begin{aligned}
- (d+1) \int_{\Omega} w^{k-1} \varphi_1'(u) \nabla u \cdot \nabla w & \leq \epsilon \frac{(d+1)}{2} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \frac{(d+1)}{2\epsilon} \int_{\Omega} w^k \frac{\varphi_1'^2(u)}{\varphi_1(u)} |\nabla u|^2 \\
& \leq \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \frac{(d+1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_1'^2(u)}{\varphi_1(u)} |\nabla u|^2,
\end{aligned} \tag{3.7}$$

and

$$\chi_1 (k-1) \int_{\Omega} w^{k-1} \varphi_1(u) \nabla w \cdot \nabla u \leq \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla u|^2, \tag{3.8}$$

$$\chi_2 (k-1) \int_{\Omega} w^{k-1} \varphi_1(u) \nabla w \cdot \nabla v \leq \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla v|^2. \tag{3.9}$$

Again,

$$\begin{aligned}
\chi_2 \int_{\Omega} w^k \varphi_1'(u) \nabla u \cdot \nabla v & \leq \frac{\epsilon}{2} \chi_2 \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2 + \frac{\chi_2}{2\epsilon} \int_{\Omega} w^k \varphi_1'(u) |\nabla v|^2 \\
& \leq \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2 + \frac{\chi_2^2}{4} \int_{\Omega} w^k \varphi_1'(u) |\nabla v|^2.
\end{aligned} \tag{3.10}$$

Substituting (3.7)–(3.10) into (3.6), one has

$$\begin{aligned}
& \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_1(u) + \frac{(k-1)}{4} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \frac{d}{k} \int_{\Omega} w^k \varphi_1''(u) |\nabla u|^2 \\
& \leq \frac{(d+1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_1'^2(u)}{\varphi_1(u)} |\nabla u|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla u|^2 + \chi_1 \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2
\end{aligned}$$

$$\begin{aligned}
& +\chi_1^2(k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla v|^2 + \chi_1 \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2 + \frac{\chi_2^2}{4\chi_1} \int_{\Omega} w^k \varphi_1'(u) |\nabla v|^2 \\
& + C \int_{\Omega} w^k \varphi_1(u) + \frac{2\beta_1^2}{k} \int_{\Omega} w^k \varphi_1(u) u^2.
\end{aligned} \tag{3.11}$$

Now, we multiply the third equation of (1.3) by $\varphi_2(v)$, we have

$$\begin{aligned}
\frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_2(v) & = \int_{\Omega} w^{k-1} \varphi_2(v) w_t + \frac{1}{k} \int_{\Omega} w^k \varphi_2'(v) v_t \\
& \leq \int_{\Omega} w^{k-1} \varphi_2(v) \Delta w - \chi_1 \int_{\Omega} w^{k-1} \varphi_2(v) \nabla \cdot (w \nabla u) - \chi_2 \int_{\Omega} w^{k-1} \varphi_2(v) \nabla \cdot (w \nabla v) \\
& \quad + C \int_{\Omega} w^k \varphi_2(v) + \frac{\eta d}{k} \int_{\Omega} w^k \varphi_2'(v) \Delta v + \frac{r}{k} \int_{\Omega} w^k \varphi_2'(v) v \\
& \leq -(k-1) \int_{\Omega} w^{k-2} \varphi_2(v) |\nabla w|^2 - (\eta d + 1) \int_{\Omega} w^{k-1} \varphi_2'(v) \nabla w \cdot \nabla v \\
& \quad + \chi_1(k-1) \int_{\Omega} w^{k-1} \varphi_2(v) \nabla w \cdot \nabla u + \chi_1 \int_{\Omega} w^k \varphi_2'(v) \nabla v \cdot \nabla u \\
& \quad + \chi_2(k-1) \int_{\Omega} w^{k-1} \varphi_2(v) \nabla w \cdot \nabla v + \chi_2 \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 + B_3 \int_{\Omega} w^k \varphi_2(v) \\
& \quad + \frac{2r\beta_2^2}{k} \int_{\Omega} w^k \varphi_2(v) v^2.
\end{aligned} \tag{3.12}$$

By using Young's inequality, we arrive at

$$-(\eta d + 1) \int_{\Omega} w^{k-1} \varphi_2'(v) \nabla w \cdot \nabla v \leq \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_2(v) |\nabla w|^2 + \frac{(\eta d + 1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_2'^2(v)}{\varphi_2(v)} |\nabla v|^2, \tag{3.13}$$

and

$$\chi_1(k-1) \int_{\Omega} w^{k-1} \varphi_2(v) \nabla w \cdot \nabla u \leq \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_2(v) |\nabla w|^2 + \chi_1^2(k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla u|^2 \tag{3.14}$$

$$\chi_2(k-1) \int_{\Omega} w^{k-1} \varphi_2(v) \nabla w \cdot \nabla v \leq \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_2(v) |\nabla w|^2 + \chi_2^2(k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla v|^2 \tag{3.15}$$

$$\begin{aligned}
\chi_1 \int_{\Omega} w^k \varphi_2'(v) \nabla u \cdot \nabla v & \leq \frac{\epsilon \chi_1}{2} \int_{\Omega} w^k \varphi_2'(v) |\nabla u|^2 + \frac{\chi_1}{2\epsilon} \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 \\
& \leq \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 + \frac{\chi_1^2}{4} \int_{\Omega} w^k \varphi_2'(v) |\nabla u|^2.
\end{aligned} \tag{3.16}$$

Substituting (3.13)–(3.16) into (3.12), we derive that

$$\begin{aligned}
& \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_2(v) + \frac{(k-1)}{4} \int_{\Omega} w^{k-2} \varphi_2(u) |\nabla w|^2 + \frac{\eta d}{k} \int_{\Omega} w^k \varphi_2''(v) |\nabla v|^2 \\
& \leq \frac{(\eta d + 1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_2''(v)}{\varphi_2(v)} |\nabla v|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla u|^2 + \chi_2 \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 \\
& \quad + \frac{\chi_1^2}{4\chi_2} \int_{\Omega} w^k \varphi_2'(v) |\nabla u|^2 + \chi_2^2 (k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla v|^2 + \chi_2 \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 + C \int_{\Omega} w^k \varphi_2(v) \\
& \quad + \frac{2r\beta_2^2}{k} \int_{\Omega} w^k \varphi_2(v) v^2. \tag{3.17}
\end{aligned}$$

Now, adding (3.11) and (3.17), the resulting inequality becomes

$$\begin{aligned}
& \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_1(u) + \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_2(v) + \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_2(v) |\nabla w|^2 \\
& \quad + \frac{d}{k} \int_{\Omega} w^k \varphi_1''(u) |\nabla u|^2 + \frac{d}{k} \int_{\Omega} w^k \varphi_2''(v) |\nabla v|^2 \\
& \leq \frac{(d+1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_1''(u)}{\varphi_1(u)} |\nabla u|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla u|^2 + (1 + \chi_1) \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2 \\
& \quad + \chi_2^2 (k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla v|^2 + \frac{\chi_2^2}{4} \int_{\Omega} w^k \varphi_1'(u) |\nabla v|^2 + B_3 \int_{\Omega} w^k \varphi_1(u) + \frac{2\beta_1^2}{k} \int_{\Omega} w^k \varphi_1(u) u^2 \\
& \quad + \frac{(\eta d + 1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_2''(v)}{\varphi_2(v)} |\nabla v|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla u|^2 + \chi_2^2 (k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla v|^2 \\
& \quad + (1 + \chi_2) \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 + \frac{\chi_1^2}{4} \int_{\Omega} w^k \varphi_2'(v) |\nabla u|^2 + C \int_{\Omega} w^k \varphi_2(v) + \frac{2r\beta_2^2}{k} \int_{\Omega} w^k \varphi_2(v) v^2. \tag{3.18}
\end{aligned}$$

Now, we do some computation to show that the terms involving $|\nabla u|^2$ and $|\nabla v|^2$ on the right-hand side of above inequality are dominated by $\int_{\Omega} w^k \varphi_1'' |\nabla u|^2$ and $\int_{\Omega} w^k \varphi_2'' |\nabla v|^2$, respectively. For $s \geq 0$, define

$$j_1(s) = \frac{(d+1)^2 \varphi_1''(u)}{(k-1) \varphi_1(u)} = \frac{4\beta_1^4 s^2 \varphi_1''(s)}{k-1}, \quad j_2(s) = \chi_1^2 (k-1) \varphi_1(s), \quad j_3(s) = 2(1 + \chi_1) \beta_1^2 s \varphi_1(s) \tag{3.19}$$

$$j_4(s) = \chi_1^2 (k-1) \varphi_2(s), \quad j_5(s) = \frac{\chi_1^2 \beta_2^2 s \varphi_2(s)}{2}, \quad j_6(s) = 2 \frac{d}{k} \beta_1^2 \varphi_1(s) + 4 \frac{d}{k} \beta_1^4 s^2 \varphi_1(s) \tag{3.20}$$

and

$$i_1(s) = \frac{4(\eta d + 1)^2 \beta_2^4 s^2 \varphi_2(s)}{(k-1)}, \quad i_2(s) = \chi_2^2 (k-1) \varphi_1(s), \quad i_3(s) = \frac{\chi_2^2 \beta_1^2 s \varphi_1(s)}{2} \tag{3.21}$$

$$i_4(s) = \chi_2^2 (k-1) \varphi_2(s), \quad i_5(s) = 2(1 + \chi_2) \beta_2^2 s \varphi_2(s). \tag{3.22}$$

Now, combining (3.19) and (3.20), one has

$$\frac{(d+1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_1''(u)}{\varphi_1(u)} |\nabla u|^2 + \chi_1^2 (k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla u|^2 + (1 + \chi_1) \int_{\Omega} w^k \varphi_1'(u) |\nabla u|^2$$

$$\begin{aligned}
& + \chi_1^2(k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla u|^2 + \frac{\chi_1^2}{4} \int_{\Omega} w^k \varphi_2'(v) |\nabla u|^2 \\
& \leq \frac{d}{k} \int_{\Omega} w^k \varphi_1''(u) |\nabla u|^2.
\end{aligned} \tag{3.23}$$

Similarly, we combine (3.21) and (3.22), we have that

$$\begin{aligned}
& \chi_2^2(k-1) \int_{\Omega} w^k \varphi_1(u) |\nabla v|^2 + \frac{\chi_2^2}{4} \int_{\Omega} w^k \varphi_1'(u) |\nabla v|^2 + \frac{(\eta d + 1)^2}{k-1} \int_{\Omega} w^k \frac{\varphi_2'^2(v)}{\varphi_2(v)} |\nabla v|^2 \\
& + \chi_2^2(k-1) \int_{\Omega} w^k \varphi_2(v) |\nabla v|^2 + (1 + \chi_2) \int_{\Omega} w^k \varphi_2'(v) |\nabla v|^2 \\
& \leq \frac{d}{k} \int_{\Omega} w^k \varphi_2''(v) |\nabla v|^2.
\end{aligned} \tag{3.24}$$

Substituting (3.23) and (3.24) into (3.18), one has

$$\begin{aligned}
& \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_1(u) + \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_2(v) + \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 + \frac{k-1}{4} \int_{\Omega} w^{k-2} \varphi_1(v) |\nabla w|^2 \\
& \leq B_3 \int_{\Omega} w^k \varphi_1(u) + \frac{2\beta_1^2}{k} \int_{\Omega} w^k \varphi_1(u) u^2 + B_3 \int_{\Omega} w^k \varphi_2(v) + \frac{2r\beta_2^2}{k} \int_{\Omega} w^k \varphi_2(v) v^2 \\
& \leq \left(C + \frac{2\beta_1^2 K_0^2}{k} \right) \int_{\Omega} w^k \varphi_1(u) + \left(B_3 + \frac{2r\beta_2^2 K_1^2}{k} \right) \int_{\Omega} w^k \varphi_2(v) \\
& \leq c_1 \int_{\Omega} w^k \varphi_1(u) + c_2 \int_{\Omega} w^k \varphi_2(v)
\end{aligned} \tag{3.25}$$

where $c_1 = C + \frac{2\beta_1^2 K_0^2}{k}$ and $c_2 = C + \frac{2r\beta_2^2 K_1^2}{k}$. Using Lemma 2.1, Lemma 2.2, and (3.3), we get the estimate

$$\begin{aligned}
\int_{\Omega} w^k \varphi_1(u) & \leq h_1 \int_{\Omega} w^k = h_1 \|w^{\frac{k}{2}}\|_{L^2}^2 \\
& \leq h_1 C_4 \|w^{\frac{k}{2}}\|_{W^{1,2}}^{2a} \|w^{\frac{k}{2}}\|_{L^{\frac{k}{2}}}^{2(1-a)} \\
& \leq h C_4 \left(C_5 \|\nabla w^{\frac{k}{2}}\|_{L^2} + \|w^{\frac{k}{2}}\|_{L^2} \right)^{2a} \|w^{\frac{k}{2}}\|_{L^{\frac{k}{2}}}^{2(1-a)} \\
& \leq h_1 C_4 C_5 \left(\|\nabla w^{\frac{k}{2}}\|_{L^2} + \|w^{\frac{k}{2}}\|_{L^1}^{\frac{k}{2}} \right)^{2a} \|w\|_{L^1}^{k(1-a)} \\
& \leq h_1 C_4 C_5 \left(\|\nabla u^{\frac{k}{2}}\|_{L^2} + K_2^{\frac{k}{2}} \right)^{2a} K_2^{k(1-a)} \\
& \leq C_6 \left(\|\nabla u^{\frac{k}{2}}\|_{L^2}^2 + 1 \right)^a,
\end{aligned} \tag{3.26}$$

where $a = \frac{\frac{kn}{2} - \frac{n}{2}}{\frac{kn}{2} + 1 - \frac{n}{2}} \in (0, 1)$. Now using the fact (3.3) and from (3.26), one can obtain

$$\begin{aligned}
\int_{\Omega} w^{k-2} \varphi_1(u) |\nabla w|^2 &\geq \int_{\Omega} w^{k-2} |\nabla w|^2 \\
&\geq \frac{4}{k^2} \int_{\Omega} |\nabla w^{\frac{k}{2}}|^2 \\
&\geq \frac{4}{k^2} \left[\frac{1}{C_6^{1/a}} \left(\int_{\Omega} w^k \varphi_1(u) \right)^{\frac{1}{a}} - 1 \right] \\
&\geq \frac{4}{k^2 C_6^{1/a}} \left(\int_{\Omega} w^k \varphi_1(u) \right)^{\frac{1}{a}} - \frac{4}{k^2}.
\end{aligned} \tag{3.27}$$

Similarly, we get

$$\int_{\Omega} w^{k-2} \varphi_2(v) |\nabla w|^2 \geq \int_{\Omega} w^{k-2} |\nabla w|^2 \geq \frac{4}{k^2 C_7^{1/a}} \left(\int_{\Omega} w^k \varphi_2(v) \right)^{\frac{1}{a}} - \frac{4}{k^2}. \tag{3.28}$$

From (3.27) and (3.28), we have

$$\frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_1(u) + \frac{1}{k} \frac{d}{dt} \int_{\Omega} w^k \varphi_2(v) \leq -\frac{(k-1)}{k^2 C_6^{1/a}} \left(\int_{\Omega} w^k \varphi_1(u) \right)^{\frac{1}{a}} - \frac{(k-1)}{k^2 C_7^{1/a}} \left(\int_{\Omega} w^k \varphi_2(v) \right)^{\frac{1}{a}} + \frac{2(k-1)}{k^2} \tag{3.29}$$

for all $t \in (0, T_{max})$ and where $\frac{1}{a} > 1$. Set $y(t) := \frac{1}{k} \int_{\Omega} w^k (\varphi_1(u) + \varphi_2(v))$. By using the inequality $x^p + y^p \geq n^{1-p} (x+y)^p$, $p \geq 1$, we get

$$y'(t) \leq -C_8 y^{\frac{1}{a}}(t) + \frac{2(k-1)}{k^2} \quad \text{for all } t \in (0, T_{max}), \tag{3.30}$$

where $C_8 := \min \left\{ \frac{(k-1)}{k^2 C_6^{1/a}}, \frac{(k-1)}{k^2 C_7^{1/a}} \right\} n^{1-\frac{1}{a}}$ and $\frac{1}{a} > 1$. By using Lemma 2.3 and the fact that (3.3), one has

$$\|w(\cdot, t)\|_{L^k} \leq \left(\int_{\Omega} w^k (\varphi_1(u) + \varphi_2(v)) \right)^{\frac{1}{k}} \leq C \tag{3.31}$$

for all $t \in (0, T_{max})$, where $C(u_0, v_0, C_8, \tau) > 0$. The proof of Lemma 3.1 is complete. \square

Lemma 3.2. *Let (u, v, w) be a solution of the system (1.3). Then, there exists a positive constant $c > 0$, such that*

$$\|w(\cdot, t)\|_{L^\infty} \leq c \text{ for all } t \in (0, T_{max}). \tag{3.32}$$

Proof. To obtain the L^∞ -bound of w , we use the semigroup estimates. In order to do this, we first obtain that for any $\tau \in (0, T_{max})$, there exists a constant $C > 0$, such that

$$\|(u(\cdot, t), v(\cdot, t))\|_{W^{1,\infty}} \leq C(\tau) \text{ for all } t \in (\tau, T_{max}) \tag{3.33}$$

Let $\tau \in (0, T_{max})$ be given such that $\tau < 1$ and also let $q := n + 1$ and $\theta \in \left(\frac{1}{2}(1 + \frac{n}{q}), 1\right)$. To begin with, we rewrite the first equation of (1.3) as follows:

$$u_t = d\Delta u - u + g(x, t), \quad (3.34)$$

with $g(x, t) := u(1 - u) - \frac{a_1 u w}{1 + a_2 u + a_3 v} + u$. Then, by Lemma 3.1 and the fact that $0 < u \leq K_0, 0 < v \leq K_1$ (see Lemma 2.5) and $\frac{a_1 \tilde{u}}{1 + a_2 u + a_3 v} \leq \tilde{K}, \tilde{K} > 0$, we have

$$\begin{aligned} \|g(x, t)\|_{L^q} &= \|u(1 - u) - \frac{a_1 u w}{1 + a_2 u + a_3 v} + u\|_{L^q} \\ &\leq K_0(1 + K_0)|\Omega|^{\frac{1}{q}} + a_1 \tilde{K} \|w(\cdot, t)\|_{L^q} + K_0 |\Omega|^{\frac{1}{q}} \\ &\leq [K_0(1 + K_0) + K_0] |\Omega|^{\frac{1}{q}} + a_1 \tilde{K} \|w(\cdot, t)\|_{L^q} \\ &\leq K_0[2 + K_0] |\Omega|^{\frac{1}{q}} + a_1 \tilde{K} \|w(\cdot, t)\|_{L^q} \\ &\leq K_0[2 + K_0] |\Omega|^{\frac{1}{q}} + a_1 \tilde{K} c_0. \end{aligned} \quad (3.35)$$

We apply the variation-of-constants formula to (3.34) and obtain

$$u(\cdot, t) = e^{-t(A_d+1)} u_0 + \int_0^t e^{-(A_d+1)(t-s)} g(\cdot, s) ds, \quad (3.36)$$

where $A_d = -d\Delta$. Then using (2.6), (2.7) and the estimate (3.35) in (3.36), one can derive

$$\begin{aligned} \|u(\cdot, t)\|_{W^{1,\infty}} &\leq C_1 \|(A_d + 1)^\theta u(\cdot, t)\|_{L^q} \\ &\leq C_1 t^{-\theta} e^{-\mu t} \|u_0\|_{L^q} + C_1 \int_0^t (t-s)^{-\theta} e^{-\mu(t-s)} \|g(\cdot, s)\|_{L^q} ds \\ &\leq C_1 t^{-\theta} e^{-\mu t} \|u_0\|_{L^q(\Omega)} + C_1 \int_0^t (t-s)^{-\theta} e^{-\mu(t-s)} [K_0[2 + K_0] |\Omega|^{\frac{1}{q}} + a_1 \tilde{K} c_0] ds \\ &\leq C_1 t^{-\theta} e^{-\mu t} \|u_0\|_{L^q(\Omega)} + C_1 [K_0[2 + K_0] |\Omega|^{\frac{1}{q}} + a_1 \tilde{K} c_0] \int_0^\infty \sigma^{-\theta} e^{-\mu \sigma} d\sigma \\ &\leq C_1 t^{-\theta} \|u_0\|_{L^q(\Omega)} + C_1 [K_0[2 + K_0] |\Omega|^{\frac{1}{q}} + a_1 \tilde{K} c_0] \mu^\theta \Gamma(1 - \theta) \\ &\leq C_1 t^{-\theta} + C_1 \end{aligned} \quad (3.37)$$

for all $t \in (\tau, T_{max})$, where $\Gamma(1 - \theta) > 0$. From the last inequality (3.37), we get the desired estimate

$$\|u(\cdot, t)\|_{W^{1,\infty}} \leq C_1 (\tau^{-\theta} + 1) := C(\tau) \text{ for all } t \in (\tau, T_{max}). \quad (3.38)$$

where C_1 is a generic constant which may vary line to line. Next, we obtain the bound for $\|v(\cdot, t)\|_{W^{1,\infty}}$. To this end, we rewrite the second equation of (1.3) as follows:

$$v_t = \eta d\Delta v - v + h(x, t), \quad (3.39)$$

with $h(x, t) := rv(1-v) - \frac{a_4vw}{1+a_2u+a_3v} + v$. Then, by Lemma 3.1 and the fact that $0 < u \leq K_0$, $0 < v \leq K_1$ (see Lemma 2.5) and $\frac{a_4v}{1+a_2u+a_3v} \leq \bar{K}$, $\bar{K} > 0$, we have

$$\begin{aligned} \|h(x, t)\|_{L^q} &= \|rv(1-v) - \frac{a_4vw}{1+a_2u+a_3v} + v\|_{L^q} \\ &\leq K_1(r(1+K_1)+1)|\Omega|^{\frac{1}{q}} + a_4\bar{K}\|w(\cdot, t)\|_{L^q} \\ &\leq K_1(r(1+K_1)+1)|\Omega|^{\frac{1}{q}} + a_4\bar{K}c_0. \end{aligned} \quad (3.40)$$

We apply the variation-of-constants formula to (3.34) and obtain

$$v(\cdot, t) = e^{-t(A_{\eta d}+1)}v_0 + \int_0^t e^{-(A_{\eta d}+1)(t-s)}h(\cdot, s)ds. \quad (3.41)$$

Then, using (2.6), (2.7) and the estimate (3.40) in (3.41), we find

$$\begin{aligned} \|v(\cdot, t)\|_{W^{1,\infty}} &\leq C_1\|(A_{\eta d}+1)^\theta v(\cdot, t)\|_{L^q} \\ &\leq C_1t^{-\theta}e^{-\mu t}\|v_0\|_{L^q} + C_1 \int_0^t (t-s)^{-\theta}e^{-\mu(t-s)}\|h(\cdot, s)\|_{L^q}ds \\ &\leq C_1t^{-\theta}e^{-\mu t}\|v_0\|_{L^q(\Omega)} + C_1 \int_0^t (t-s)^{-\theta}e^{-\mu(t-s)}[K_1(r(1+K_1)+1)|\Omega|^{\frac{1}{q}} + a_4\bar{K}c_0]ds \\ &\leq C_1t^{-\theta}e^{-\mu t}\|v_0\|_{L^q(\Omega)} + C_1[K_1(r(1+K_1)+1)|\Omega|^{\frac{1}{q}} + a_4\bar{K}c_0] \int_0^\infty \sigma^{-\theta}e^{-\mu\sigma}d\sigma \\ &\leq C_1t^{-\theta}\|v_0\|_{L^q(\Omega)} + C_1[K_1(r(1+K_1)+1)|\Omega|^{\frac{1}{q}} + a_4\bar{K}c_0]\mu^\theta\Gamma(1-\theta) \\ &\leq C_1t^{-\theta} + C_1 \end{aligned} \quad (3.42)$$

for all $t \in (\tau, T_{max})$ where $\Gamma(1-\theta) > 0$. From the last inequality (3.42), we obtain

$$\|v(\cdot, t)\|_{W^{1,\infty}} \leq C_1(\tau^{-\theta} + 1) := C(\tau) \text{ for all } t \in (\tau, T_{max}). \quad (3.43)$$

Next we derive the L^∞ -bound of $w(\cdot, t)$. We rewrite the third equation of (1.3) as follows:

$$w_t = \Delta w - w - \nabla \cdot (\chi_1 w \nabla u + \chi_2 w \nabla v) + \frac{a_5uw}{1+a_2u+a_3v} + \frac{a_6vw}{1+a_2u+a_3v} + w - \mu w. \quad (3.44)$$

Then, applying the variation-of-constants formula to (3.44), one has

$$\begin{aligned} w(\cdot, t) &= e^{-t(A+1)}w_0 - \chi_1 \int_0^t e^{-(t-s)(A+1)}\nabla \cdot (w\nabla u)ds - \chi_2 \int_0^t e^{-(t-s)(A+1)}\nabla \cdot (w\nabla v)ds \\ &\quad + \int_0^t e^{-(t-s)(A+1)}f(u, v, w)ds := I_1 + I_2 + I_3 + I_4, \end{aligned} \quad (3.45)$$

where $f(u, v, w) = \frac{a_5uw + a_6vw}{1+a_2u+a_3v} + (1-\mu)w$. Now, we take the L^∞ -norm on both sides of the above equation, we have

$$\|w(\cdot, t)\|_{L^\infty} \leq \|I_1\|_{L^\infty} + \|I_2\|_{L^\infty} + \|I_3\|_{L^\infty} + \|I_4\|_{L^\infty}. \quad (3.46)$$

First, we estimate the term I_1 as follows:

$$\|I_1\|_{L^\infty} \leq C_2 \tau^{-\theta} e^{-\mu t} \|u_0\|_{L^\infty} \leq C_2 \tau^{-\theta} \|u_0\|_{L^\infty} \quad (3.47)$$

for all $t \in (\tau, T_{max})$ and $\theta \in (\frac{n}{2q}, 1)$ and $\mu > 0$. In order to estimate the term I_2 , we set $m = 0$, $q = n + 1$, $p = \infty$ for (2.8) and we use (3.33) and (3.1), one has

$$\begin{aligned} \|I_2\|_{L^\infty} &\leq C_3 \chi_1 \int_0^t \|(A+1)^\theta e^{-(t-s)(A+1)} \nabla \cdot (w \nabla u)\|_{L^q} ds \\ &\leq C_3 \chi_1 \int_0^t e^{-(t-s)} \|(A+1)^\theta e^{-(t-s)A} \nabla \cdot (w \nabla u)\|_{L^q} ds \\ &\leq C_4 \int_0^t (t-s)^{\theta-\frac{1}{2}-\epsilon} e^{-(\mu+1)(t-s)} \|w \nabla u\|_{L^q} ds \\ &\leq C_0 C_3 C(\tau) \int_0^t (t-s)^{\theta-\frac{1}{2}-\epsilon} e^{-(\mu+1)(t-s)} ds \\ &\leq C_4 \int_0^\infty \rho^{-\theta-\frac{1}{2}-\epsilon} e^{-(\mu+1)\rho} d\rho \\ &\leq C_5 \Gamma\left(\frac{1}{2} - \theta - \epsilon\right), \end{aligned} \quad (3.48)$$

where $\Gamma(\frac{1}{2} - \theta - \epsilon)$ is a Gamma function which is positive since $\frac{1}{2} - \theta - \epsilon > 0$ and $\mu, C_5 > 0$.

Next, we obtain the bound for I_3 . As in the estimate of I_2 , we set $m = 0$, $q = n + 1$, $p = \infty$ for (2.8) and we use (3.33) and (3.1), one has

$$\begin{aligned} \|I_3\|_{L^\infty} &\leq C_3 \chi_1 \int_0^t \|(A+1)^\theta e^{-(t-s)(A+1)} \nabla \cdot (w \nabla v)\|_{L^q} ds \\ &\leq C_3 \chi_2 \int_0^t e^{-(t-s)} \|(A+1)^\theta e^{-(t-s)A} \nabla \cdot (w \nabla v)\|_{L^q} ds \\ &\leq C_6 \int_0^t (t-s)^{\theta-\frac{1}{2}-\epsilon} e^{-(\mu+1)(t-s)} \|w \nabla v\|_{L^q} ds \\ &\leq C_0 C_6 C(\tau) \int_0^t (t-s)^{\theta-\frac{1}{2}-\epsilon} e^{-(\mu+1)(t-s)} ds \\ &\leq C_7 \int_0^\infty \rho^{-\theta-\frac{1}{2}-\epsilon} e^{-(\mu+1)\rho} d\rho \\ &\leq C_8 \Gamma\left(\frac{1}{2} - \theta - \epsilon\right), \end{aligned} \quad (3.49)$$

where $\Gamma(\frac{1}{2} - \theta - \epsilon)$ is a Gamma function which is positive since $\frac{1}{2} - \theta - \epsilon > 0$ and $\mu, C_8 > 0$. Finally, we obtain the bound for I_4 . To this end, we use (2.6) and (2.7) and let $m = 1$, $p = (n, \infty]$ and $q = n + 1$. Hence, we can choose $\theta \in (\frac{1}{2}(1 - \frac{n}{p} + \frac{n}{q}), 1)$. Then one has

$$\begin{aligned} \|I_4\|_{W^{1,p}} &\leq C_1 \|(A+1)^\theta I_4\|_{L^q} \\ &\leq C_1 C_2 \int_0^t (t-s)^{-\theta} e^{-\nu t} \|f(u, v, w)\|_{L^q} ds. \end{aligned} \quad (3.50)$$

Using the fact that $0 < u \leq K_0, 0 < v \leq K_1$ and (3.1), we can get

$$\begin{aligned} \left\| \frac{a_5 u w + a_6 v w}{1 + a_2 u + a_3 v} + (1 - \mu) w \right\|_{L^q} &\leq \tilde{K} \|w\|_{L^q} + (1 + \mu) \|w\|_{L^q} \\ &\leq [\tilde{K} + (1 + \mu)] \|w\|_{L^q} \\ &\leq [\tilde{K} + (1 + \mu)] C(\tau) \end{aligned} \quad (3.51)$$

for all $t \in (\tau, T_{max})$. Hence, we have

$$\begin{aligned} \|I_4\|_{W^{1,p}} &\leq C_1 C_2 [\tilde{K} + (1 + \mu)] C(\tau) \int_0^t (t-s)^{-\theta} e^{-\nu t} ds \\ &\leq C_1 C_2 [\tilde{K} + (1 + \mu)] C(\tau) \int_0^\infty \sigma^{-\theta} e^{-\nu t} d\sigma \\ &\leq C_1 C_2 [\tilde{K} + (1 + \mu)] C(\tau) \nu^\theta \Gamma(1 - \theta) \end{aligned} \quad (3.52)$$

for all $t \in (\tau, T_{max})$ and where $\Gamma(1 - \theta)$ is a Gamma function and it is positive since $1 - \theta > 0$ and $\nu > 0$. Since $p > n$, Sobolev embedding theorem yields that

$$\|I_4\|_{L^\infty} \leq C_9 \text{ for all } t \in (\tau, T_{max}). \quad (3.53)$$

Substituting the estimates (3.47), (3.48), (3.49), (3.53) into (3.46) which yields (3.32). Hence, this completes the proof. \square

Proof of Theorem 1.1. From Lemma 2.5, we obtain $\|(u(\cdot, t), v(\cdot, t))\|_{L^\infty(\Omega)} \leq C$. Further, we also obtain the bound for $\|w(\cdot, t)\|_{L^\infty}$ from Lemma 3.2. By noticing these results now, we can conclude that

$$\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{L^\infty} + \|w(\cdot, t)\|_{L^\infty} \leq c \text{ for all } t \in (0, T_{max}), \quad (3.54)$$

where c is a positive constant. From the criterion (2.9), we obtain that $T_{max} = \infty$ and hence $\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{L^\infty} + \|w(\cdot, t)\|_{L^\infty} \leq c$ for all $t \in (0, \infty)$. The proof of Theorem 1.1 is complete.

4. Global stability of solutions

In this section, we shall prove the global stability of solutions of (1.3) by constructing some suitable Lyapunov functionals, and then we use the LaSalle's principle.

4.1. Global stability of Prey only state

Lemma 4.1. Let (u, v, w) be the solution of (1.3) and let $\Gamma_1 = \frac{(a_5(1+a_3-a_2a_6))}{a_1(1+a_2+a_3)}, \Gamma_2 = \frac{(a_6(1+a_2)-a_3a_5)}{a_4(1+a_2+a_3)}$. Then, if $\mu \geq \frac{a_5+a_6}{1+a_2+a_3}$ and $a_4 < \frac{a_1 r(a_5+a_3a_5-a_2a_6)}{a_3a_5-a_6-a_2a_6}$, it holds that

$$\lim_{t \rightarrow \infty} (\|u(\cdot, t) - 1\|_{L^\infty} + \|v(\cdot, t) - 1\|_{L^\infty} + \|w(\cdot, t)\|_{L^\infty}) = 0. \quad (4.1)$$

Proof. Let us define the energy functional

$$\mathcal{E}(t) := \Gamma_1 \int_{\Omega} (u - 1 - \ln u) + \Gamma_2 \int_{\Omega} (v - 1 - \ln v) + \int_{\Omega} w. \quad (4.2)$$

First we need to prove that $\mathcal{E}(t) \geq 0$ and $\mathcal{E}(t) = 0$ iff $(u, v, w) = (1, 1, 0)$. To this end, let $\psi(x) = x - g_* \ln x$ and by Taylor's formula, one has

$$\begin{aligned} g - g_* - g_* \ln \frac{g}{g_*} &= \psi(g) - \psi(g_*) = \psi'(g_*)(g - g_*) + \frac{1}{2} \psi''(\delta)(g - g_*)^2 \\ &= \frac{g_*}{2\delta^2}(g - g_*), \end{aligned}$$

where we choose δ in between g_* and g . Now putting $g = u$ and $g_* = 1$ in the last equation, we have

$$u - 1 - \ln u = \frac{1}{2\delta^2}(u - 1)^2 \geq 0.$$

Similarly, we can show that

$$v - 1 - \ln v = \frac{1}{2\delta^2}(v - 1)^2 \geq 0.$$

Therefore, we get $\mathcal{E}(u, v, w) = \mathcal{E}(1, 1, 0) = 0$ and $\mathcal{E}(u, v, w) \geq 0$ for $(u, v, w) \neq (1, 1, 0)$. Differentiating (4.2) with respect to t , and then substituting equations from (1.3), we have

$$\begin{aligned} \frac{d\mathcal{E}(t)}{dt} &= \Gamma_1 \int_{\Omega} \frac{u-1}{u} u_t + \Gamma_2 \int_{\Omega} \frac{v-1}{v} v_t + \int_{\Omega} w_t \\ &= -\Gamma_1 d \int_{\Omega} \frac{|\nabla u|^2}{u^2} + \Gamma_1 \int_{\Omega} \left[1 - u - \frac{a_1 w}{1 + a_2 u + a_3 v} \right] (u-1) + \Gamma_2 \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} \\ &\quad + \Gamma_2 \int_{\Omega} \left[r(1-v) - \frac{a_4 w}{1 + a_2 u + a_3 v} \right] (v-1) + \int_{\Omega} \left[\frac{a_5 u + a_6 v}{1 + a_2 u + a_3 v} - \mu \right] w \\ &\leq -\Gamma_1 d \int_{\Omega} \frac{|\nabla u|^2}{u^2} + \Gamma_1 \int_{\Omega} \left[1 - u - \frac{a_1 w}{1 + a_2 u + a_3 v} \right] (u-1) - \Gamma_2 \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} \\ &\quad + \Gamma_2 \int_{\Omega} \left[r(1-v) - \frac{a_4 w}{1 + a_2 u + a_3 v} \right] (v-1) + \int_{\Omega} \left[\frac{a_5 u + a_6 v}{1 + a_2 u + a_3 v} - \frac{a_5 + a_6}{1 + a_2 + a_3} \right] w \\ &\leq -\Gamma_1 d \int_{\Omega} \frac{|\nabla u|^2}{u^2} + \Gamma_1 \int_{\Omega} \left[1 - u - \frac{a_1 w}{1 + a_2 u + a_3 v} \right] (u-1) - \Gamma_2 \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} \\ &\quad + \Gamma_2 \int_{\Omega} \left[r(1-v) - \frac{a_4 w}{1 + a_2 u + a_3 v} \right] (v-1) \\ &\quad + \int_{\Omega} \left[\frac{a_5(u-1) + a_3 a_5(u-v)}{(1 + a_2 u + a_3 v)(1 + a_2 + a_3)} + \frac{a_6(v-1) + a_2 a_6(v-u)}{(1 + a_2 + a_3)(1 + a_2 u + a_3 v)} \right] w. \end{aligned} \quad (4.3)$$

By using the assumptions of Γ_1 and Γ_2 in (4.3), one has

$$\frac{d\mathcal{E}(t)}{dt} \leq -\Gamma_1 d \int_{\Omega} \frac{|\nabla u|^2}{u^2} - \Gamma_2 \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} - \Gamma_1 \int_{\Omega} (u-1)^2 - \Gamma_2 r \int_{\Omega} (v-1)^2, \quad (4.4)$$

which yields

$$\frac{d\mathcal{E}(t)}{dt} \leq 0, \quad (4.5)$$

for all (u, v, w) and also the equality holds when $(u, v, w) = (1, 1, 0)$. At last, the LaSalle's invariance principle (cf. [43], Theorem 3) yields that the solutions (u, v, w) converge to the constant steady state $(1, 1, 0)$ as time $t \rightarrow \infty$. \square

4.2. Global stability of coexistence state

Lemma 4.2. Let $\Gamma_1 = \frac{a_5 + (a_3 a_5 - a_2 a_6) v_*}{a_1 (1 + a_2 u_* + a_3 v_*)}$ and $\Gamma_2 = \frac{a_6 + (a_2 a_6 - a_3 a_5) u_*}{a_4 (1 + a_2 u_* + a_3 v_*)}$ be positive constants. Let (u, v, w) be the solution of (1.3). If $\mu < \frac{a_5 + a_6}{1 + a_2 + a_3}$, $a_4 > \frac{a_1 r (a_5 + a_3 a_5 - a_2 a_6)}{a_3 a_5 - a_6 - a_2 a_6}$ and

$$\frac{\Gamma_1 (2a_2 + a_3)}{2} + \frac{\Gamma_2 a_2 r}{2} < \Gamma_1 + \frac{\Gamma_1 (2a_2 + a_3) u_*}{2} + \frac{\Gamma_2 a_2 r v_*}{2}, \quad (4.6)$$

$$\frac{\Gamma_2 r (2a_3 + a_2)}{2} + \frac{\Gamma_1 a_3}{2} < r \Gamma_2 + \frac{\Gamma_2 r (2a_3 + a_2) v_*}{2} + \frac{\Gamma_1 a_3 u_*}{2}, \quad (4.7)$$

$$4d\Gamma_1 \Gamma_2 \eta u_* v_* > \Gamma_1 \chi_2^2 u_* w_* K_1^2 + \chi_1^2 \eta \Gamma_2 K_0^2 v_* w_*, \quad (4.8)$$

then it holds that

$$\lim_{t \rightarrow \infty} (\|u(\cdot, t) - u_*\|_{L^\infty} + \|v(\cdot, t) - v_*\|_{L^\infty} + \|w(\cdot, t) - w_*\|_{L^\infty}) = 0. \quad (4.9)$$

Proof. The coexistence steady state (u_*, v_*, w_*) of (1.3) satisfies equations

$$\begin{aligned} (1 - u_*) - \frac{a_1 w_*}{1 + a_2 u_* + a_3 v_*} &= 0, \\ r(1 - v_*) - \frac{a_4 w_*}{1 + a_2 u_* + a_3 v_*} &= 0, \\ -\mu + \frac{a_5 u_*}{1 + a_2 u_* + a_3 v_*} + \frac{a_6 v_*}{1 + a_2 u_* + a_3 v_*} &= 0. \end{aligned}$$

Let us define the Lyapunov functional $\mathcal{E}(u, v, w)$ as

$$\mathcal{E}(u, v, w) := \Gamma_1 \mathcal{F}_1(t) + \Gamma_2 \mathcal{F}_2(t) + \mathcal{F}_3(t), \quad (4.10)$$

where

$$\mathcal{F}_1(t) = \int_{\Omega} u - u_* - u_* \log\left(\frac{u}{u_*}\right), \mathcal{F}_2(t) = \int_{\Omega} v - v_* - v_* \log\left(\frac{v}{v_*}\right), \mathcal{F}_3(t) = \int_{\Omega} w - w_* - w_* \log\left(\frac{w}{w_*}\right). \quad (4.11)$$

Next, we take the derivative of $\mathcal{E}(t)$ with respect to t along the trajectory of the system (1.3) and we obtain

$$\frac{d\mathcal{E}(t)}{dt} = \Gamma_1 \frac{d\mathcal{F}_1(t)}{dt} + \Gamma_2 \frac{d\mathcal{F}_2(t)}{dt} + \frac{d\mathcal{F}_3(t)}{dt} := \Gamma_1 I_1 + \Gamma_2 I_2 + I_3. \quad (4.12)$$

Using the definition of $\mathcal{F}_1(t)$, the first equation of (1.3) and the fact that $(1 - u_*) - \frac{a_1 w_*}{1 + a_2 u_* + a_3 v_*} = 0$, we estimate the term I_1 as follows:

$$\begin{aligned}
I_1 &= \int_{\Omega} \frac{u - u_*}{u} \left(d\Delta u + u(1 - u) - \frac{a_1 u w}{1 + a_2 u + a_3 v} \right) \\
&= -u_* \int_{\Omega} \frac{|\nabla u|^2}{u^2} + \int_{\Omega} (u - u_*) \left(1 - u - \frac{a_1 w}{1 + a_2 u + a_3 v} - 1 + u_* + \frac{a_1 w_*}{1 + a_2 u_* + a_3 v_*} \right) \\
&= -u_* d \int_{\Omega} \frac{|\nabla u|^2}{u^2} + \int_{\Omega} (u - u_*) \left(-(u - u_*) - \frac{a_1 w(1 + a_2 u_* + a_3 v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \right. \\
&\quad \left. + \frac{a_1 w_*(1 + a_2 u + a_3 v)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \right) \\
&= -u_* d \int_{\Omega} \frac{|\nabla u|^2}{u^2} - \int_{\Omega} (u - u_*)^2 + \int_{\Omega} \frac{[-a_1 w(1 + a_2 u_* + a_3 v_*)](u - u_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \\
&\quad + \frac{[a_1 w_*(1 + a_2 u + a_3 v)](u - u_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)}. \tag{4.13}
\end{aligned}$$

Now, let us simplify the numerator in the integrand of the third integral on the R.H.S. of (4.13) as follows:

$$\begin{aligned}
&[-a_1 w(1 + a_2 u_* + a_3 v_*)](u - u_*) + [a_1 w_*(1 + a_2 u + a_3 v)](u - u_*) \\
&= [-a_1(w - w_*) + a_1 a_2(uw_* - wu_*) + a_1 a_3(w_* v - wv_*)](u - u_*) \\
&= [-a_1(w - w_*) + a_1 a_2(uw_* + u_* w_* - u_* w_* - wu_*) \\
&\quad + a_1 a_3(w_* v + v_* w_* - v_* w_* - wv_*)](u - u_*) \\
&= -a_1(w - w_*)(u - u_*) + a_1 a_2(w_*(u - u_*) - u_*(w - w_*))(u - u_*) \\
&\quad + a_1 a_3(w_*(v - v_*) - v_*(w - w_*))(u - u_*). \tag{4.14}
\end{aligned}$$

Substituting (4.14) into the last integral on the R.H.S of (4.13), we obtain

$$\int_{\Omega} \frac{[-a_1 w(1 + a_2 u_* + a_3 v_*) + a_1 w_*(1 + a_2 u + a_3 v)](u - u_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} = \int_{\Omega} \frac{a_1 a_2 w_*(u - u_*)^2}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \tag{4.15}$$

$$- \int_{\Omega} \frac{a_1 [1 + a_2 u_* + a_3 v_*](u - u_*)(w - w_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} + \int_{\Omega} \frac{a_1 a_3 w_*(u - u_*)(v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)}. \tag{4.16}$$

Again, inserting (4.16) into (4.13), we end up with

$$\begin{aligned}
I_1 &= -u_* d \int_{\Omega} \frac{|\nabla u|^2}{u^2} - \int_{\Omega} (u - u_*)^2 + \int_{\Omega} \frac{a_1 a_2 w_*(u - u_*)^2}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \\
&\quad - \int_{\Omega} \frac{a_1 [1 + a_2 u_* + a_3 v_*](u - u_*)(w - w_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} + \int_{\Omega} \frac{a_1 a_3 w_*(u - u_*)(v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \\
&= -u_* d \int_{\Omega} \frac{|\nabla u|^2}{u^2} + \int_{\Omega} \left(\frac{a_1 a_2 w_*}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} - 1 \right) (u - u_*)^2 \\
&\quad - \int_{\Omega} \frac{a_1 [1 + a_2 u_* + a_3 v_*](u - u_*)(w - w_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} + \int_{\Omega} \frac{a_1 a_3 w_*(u - u_*)(v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)}. \tag{4.17}
\end{aligned}$$

Similarly, we estimate the term I_2 . Using the definition of $\mathcal{F}_2(t)$, the second equation of (1.3) and the fact that $r(1 - v_*) - \frac{a_4 w_*}{1 + a_2 u_* + a_3 v_*} = 0$, we estimate I_2 as follows:

$$\begin{aligned}
 I_2 &= \int_{\Omega} \frac{v - v_*}{v} \left(\eta d \Delta v + rv(1 - v) - \frac{a_4 v w}{1 + a_2 u + a_3 v} \right) \\
 &= -v_* \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} + \int_{\Omega} (v - v_*) \left(-rv - \frac{a_4 w}{1 + a_2 u + a_3 v} + rv_* + \frac{a_4 w_*}{1 + a_2 u_* + a_3 v_*} \right) \\
 &= -v_* \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} + \int_{\Omega} (v - v_*) \left(-r(v - v_*) - \frac{a_4 w(1 + a_2 u_* + a_3 v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \right. \\
 &\quad \left. + \frac{a_4 w_*(1 + a_2 u + a_3 v)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \right) \\
 &= -v_* \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} - r \int_{\Omega} (v - v_*)^2 + \int_{\Omega} \frac{[-a_4 w(1 + a_2 u_* + a_3 v_*)](v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \\
 &\quad + \frac{[a_4 w_*(1 + a_2 u + a_3 v)](v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)}. \tag{4.18}
 \end{aligned}$$

Now, let us simplify the numerator in the integrand of the third integral on the R.H.S. of (4.18) as follows:

$$\begin{aligned}
 [-a_4 w(1 + a_2 u_* + a_3 v_*) + a_4 w_*(1 + a_2 u + a_3 v)](u - u_*) &= a_3 a_4 w_*(v - v_*)^2 \\
 - a_4 [1 + a_2 u_* + a_3 v_*](v - v_*)(w - w_*) + a_2 a_4 w_*(u - u_*)(v - v_*) &. \tag{4.19}
 \end{aligned}$$

Substituting (4.19) into (4.18) and simplifying, we obtain

$$\begin{aligned}
 I_2 &= -v_* \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} - r \int_{\Omega} (v - v_*)^2 + \int_{\Omega} \frac{a_2 a_3 w_*(v - v_*)^2}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \\
 &\quad - \int_{\Omega} \frac{a_4 [1 + a_2 u_* + a_3 v_*](v - v_*)(w - w_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} + \int_{\Omega} \frac{a_2 a_4 w_*(u - u_*)(v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} \\
 &= -v_* \eta d \int_{\Omega} \frac{|\nabla v|^2}{v^2} + \int_{\Omega} \left(\frac{a_2 a_3 w_*}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} - r \right) (v - v_*)^2 \\
 &\quad - \int_{\Omega} \frac{a_4 [1 + a_2 u_* + a_3 v_*](v - v_*)(w - w_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)} + \int_{\Omega} \frac{a_2 a_4 w_*(u - u_*)(v - v_*)}{(1 + a_2 u + a_3 v)(1 + a_2 u_* + a_3 v_*)}. \tag{4.20}
 \end{aligned}$$

Finally, we estimate the term I_3 . Using the definition of $\mathcal{F}_3(t)$, the third equation of (1.3) and the fact that $-\mu + \frac{a_5 u_*}{1 + a_2 u_* + a_3 v_*} + \frac{a_6 v_*}{1 + a_2 u_* + a_3 v_*} = 0$, we find

$$\begin{aligned}
 I_3 &= \int_{\Omega} \frac{w - w_*}{w} (\Delta w - \chi_1 \nabla \cdot (w \nabla u) - \chi_2 \nabla \cdot (w \nabla v)) \\
 &\quad + \int_{\Omega} (w - w_*) \left(-\mu + \frac{a_5 u + a_6 v}{1 + a_2 u + a_3 v} \right) \\
 &= -w_* \int_{\Omega} \frac{|\nabla w|^2}{w^2} + \chi_1 w_* \int_{\Omega} \frac{\nabla u \nabla w}{w} + \chi_2 w_* \int_{\Omega} \frac{\nabla v \nabla w}{w} \\
 &\quad + \int_{\Omega} (w - w_*) \left(\frac{a_5 u + a_6 v}{1 + a_2 u + a_3 v} - \frac{a_5 u_* + a_6 v_*}{1 + a_2 u_* + a_3 v_*} \right). \tag{4.21}
 \end{aligned}$$

Further, we simplify the numerator in the integrand of the third integral on the R.H.S. of (4.21), one has that

$$\begin{aligned} & (a_5u + a_6v)(1 + a_2u_* + a_3v_*) - (a_5u_* + a_6v_*)(1 + a_2u + a_3v) \\ &= a_5(u - u_*) + a_6(v - v_*) + a_3a_5[v_*(u - u_*) - u_*(v - v_*)] + a_2a_6[u_*(v - v_*) - v_*(u - u_*)] \\ &= [a_5 + (a_3a_5 - a_2a_6)v_*](u - u_*) + [a_6 + (a_2a_6 - a_3a_5)u_*](v - v_*). \end{aligned}$$

Now, we rewrite the fourth term in the R.H.S of (4.21) using the above estimate, we obtain

$$\begin{aligned} \int_{\Omega} (w - w_*) \left(\frac{a_5u + a_6v}{1 + a_2u + a_3v} - \frac{a_5u_* + a_6v_*}{1 + a_2u_* + a_3v_*} \right) &= \int_{\Omega} \frac{[a_5 + (a_3a_5 - a_2a_6)v_*](u - u_*)(w - w_*)}{(1 + a_2u + a_3v)(1 + a_2u_* + a_3v_*)} \\ &+ \int_{\Omega} \frac{[a_6 + (a_2a_6 - a_3a_5)u_*](v - v_*)(w - w_*)}{(1 + a_2u + a_3v)(1 + a_2u_* + a_3v_*)}. \end{aligned} \quad (4.22)$$

Substituting (4.22) into (4.21), we have

$$\begin{aligned} I_3 &= -w_* \int_{\Omega} \frac{|\nabla w|^2}{w^2} + \chi_1 w_* \int_{\Omega} \frac{\nabla u \cdot \nabla w}{w} + \chi_2 w_* \int_{\Omega} \frac{\nabla v \cdot \nabla w}{w} + \int_{\Omega} \frac{[a_5 + (a_3a_5 - a_2a_6)v_*](u - u_*)(w - w_*)}{(1 + a_2u + a_3v)(1 + a_2u_* + a_3v_*)} \\ &+ \int_{\Omega} \frac{[a_6 + (a_2a_6 - a_3a_5)u_*](v - v_*)(w - w_*)}{(1 + a_2u + a_3v)(1 + a_2u_* + a_3v_*)}. \end{aligned} \quad (4.23)$$

Furthermore, inserting (4.17), (4.20) and (4.23) into (4.12) and let $p(u, v) = \frac{1}{(1+a_2u+a_3v)(1+a_2u_*+a_3v_*)}$, we arrive at

$$\begin{aligned} \frac{d\mathcal{E}(t)}{dt} &= -u_* d\Gamma_1 \int_{\Omega} \frac{|\nabla u|^2}{u^2} - v_* \eta d\Gamma_2 \int_{\Omega} \frac{|\nabla v|^2}{v^2} - w_* \int_{\Omega} \frac{|\nabla w|^2}{w^2} + \chi_1 w_* \int_{\Omega} \frac{\nabla u \cdot \nabla w}{w} + \chi_2 w_* \int_{\Omega} \frac{\nabla v \cdot \nabla w}{w} \\ &+ \Gamma_1 \int_{\Omega} \left(\frac{a_1 a_2 w_*}{p(u, v)} - 1 \right) (u - u_*)^2 + w_* (\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4) \int_{\Omega} \frac{(u - u_*)(v - v_*) w_*}{p(u, v)} \\ &+ \Gamma_2 \int_{\Omega} \left(\frac{a_4 a_3 w_*}{p(u, v)} - r \right) (v - v_*)^2. \end{aligned} \quad (4.24)$$

Applying the Cauchy's inequality, we get

$$\int_{\Omega} \frac{(u - u_*)(v - v_*)}{p(u, v)} \leq \frac{1}{2} \int_{\Omega} \frac{(u - u_*)^2}{p(u, v)} + \frac{1}{2} \int_{\Omega} \frac{(v - v_*)^2}{p(u, v)}. \quad (4.25)$$

Inserting the last inequality (4.25) into (4.24), and letting $Z = \left(\frac{|\nabla u|}{u}, \frac{|\nabla v|}{v}, \frac{|\nabla w|}{w} \right)$, we obtain

$$\begin{aligned} \frac{d\mathcal{E}(t)}{dt} &\leq \underbrace{-u_* d\Gamma_1 \int_{\Omega} \frac{|\nabla u|^2}{u^2} - v_* \eta d\Gamma_2 \int_{\Omega} \frac{|\nabla v|^2}{v^2} - w_* \int_{\Omega} \frac{|\nabla w|^2}{w^2} + \chi_1 w_* \int_{\Omega} \frac{\nabla u \cdot \nabla w}{w} + \chi_2 w_* \int_{\Omega} \frac{\nabla v \cdot \nabla w}{w}}_{I_4} \\ &+ \Gamma_1 \int_{\Omega} \left(\frac{a_1 a_2 w_*}{p(u, v)} - 1 \right) (u - u_*)^2 + \frac{w_* (\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2} \int_{\Omega} \frac{(u - u_*)^2}{p(u, v)} \\ &+ \frac{w_* (\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2} \int_{\Omega} \frac{(v - v_*)^2}{p(u, v)} + \Gamma_2 \int_{\Omega} \left(\frac{a_3 a_4 w_*}{p(u, v)} - r \right) (v - v_*)^2 \\ &\leq I_4 + \int_{\Omega} \left(\frac{2\Gamma_1 a_1 a_2 w_* + w_* (\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2p(u, v)} - \Gamma_1 \right) (u - u_*)^2 \end{aligned}$$

$$+ \int_{\Omega} \left(\frac{2\Gamma_2 a_3 a_4 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2p(u, v)} - r\Gamma_2 \right) (v - v_*)^2, \quad (4.26)$$

where

$$I_4 = - \int_{\Omega} Z^T B Z,$$

and the symmetric matrix is denoted by

$$B = \begin{bmatrix} d\Gamma_1 u_* & 0 & -\frac{\chi_1 w_* u}{2} \\ 0 & \eta d\Gamma_2 v_* & -\frac{\chi_2 w_* v}{2} \\ -\frac{\chi_1 w_* u}{2} & -\frac{\chi_2 w_* v}{2} & w_* \end{bmatrix}. \quad (4.27)$$

The above matrix B is positive definite if (4.8) holds. Therefore, we check that

$$\begin{vmatrix} dd\Gamma_1 u_* & 0 \\ 0 & \eta d\Gamma_2 v_* \end{vmatrix} = d^2 \eta^2 \Gamma^2 u_* v_* > 0 \quad (4.28)$$

and

$$|B| = \frac{dw_*}{4} [4d\Gamma_1 \Gamma_2 \eta u_* v_* - \Gamma_1 \chi_2^2 u_* w_* v_*^2 - \chi_1^2 \eta \Gamma_2 u^2 v_* w_*] > 0. \quad (4.29)$$

Hence, there exists a positive constant α , such that

$$\begin{aligned} \frac{d\mathcal{E}(t)}{dt} &\leq -\alpha \int_{\Omega} \left(\frac{|\nabla u|^2}{u^2} + \frac{|\nabla v|^2}{v^2} + \frac{|\nabla w|^2}{w^2} \right) + \int_{\Omega} \left(\frac{2\Gamma_1 a_1 a_2 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2p(u, v)} - \Gamma_1 \right) (u - u_*)^2 \\ &\quad + \int_{\Omega} \left(\frac{2\Gamma_2 a_2 a_3 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2p(u, v)} - r\Gamma_2 \right) (v - v_*)^2. \end{aligned} \quad (4.30)$$

Noting the facts that $1 - u_* - \frac{a_1 w_*}{1 + a_2 u_* + a_3 v_*} = 0$ and $r(1 - v_*) - \frac{a_4 w_*}{1 + a_2 u_* + a_3 v_*} = 0$, one has that

$$\begin{aligned} -\Gamma_1 + \frac{2\Gamma_1 a_1 a_2 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2(1 + a_2 u_* + a_3 v_*)} &\leq -\Gamma_1 + \frac{2\Gamma_1 a_1 a_2 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2(1 + a_2 u_* + a_3 v_*)} \\ &= -\Gamma_1 + \Gamma_1 a_2 (1 - u_*) + \frac{\Gamma_2 a_2 r (1 - v_*)}{2} + \frac{\Gamma_1 a_3 (1 - u_*)}{2} \\ &\leq 0, \end{aligned}$$

and

$$\begin{aligned} -r\Gamma_2 + \frac{2\Gamma_2 a_2 a_3 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2p(u, v)} &\leq -r\Gamma_2 + \frac{2\Gamma_2 a_2 a_3 w_* + w_*(\Gamma_1 a_1 a_3 + \Gamma_2 a_2 a_4)}{2(1 + a_2 u_* + a_3 v_*)} \\ &= -r\Gamma_2 + \Gamma_2 a_3 r (1 - v_*) + \frac{\Gamma_1 a_3 (1 - u_*)}{2} + \frac{\Gamma_2 a_2 r (1 - v_*)}{2} \\ &\leq 0, \end{aligned}$$

where we used the assumptions (4.6) and (4.7). Hence, we can conclude that

$$\frac{d}{dt} \mathcal{E}(t) \leq -c \int_{\Omega} \left(\frac{|\nabla u|^2}{u^2} + \frac{|\nabla v|^2}{v^2} + \frac{|\nabla w|^2}{w^2} \right), \quad (4.31)$$

which yields that $\frac{d}{dt}\mathcal{E}(t) \leq 0$ for all u, v, w and the equality holds if $\nabla u = \nabla v = \nabla w = 0$. Therefore by applying LaSalle's invariance principle (cf. [43], Theorem 3) we can say that the solutions of (1.3) converges to the coexistence steady state (u_*, v_*, w_*) as time $t \rightarrow \infty$. \square

Proof of Theorem 1.2. Theorem 1.2 is a consequence of Lemma 4.1 and Lemma 4.2.

Remark 4.1. We note that the condition (4.8) is a strong requirement, which implies that χ_1 and χ_2 have to be small enough.

Acknowledgments

The author is grateful to the referees for insightful comments which largely help improve the exposition of this paper. The author also thanks Prof. Zhi-An Wang from Hong Kong Polytechnic University for the fruitful discussions and valuable suggestions. This work was supported by the Postdoc Matching Fund Schemes of the Hong Kong Polytechnic University (Project ID P0036730 and a/c no. W18M).

Conflict of interest

The author declares that there are no conflicts of interest.

References

1. N. Sapoukhina, Y. Tyutyunov, R. Arditi, The role of prey taxis in biological control: A spatial theoretical model, *Am. Nat.*, **162** (2003), 61–76. <https://doi.org/10.1086/375297>
2. A. Mondal, A. K. Pal, P. Dolai, G.P. Samanta, A system of two competitive prey species in presence of predator under the influence of toxic substances, *Filomat*, **36** (2) (2022), 361–385. <https://doi.org/10.2298/FIL2202361M>
3. M. A. Ragusa, A. Razani, Existence of a periodic solution for a coupled system of differential equations, in *AIP Conference Proceedings*, AIP Publishing LLC, 2022, 370004. <https://doi.org/10.1063/5.0081381>
4. J. Tian, P. Liu, Global dynamics of a modified Leslie-Gower predator-prey model with Beddington-DeAngelis functional response and prey-taxis, *Elec. Res. Arch.*, **30** (2022), 929–942. <https://doi.org/10.3934/era.2022048>
5. P. Kareiva, G. T. Odell, Swarms of predators exhibit preytaxis if individual predators use area-restricted search, *Am. Nat.*, **130** (1987), 233–270.
6. J. R. Beddington, Mutual interference between parasites or predators and its effect on searching efficiency, *J. Anim. Ecol.*, **44** (1975), 331–340.
7. D. L. DeAngelis, R. A. Goldstein, R. V. O'Neill, A model for trophic interaction, *Ecology*, **56** (1975), 881–892. <https://doi.org/10.2307/1936298>

8. P. A. Abrams, L. R. Ginzburg, The nature of predation: prey dependent, ratio dependent or neither?, *Trends Ecol. Evol.*, **15** (2000), 337–341. [https://doi.org/10.1016/S0169-5347\(00\)01908-X](https://doi.org/10.1016/S0169-5347(00)01908-X)
9. B. Ainseba, M. Bendahmane, A. Noussair, A reaction-diffusion system modeling predator-prey with prey-taxis, *Nonlinear Anal. Real World Appl.*, **9** (2008), 2086–2105. <https://doi.org/10.1016/j.nonrwa.2007.06.017>
10. M. Bendahmane, Analysis of a reaction-diffusion system modeling predator-prey with prey-taxis, *Netw. Heterog. Media*, **3** (2008), 863–879. <https://doi.org/10.3934/nhm.2008.3.863>
11. Y. Cai, Q. Cao, Z. A. Wang, Asymptotic dynamics and spatial patterns of a ratio-dependent predator-prey system with prey-taxis, *Appl. Anal.*, **101** (2022), 81–99. <https://doi.org/10.1080/00036811.2020.1728259>
12. H. Y. Jin, Z. A. Wang, Global dynamics and spatio-temporal patterns of predator-prey systems with density-dependent motion, *Eur. J. Appl. Math.*, **32** (2021), 652–682. <https://doi.org/10.1017/S0956792520000248>
13. D. Li, Global stability in a multi-dimensional predator-prey system with prey-taxis, *Discrete Contin. Dyn. Syst. Ser.*, **41** (2021), 1681–1705. <https://doi.org/10.3934/dcds.2020337>
14. S. Li, R. Mu, Positive steady-state solutions for predator-prey systems with prey-taxis and Dirichlet conditions, *Nonlinear Anal. Real World Appl.*, **68** (2022), 103669. <https://doi.org/10.1016/j.nonrwa.2022.103669>
15. D. Luo, Global bifurcation for a reaction-diffusion predator-prey model with Holling-II functional response and prey-taxis, *Chaos Soliton. Fract.*, **147** (2021), 110975. <https://doi.org/10.1016/j.chaos.2021.110975>
16. X. L. Wang, W. D. Wang, G. H. Zhang, Global bifurcation of solutions for a predator-prey model with prey-taxis, *Math. Methods Appl. Sci.*, **3** (2014), 431–443. <https://doi.org/10.1002/mma.3079>
17. T. Xiang, Global dynamics for a diffusive predator-prey model with prey-taxis and classical Lotka-Volterra kinetics, *Nonlinear Anal. Real World Appl.*, **39** (2018) 278–299. <https://doi.org/10.1016/j.nonrwa.2017.07.001>
18. L. Zhang, S. Fu, Global bifurcation for a Holling Tanner predator-prey model, *Nonlinear Anal. Real World Appl.*, **47** (2019), 460–472. <https://doi.org/10.1016/j.nonrwa.2018.12.002>
19. X. He, S. Zheng, Global boundedness of solutions in a reaction-diffusion system of predator-prey model with prey-taxis, *Appl. Math. Lett.*, **49** (2015), 73–77. <https://doi.org/10.1016/j.aml.2015.04.017>
20. W. Choi, I. Ahn, Predator invasion in predator-prey model with prey-taxis in spatially heterogeneous environment, *Nonlinear Anal. Real World Appl.*, **65** (2022), 103495. <https://doi.org/10.1016/j.nonrwa.2021.103495>
21. H. Y. Jin, Z. A. Wang, Global stability of prey-taxis systems, *J. Differ. Equation*, **262** (2017), 1257–1290. <https://doi.org/10.1016/j.jde.2016.10.010>

22. Y. Tao, Global existence of classical solutions to a predator-prey model with nonlinear prey-taxis, *Nonlinear Anal. Real World Appl.*, **11** (2010), 2056–2064. <https://doi.org/10.1016/j.nonrwa.2009.05.005>
23. S. N. Wu, J. P. Shi, B. Y. Wu, Global existence of solutions and uniform persistence of a diffusive predator-prey model with prey-taxis, *J. Differ. Equation*, **260** (2016), 5847–5874. <https://doi.org/10.1016/j.jde.2015.12.024>
24. J. Wang, M. X. Wang, Boundedness and global stability of the two-predator and one-prey models with nonlinear prey-taxis, *Z. Angew. Math. Phys.* **69** (2018), 63. <https://doi.org/10.1007/s00033-018-0960-7>
25. Z. Feng, M. Zhang, Boundedness and large time behavior of solutions to a prey-taxis system accounting in liquid surrounding, *Nonlinear Anal., Real World Appl.*, **57** (2021), 103197. <https://doi.org/10.1016/j.nonrwa.2020.103197>
26. E. C. Haskel, J. Bell, Pattern formation in a predator-mediated coexistence model with prey-taxis, *Dis. Cont. Dyn. Sys.*, **25** (2020), 2895–2921. <https://doi.org/10.3934/dcdsb.2020045>
27. E. C. Haskel, J. Bell, Bifurcation analysis for a one predator and two prey model with prey-taxis, *J. Bio. Sys.*, **29**, 495–524. <https://doi.org/10.1142/S0218339021400131>
28. X. Xu, Y. Wang, Y. Wang, Local bifurcation of a Ronsenzwing-MacArthur predator prey model with two prey-taxis, *Math. Bio. Eng.*, **16** (2019), 1786–1797. <https://doi.org/10.3934/mbe.2019086>
29. H. Y. Jin, Z. A. Wang, L. Y. Wu, Global dynamics of a three species spatial food chain model, *J. Differ. Equation*, **333** (2022), 144–183. <https://doi.org/10.1016/j.jde.2022.06.007>
30. X. D. Zhao, F. Y. Yang, W. T. Li, Traveling waves for a nonlocal dispersal predator-prey model with two preys and one predator, *Z. Angew. Math. Phys.*, **73** (2022), 124. <https://doi.org/10.1007/s00033-022-01753-5>
31. P. Amorim, R. Bürger, R. Ordoñez, L. M. Villada, Global existence in a food chain model consisting of two competitive preys, one predator and chemotaxis, *Nonlinear Anal. Real World Appl.*, **69** (2023), 103703. <https://doi.org/10.1016/j.nonrwa.2022.103703>
32. Y. Min, C. Song, Z. Wang, Boundedness and global stability of the predator -prey model with prey-taxis and competition, *Nonlinear Anal. Real World Appl.*, **66** (2022), 103521. <https://doi.org/10.1016/j.nonrwa.2022.103521>
33. X. Wang, R. Li, Y. Shi, Global generalized solutions to a three species predator-prey model with prey-taxis, *Discrete Contin. Dyn. Syst. Ser.-B*, **27** (2022), 7012–7042. <https://doi.org/10.3934/dcdsb.2022031>
34. H. Y. Jin, Z. A. Wang, Global stabilization of the full attraction-repulsion Keller-Segel system, *Discrete Conti. Dyn. Sys.*, **40** (2020), 3509–3527. <https://doi.org/10.3934/dcds.2020027>
35. P. Liu, J. P. Shi, Z. A. Wang, Pattern formation of the attraction-repulsion Keller-Segel system, *Discrete Contin. Dyn. Syst-Series B*, **18** (10) (2013), 2597–2625. <https://doi.org/10.3934/dcdsb.2013.18.2597>

36. M. Luca, A. Chavez-Ross, L. Edelstein, A. Mogilner, Chemotactic signalling, microglia, and Alzheimer's disease senile plaques: is there a connection?, *Bull. Math. Biol.*, **65** (2021), 110975. <https://doi.org/10.1016/j.chaos.2021.110975>
37. H. Amann, Dynamic theory of quasilinear parabolic equations. II. Reaction-diffusion systems, *Differ. Integral. Equation*, **3** (1990), 13–75.
38. H. Amann, Nonhomogeneous linear and quasilinear elliptic and parabolic boundary value problems, in *Function Spaces, Differential Operators and Nonlinear Analysis*, Springer Fachmedien, (1993), 13–75. https://doi.org/10.1007/978-3-663-11336-2_1
39. D. Horstmann, M. Winkler, Boundedness vs. blow-up in a chemotaxis system, *J. Differ. Equation*, **215** (2005), 52–107. <https://doi.org/10.1016/j.jde.2004.10.022>
40. C. Jin, Boundedness and global solvability to a chemotaxis-haptotaxis model with slow and fast diffusion, *Dis. Cont. Dyn. Sys. B*, **23** (2018), 1675–1688. <https://doi.org/10.3934/dcdsb.2018069>
41. H. Amann, Dynamic theory of quasilinear parabolic systems III. Global existence, *Math. Z.*, **202** (1989), 219–250. <https://doi.org/10.1007/BF01215256>
42. M. Winkler, Absence of collapse in parabolic chemotaxis system with signal-dependent sensitivity, *Math. Z.*, **283** (2010), 1664–1673. <https://doi.org/10.1002/mana.200810838>
43. J. LaSalle, Some extensions of Liapunov's second method, *IRE Trans. Circuit Theory*, **7** (1960), 520–527. <https://doi.org/10.1109/TCT.1960.1086720>



AIMS Press

©2023 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)