

Spectral correlation study of skin blood flow oscillation

T.X. Chi , Y. Liu and X.Y. Luo

Abstract Spontaneous rhythmic oscillations in microvessel diameter are known as vasomotion. Vasomotion is the intrinsic property of small arteries and arterioles and is dependent of heartbeat, respiration, or neuronal input. The skin microcirculation is an anastomotic network of vessels with many crucial functions in which the blood flow must be finely regulated and tuned in order to fulfill all the demands of the organism. The laser Doppler Flowmetry (LDF) can be used to measure dynamic changes in skin blood flow over a small area. Traditional Chinese medicine (TCM) considers that there exist acupoints around body which are connected by meridian, and stimulating the acupoint is a typical therapeutic technique in TCM. It was found that stimulating one acupoint could enhance the vasomotion in another acupoint significantly [1]. We argue that the vasomotions in these two acupoints could be correlated to each other. To verify our argument, we used the laser Doppler Flowmetry (LDF) to measure the skin blood flow at two acupoints, and carried out spectral and correlation analyses. It is found that the vasomotions related to myogenic activity are quite strong at acupoints and the correlation is discernable. The vasomotion related to myogenic activity at non-acupoint is much weaker and there is no significant correlation between the acupoint and non-acupoint.


1. INTRODUCTION

The spontaneous time-dependent contraction and relaxation of small arteries and arterioles are termed as vasomotion which may have considerable implications in the physiology and pathology of the arterial system. The spectral analysis of human skin blood flow oscillation has revealed five frequency components of vasomotion: the band 0.6 - 1.8 Hz is due to heart activity, the band 0.2 - 0.6 Hz is related to respiration, the band 0.06 - 0.2 Hz is related to myogenic activity, the band 0.02 - 0.06 Hz is associated to sympathetic activity and the band 0.007 - 0.02 Hz is more specifically related to endothelial activity [2]. It was found that the vasomotion propagates along arterioles, capillaries and venules in both upstream and downstream directions. Moreover, there is evidence that microcirculation may mirror the vascular function of other parts of the body and the microvascular abnormalities may initiate the pathogenesis sequence in some diseases [3].

CHI Tianxi
The HONG KONG Polytechnic University, email: 15900011r@connect.polyu.hk
LIU Yang
The HONG KONG Polytechnic University, email: mmyliu@polyu.edu.hk
LUO Xiaoyu
University of Glasgow, UK, email: Xiaoyu.Luo@glasgow.ac.uk

In TCM, the acupoints are interrelated to each other, and acupuncture at one acupoint may affect another acupoint. Several in vivo human studies have found that acupuncture at one acupoint may increase significantly the myogenic component of flowmotion at another acupoint [4]. To verify whether the vasomotions related to myogenic activity at acupoints are correlated to each other, we selected two acupoints along the same meridian (A & B). In addition, another non-acupoint (point C) is selected as reference.

Figure 1. Two selected acupoints (named Chize (A) and Kongzui (B)) and one non-acupoint (C) are measured simultaneously using LDF



2. MATERIALS AND METHODS

2.1 Measurement Procedure

Two LDF probes were positioned on two selected points. The ambient temperature is set at $23 \pm 1^\circ\text{C}$ in room and sampling rate is 20 Hz equivalent to 0.05s time interval. Points A-B and Points A-C were measured simultaneously. The measurement time was 20 mins.


2.2 Spectral Analysis

The mainly analytic tools analyzing the LDF signal are usually considered as the linear analysis which include Fast Fourier transform analysis (FFT) and Morlet wavelet analysis [5]. Periodic oscillations of skin blood flow can be quantified directly by FFT which could reveal the amplitude (energy) of the whole signal at given frequencies. For wavelet analysis, its feature allows us to determine both the dominant modes of blood flow oscillation and how those modes vary in time. For skin blood flow signals, Morlet wavelet analysis breaks down the steady fluctuating time series into its frequency elements and computes the power of signal components in predetermined frequency bands, allowing to measure the amplitude of different flowmotion waves in PU/Hz [6].

3. RESULTS AND DISCUSSION

3.1 FFT and Wavelet analysis

Figure 2(a)
FFT analysis
of acupoint A
& acupoint B






Figure 2(b) FFT analysis for one acupoint (A) & nonacupoint (C)





Figure 3(a) Morlet Wavelet analysis acupoint A & acupoint B

Figure 3(b) Morlet Wavelet analysis for acupoint A & nonacupoint C



From FFT analysis, the PSD of acupoint A and B at ~ 0.1 Hz are dominant on Figure 2(a) & (b), but not for the non-acupoint C. Thus It could be explained that the skin blood flow oscillation on these two acupoints (A & B) are influenced greatly by the myogenic factor (~ 0.1 Hz). In addition, all the PSDs of the lower frequency are higher than that of 1 Hz for these three points, indicating the myogenic component of vasomotion is significant. Comparing with the FFT analysis, the wavelet transform does not lose the time-series information of characteristic frequency, and it is obviously that the energy at different frequency varied with time. From wavelet analysis, both two Figures (3a & 3b) shown that the PSD of acupoint A & B at ~ 0.1 Hz exhibits periodic behaviour and follows a certain pattern, and the non-acupoint have no significance periodic pattern occurred at ~ 0.1 Hz. Therefore, the periodic behavior at ~ 0.1 Hz only exists in acupoints (A & B), and this periodic behavior should follow a certain transmitting direction.

3.2 Extracting Characteristic from wavelet & Correlation analysis

Since the PSD at 0.1 Hz in wavelet follows a periodic pattern, if the myogenic vasomotions at acupoints are correlated to each other, the

wavelet PSD variations with time should be correlated as well. We extracted the PSD signal at 0.1 Hz from wavelet analyses in Figures 3(a) & 3(b), and carried out cross-correlation analysis between these two extracted time series. The PSDs of both acupoints A and B are comparable (Fig. 4(a)), but the PSD of acupoint A is much higher than that of non-acupoint C (Fig. 4(b)). From point A to point B, there is no significant correlation identified; from point B to point A, the maximum correlation can be 0.6 with a phase lag 0.93π (Fig. 4a). Thus, it could be stated that the myogenic factor transmitted from point B to point A. For point A and point C, there is no significant correlation found, indicating the myogenic components in acupoint and non-acupoint are not correlated. Therefore, of the correlation of myogenic related vasomotion only existed between acupoints A and B.





Figure 4(a) PSD variation at 0.113 Hz from wavelet analysis and cross-correlation between acupoints A and B

Figure 4(b) extracted PSD from wavelet analysis and phase lag of PSD variation for acupoint A and non-acupoint C



4. CONCLUSION

The LDF measurement was carried out at acupoint and non-acupoint simultaneously, the spectrum and correlation analyses lead to following conclusion:

- (i) The myogenic component of vasomotion is dominant in term of PSD;
- (ii) The PSD of myogenic component is much higher in acupoint than that in non-acupoint;
- (iii) There exist correlation of myogenic components between acupoints, but there is no correlation between acupoint and non-acupoint.

5. ACKNOWLEDGEMENT

Supports given by HKRGC PolyU 5202/13E and PolyU G-YBG9 are gratefully acknowledged.

6. REFERENCE

- [1] Xiu RJ, Cheng J, Zhang J (1988) Effects of acupuncture on microvascular vasomotion. *Chinese Medical Journal* 68: 489-490.
- [2] Rossi, M. et al. 2006 Spectral analysis of laser Doppler skin blood flow oscillations in human essential arterial hypertension *Microvascular research*, **72**, pp. 34-41.
- [3] Secomb, T.W., Intaglietta, M. and Gross, J.F. 1989 Effects of vasomotion on microcirculatory mass transport: theoretical predictions. *Prog. Appl. Microcirc.* **15**, 49-61.
- [4] Mu X, Duan HQ, Chen W, Gao Y, Yang ZJ (2001) Physiological study on the relationship between the essence of acupuncture point and the microvasculum. *China Journal of Basic Medicine in Traditional Chinese Medicine* 7: 927-932.
- [5] Jan YK, Brienza DM, Geyer MJ, Karg P. 2008. Wavelet-based spectrum analysis of sacral skin blood flow response to alternating pressure. *Archives of Physical Medicine & Rehabilitation*. 89(1):137–45. [PubMed: 18164343]
- [6] Young J.D. And Cameron E.M. 1995. Dynamics of skin blood flow in human sepsis. *Intensive care medicine*, 21, 669-674.