

# Heavy Metal and Pb Isotopic Compositions of Aquatic Organisms in the Pearl River Estuary, South China

C.C.M. Ip<sup>1</sup>, X.D. Li<sup>1\*</sup>, G. Zhang<sup>2</sup>, C.S.C. Wong<sup>1</sup> and W.L. Zhang<sup>2</sup>

<sup>1</sup> *Department of Civil and Structural Engineering, The Hong Kong Polytechnic  
University, Hung Hom, Kowloon, Hong Kong*

<sup>2</sup> *State Key Laboratory of Organic Geochemistry, Guangzhou Institute of  
Geochemistry, Chinese Academy of Sciences, Guangzhou 510640, China*

**“Capsule”:** *Relative high concentrations of Cd were found in crab, shrimp and shellfish samples, while high concentration of Pb was found in fish, particularly from the anthropogenic inputs.*

## Abstract

The accumulation of trace metals in aquatic organisms may lead to serious health problems through the food chain. The present research project aims to study the accumulation and potential sources of trace metals in aquatic organisms of the Pearl River Estuary (PRE). Four groups of aquatic organisms, including fish, crab, shrimp, and shellfish, were collected in the PRE for trace metal and Pb isotopic analyses. The trace metal concentrations in the aquatic organism samples ranged from 0.01 to 2.10 mg/kg Cd, 0.02 to 4.33 mg/kg Co, 0.08 to 4.27 mg/kg Cr, 0.15 to 77.8 mg/kg Cu, 0.17 to 31.0 mg/kg Ni, 0.04 to 30.7 mg/kg Pb, and 8.78 to 86.3 mg/kg Zn (wet weight).

---

\* Corresponding author (X.D. Li), email: [cexdli@polyu.edu.hk](mailto:cexdli@polyu.edu.hk); fax: (852) 2334 6389; tel: (852) 2766 6041.

High concentrations of Cd were found in crab, shrimp and shellfish samples, while high concentration of Pb was found in fish. In comparison with the baseline reference values in other parts of the world, fish in the PRE had the highest elevated trace metals. The results of Pb isotopic compositions indicated that the bioaccumulation of Pb in fish come from a wide variety of food sources and/ or exposure pathways, particularly the anthropogenic inputs.

*Keyword:* Heavy metals, Pb isotope, aquatic organism (fish), Pearl River Estuary, China.

## **1. Introduction**

The mixed regime of the Pearl River Estuary (PRE), with fresh and oceanic water, provides a suitable habitat for a wide variety of aquatic organisms (Chen, 1995; Wang et al., 1995). However, rapid economic development in the Pearl River Delta (PRD) region in the last few decades has led to the excessive discharge of pollutants into the PRE (Li et al., 1997). Hence, great concern has arisen in recent years over environmental pollution in this coastal region. Elevated concentrations of Cu, Pb, and Zn in the sediments of the PRE have been found (Chen and Zhou, 1992; Zheng, 1992; Li et al., 2000a, 2000b and 2001; Liu et al., 2003). Trace metal contamination of the PRE may have a significant impact on aquatic organisms, disturbing the area's delicate ecological balance and potentially contaminating the marine food chain. Trace metal analysis of aquatic organisms from the PRE can provide important information on the degree of environmental contamination, and potential impact of

seafood consumption. In addition, the Pb isotopic compositions of aquatic organisms may further assist the identification of possible sources of contamination and biological pathways.

Aquatic organisms accumulate trace metals from various sources in the environment. The possible sources of trace metals include sediments (Labonne et al., 2001; Goodwin et al., 2003), soil erosion and runoff (Gelinas and Schmit, 1997), air depositions of dust and aerosol (Gelinas and Schmit, 1997; Labonne et al., 2001), discharges of wastewater (Labonne et al., 2001; Goodwin et al., 2003), and so forth (Bryan, 1979; Blackmore et al., 1998; Hoven et al., 1999; Goodwin et al., 2003). The accumulation of trace metals in aquatic organisms can pose a long-term burden on biogeochemical cycling in the ecosphere. Once trace metals enter the food chain, they may accumulate to dangerous levels and be harmful to human health (Manahan, 2000).

Stable Pb isotopic studies have been commonly applied to assess the sources of Pb in various ecosystems, including sediments (Farmer et al., 1996; Ritson et al., 1999; Liu et al., 2003), soils (Sugden et al., 1993; Semlali et al., 2001; Wong et al., 2002), suspended matters (Hinrichs et al., 2002), and atmospheric depositions (Bollhöfer and Rosman, 2000 and 2001; Wong et al., 2003). However, only a few studies have focused on stable Pb isotopes in biological samples to trace the anthropogenic origins of Pb (Kurkjian and Flegal, 2003). Previous studies have proven that the Pb isotopic composition of biological samples can provide a fingerprint for sources of Pb (Rabinowitz, 1995; Spencer et al., 2000; Manahan, 2000; Kurkjian and Flegal, 2003).

Aquatic organisms in the PRE are one of the most important sources of seafood for people in the Pearl River Delta (PRD) region (Fu et al., 1995). Extensive

studies on the ecosystem in the PRE, such as on sediments (Li et al., 2000a, 2000b and 2001) and water (Ho and Hui, 2001), have recently been carried out. However, only a few of the studies have focused on the aquatic organisms in the PRE (Chen, 1995; Fu et al., 1995), and none on the potential pathway of metal contaminants in this region.

Studies have recently been conducted on the stable Pb isotopic compositions of various ecosystems in the PRD region, such as agricultural soils (Wong et al., 2002), air depositions (Wong et al., 2003), and sediments (Liu et al. 2003; Ip et al., 2004). These environmental media have distinctive ranges of stable Pb isotopic ratios. According to these studies, automobile emissions and industrial discharges are the major sources of anthropogenic Pb in the PRD region. These research projects have provided an important database on Pb isotopic signatures in the PRD region. This database may help in efforts to evaluate the accumulation and biological pathways of heavy metals in aquatic organisms in the PRE. Trace metal accumulation in aquatic organisms depended on several factors, including (i) the environmental concentrations of metals in water and sediments; (ii) the species of organisms; (iii) body size and age of organisms. Different concentrations of trace metals can also be found in different organs in the same biological sample. However, this study mainly focused on the general trace metal burden in aquatic organisms, and the potential major pathways for metal contaminants in the estuarine environment. The common species of seafood in the PRE were collected in the present study. The whole meat tissue of the samples was used in this study in order to examine the general situation of trace metal contamination of aquatic organisms in the PRE. Therefore, the present study aims (1) to assess the accumulation of trace metals in several groups of common aquatic organisms in the PRE, and (2) to identify possible Pb sources for aquatic organisms

using the Pb isotopic signature in aquatic organisms and various environmental media in the region.

## **2. Materials and Methods**

A total of 58 samples of aquatic organisms were collected at seven sampling sites in the PRE in April 2003 with the assistance of the South China Sea Institute of Oceanology under the Chinese Academy of Sciences. The seven sampling locations are depicted in the Fig. 1. The samples include four common estuarine groups: fish, crab, shrimp, and shellfish. Sixteen species of fish, one species of crab, two species of shrimp, and three species of shellfish were sampled in the present study. The details of the sampled species are summarized in Table 1.

All of the samples of aquatic organisms were individually stored in polyethylene bags at 4-6°C immediately after collection prior to the laboratory analysis. After washing with tap water and distilled and deionised water (DIW), the samples were stored frozen at -20°C prior to freeze-drying. The samples were freeze-dried at -45°C for 3 days. Whole tissues of the samples were grounded homogeneously. All of the freeze-dried and grounded samples were stored in a dessicator prior to undergoing further chemical analyses.

The samples of aquatic organisms were digested using strong acid digestion according to the method from the USEPA (1999) with some modifications. About 0.500g of grounded aquatic organism samples were weighed and placed in Pyrex test tubes pre-cleaned with high purity nitric acid. The nitric acid used in the present study was in high purity grade, which contained usually less than 0.1 ppb trace metals

(except Al, Ca, Mg and Zn in less than 0.5 ppb). Five ml of high purity nitric acid was added to each tube, and the tubes were left overnight to be slowly digested. Another 3 ml of high purity nitric acid and 1 ml of perchloric acid were added to each tube the next day. Each mixture was gently shaken using a vortex and then placed in an aluminium heating block (FOSS TECATOR 2000). The heating process for the digestion was set up according to the following temperature scheme: 50°C for 8 hrs, 75°C for 2 hrs, 100°C for 2 hrs, 125°C for 3 hrs, and 150°C until complete dryness was achieved. After the test tubes were removed from the heating block and cooled down, 10 ml of 5% high purity nitric acid were added to the residue. The mixture was then heated at 70°C for 1 hr. The heated mixture was shaken gently and poured into polyethylene tubes. The tubes were centrifuged with centrifugal force around 150 N for 10 minutes prior to determining the concentration of metals. The concentrations of Al, Ca, Cu, Fe, Mg, Mn, and Zn were measured by inductively coupled plasma-atomic emission spectrometry (ICP-AES; *Perkin Elmer Optima, 3300DV*) (Li and Thornton, 1992; USEPA, 1999; Li et al., 2001). Due to the low concentrations of Cd, Co, Cr, Ni, Pb, and V, the concentrations of these elements were determined by inductively coupled plasma – mass spectrometry (ICP-MS; *Perkin Elmer Sciex Elan 6100 DRC<sup>plus</sup>*). Selected samples of aquatic organisms were also analysed for Pb isotopic composition by ICP-MS. All of the analytical solutions for Pb isotopic composition were diluted to about 30 µg/L Pb using 5% high purity nitric acid.

The quality controls for the strong acid digestion method included reagent blanks, duplicate samples, and standard reference materials (NIST SRM 1566a and DORM-2). The QA/QC results showed no sign of contamination in all the analysis. The recovery rates for most of the trace metals in the reference materials were around 80% - 115%, except for Al and Ni (62% and 147%, respectively). To detect whether

there was any contamination and drift during the measurements, quality control standards were used during the determination of elemental concentration and isotopic compositions at every 10 samples for the ICP-AES analysis and every 4 samples for the ICP-MS analysis. For the Pb isotopic analysis, an international standard reference material (NIST SRM 981, common lead) was used for calibration and analytical control. The relative standard deviation of each sample measurement was  $< 0.3\%$ . The average measured ratios of  $^{204}\text{Pb}/^{207}\text{Pb}$ ,  $^{206}\text{Pb}/^{207}\text{Pb}$ , and  $^{208}\text{Pb}/^{207}\text{Pb}$  of the SRM 981 were  $0.0645 \pm 0.0003$ ,  $1.0931 \pm 0.0023$ , and  $2.3718 \pm 0.0045$ , respectively. These values were very close to the certified standard values (0.0646, 1.0933, and 2.3704, respectively).

### **3. Results and Discussion**

#### *3.1. Trace metal concentrations of the aquatic organisms*

The trace metal concentrations of the fish, crab, shrimp, and shellfish collected in the PRE are summarized in Table 2. A comparison between the data of the present study and those of previous studies conducted in Hong Kong (Tam and Mok, 1991) and China (Wei et al., 2002) is presented in Table 3, together with the guidelines for assessing the aquatic organisms in China and the baseline reference values in Norway (Green and Knutzen, 2003). The mean and median concentrations of Cd and Cu in crab, shrimp, and shellfish; and Cr in shellfish exceeded the threshold values recommended by China's assessment guidelines, suggesting that the concentrations of these trace metals in these species from the PRE were elevated. In addition, the mean Pb concentrations in fish also exceeded the China's assessment guidelines, while the median concentration of Pb in fish was below the guidelines. This indicated that a

small number of fish samples accumulated high concentrations of Pb in their bodies. The highest concentration of Pb was found in fish, *Siganus oramin* (30.7 mg/kg wet weight). This value was about 30 times higher than the recommended values in the Food Assessment Guidelines of China. According to the present results, the concentrations of Pb in 20% fish samples were above the guideline level. Furthermore, the concentrations of Pb in fish were considerably higher than in the other organisms. This group of aquatic organisms needs to pay special attention for their Pb accumulation. Although the mean concentrations of other metals in these species were below the guideline values, some samples also had high concentrations of one or more metals due to the wide ranges of metal concentrations in these aquatic organisms.

Among the four groups of aquatic organisms, the concentrations of Cd in shrimp were the highest. The highest concentration of Cd was 2.10 mg/kg wet weight in shrimp, *Metapenaeus ensis*. This value was more than 40 times higher than the assessment standard of China. The concentrations of Cd were also noticeably elevated in both crab and shellfish. However, the concentration of Cd was significantly lower in fish in comparison with other species. Among the species studied, the highest concentrations of Co, Cr, Cu, Ni, V, and Zn were observed in shellfish. The concentrations of these elements in shellfish were 3 to 20 times higher than that in other species. These findings suggest that shellfish could accumulate trace metals more efficiently from water and sediment.

In general, the average concentrations of trace metals in aquatic organisms in the PRE were higher than the reported values in other parts of China (see Tables 2 and 3). In addition, the concentrations of trace metals in the present study were higher than those in previous studies of the same region. According to Wei et al. (2002), the concentrations of Cd, Cr, Cu, Ni, Pb, and Zn in fish collected in 2000 were 0.03, 0.16,



0.18, 0.28, 0.51, and 1.01 mg/kg, respectively; while those in the fish samples in the present study were  $0.0409 \pm 0.0289$ ,  $0.667 \pm 0.756$ ,  $1.81 \pm 1.74$ ,  $0.653 \pm 0.550$ ,  $2.202 \pm 6.02$ , and  $18.4 \pm 6.25$  mg/kg, respectively. These figures represent increases of 1.4 to 18 times between the two studies. The results might indicate that concentrations of trace metals in fish in the PRE increased in the last few years.

The concentrations of trace metals in fish from the PRE were generally over 200 times higher than the reference values in Norway (see Table 2 and 3). The concentrations of Cd, Cu, Pb, and Zn in shellfish in the PRE were generally enriched. The enrichment factors for fish in the PRE were 409, 240, 73, 333, and 780 for Cd, Cu, Pb, and Zn, respectively. For shellfish in the PRE, the enrichment factors of the mean Cd, Cu, Pb, and Zn concentrations were 2.9, 22, 1.6, and 2.0, respectively.

At low concentrations, Zn and Cu are essential elements for the growth of organisms (WHO, 1996). They are normally the most abundant trace elements in aquatic organisms (Parsons, 1998; Chien et al., 2002; Wei et al., 2002; Usero et al., 2003). At low concentrations, Co, Ni, and V are probably elements essential to organisms. Cd and Pb are non-essential elements and are toxic even at low concentrations. Cd is usually present in low concentrations in different environmental media; for example, in sediments (Li et al., 2000a, 2000b and 2001; Lin et al., 2002), in soils (Wong et al., 2002), and in atmospheric depositions (Wong et al., 2003).

The four groups of aquatic organisms collected in the PRE showed quite different patterns of metal accumulation, although the metal concentrations of the aquatic organisms showed large variations within the same group. Crab and shrimp showed similar ranges of metal concentrations and relative orders of mean trace metal concentrations: Cu > Zn > Cd > Ni > Cr > V > Pb > Co for crab and Cu > Zn > Cd > Ni >

V > Cr > Pb > Co for shrimp. This exemplified the common feeding habits and living behaviours of these two aquatic organisms. However, fish and shellfish had different patterns of trace metal accumulation. The accumulations of trace metals in fish were: Zn > Pb > Cu > Cr > Ni > V > Co > Cd, while those in shellfish were: Zn > Ni > Co > Cu > Cr > Cd > Pb. The accumulation of Pb in fish was particularly significant in these samples.

The different feeding habits and living modes of shellfish, shrimp, crab, and fish as well as the different aquatic geochemistry of the trace metals significantly affect the intake, bioassimilation, and subsequent bioaccumulation of trace metals in these organisms. Although the trace metal concentrations in different species of aquatic organisms in the same group were in a wide range of variations, the aquatic organisms in different group also showed significant metal accumulation patterns (see Table 2). This demonstrated that aquatic organisms in different groups had different accumulation mechanisms for trace metals. Shellfish is a filter feeder and mainly filters fine suspended matter as its source of food. Furthermore, shellfish are benthic organisms, and are usually relatively immobile or sessile. Based on the feeding mode and living habits of shellfish, the trace metal content of shellfish most likely reflects the quality of the water and sediment in the aquatic environment, including the accumulation of both dissolved and suspended trace metals. The significantly elevated concentrations of Co, Cr, Cu, Ni, V, and Zn found in shellfish likely resulted from the fact that their primary source of food is suspended matter, in particular, suspended fine sediment near or on the sea floor. It might also be partly attributed to the solubility of these trace metals in an aquatic environment. This is because the ratio of the dissolved metal concentration to the total metal concentration (dissolved/total) generally increases in the following order: Pb < Cd < Cr < Ni < Cu < Zn (Foster and

Charlesworth, 1996). Hence, in an aquatic environment, Cr, Ni, Cu, and Zn are more soluble and bioavailable than Pb and Cd.

Similar to shellfish, crab and shrimp are also benthic organisms that generally live on or near the sea floor and are capable of travelling in distance. As scavengers, crab and shrimp have similar feeding patterns. They tend to feed on detritus and, sometimes, small crustaceans and fish on or near the sea floor as well as on floating materials. Among the different aquatic organisms, fish are probably the most mobile and capable of travelling a long distance. However, the fish samples collected in this study were mainly live near the sea floor, and with short travelling distance (e.g. *Collichthys lucidus*, *Platycephalus indicus*, *Nibea albiflora*, *Zebrias zebra*, *Cynoglossus macrolepidotus*). Furthermore, fish is also on a high trophic level in the food chain as compared to the other three types of organisms; hence, their diet is probably the most diverse of the species studied here.

Moreover, the comparatively low bioaccumulation of Pb in shellfish, crab, and shrimp showed that the bioassimilation and bioavailability of Pb is limited in an aquatic environment, especially near the sea floor. Pb generally becomes immobile or bound to organic complexes shortly after its deposition in water. Based on the Pb concentrations of the different aquatic organisms (see Table 2), the direct intake and subsequent bioassimilation of Pb by shellfish and even crustaceans, in the form of suspended matter and detritus near or on the sea floor, might be of secondary importance in the PRE. The results suggest that the primary importance of the bioaccumulation of Pb in aquatic organisms could be bioassimilation in the food chain and/ or exposure in water. As mentioned previously, fish is situated at a higher trophic level in comparison with other three groups of organisms. Not only are their sources of food the most diverse, they also require a large quantity of food compared to the

other organisms. These factors could lead to the bioassimilation and bioaccumulation of Pb in fish over time (Manahan, 2000; Jacobson et al., 2000). The particularly high Pb concentrations in fish might be due to the bioaccumulation of Pb in some species of fish, such as *Siganus oramin* (spinefoot), *Collichthys lucidus* (croaker), and *Cyneglossus macrolepidotes* (large scaled tongue sole) found in the present study. The major food sources of the abovementioned species are small shrimps and fish. A number of studies also revealed that fish have a tendency to accumulate trace metals at high levels (Allen, 1994; Karadede and Unlu, 2000).

### 3.2. Stable Pb isotope compositions in aquatic organisms

The means and ranges of the  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios in the aquatic organisms collected from the PRE are presented in Table 4. The  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios of the aquatic organisms ranged from 1.161 to 1.193 and 2.438 to 2.494, respectively. The Pb accumulated in aquatic organisms can result from Pb derived from natural processes of weathering, erosion, and transport of bedrocks, as well as from a range of anthropogenic activities in the aquatic environment. According to Zhu (1995), the Pb isotopic ratios of the background geological materials in the PRD region ranged from 1.183 to 1.199 for  $^{206}\text{Pb}/^{207}\text{Pb}$  and 2.468 to 2.497 for  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios. From the present results, a large proportion of the  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of the aquatic organisms were lower than those of the geological materials. The mean  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of the aquatic organisms descended in the following order: shellfish > crab > shrimp > fish. This is possibly related to their habits, food sources, and trophic level.

Aquatic organisms are exposed to at least four sources of trace metals in the aquatic system, including water, sediments, plankton, and detritus in the water columns (Kneip and Lauer, 1973; Stokes, 1979; Hare, 1992; Roy and Hare, 1999; Barata et al., 2002). In order to identify the potential pathways of the anthropogenic Pb that had accumulated in the aquatic organisms, the relationship between the  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios of the selected aquatic organisms in the PRE and the major sources for the input of Pb in the PRD are shown in Fig. 2. The possible sources of Pb include surface sediments in the PRE (Ip et al., 2004), atmospheric deposits (Wong et al., 2003), natural soils (Wong et al., 2002), and some other known anthropogenic sources in the PRD (Zhu et al., 2001). The atmospheric deposition in the PRD was taken into account in the assessment because atmospheric deposition is one of the principal pathways of transport for anthropogenic Pb (Jikells, 1995; Neff, 2002; Reuer and Weiss, 2002). Therefore, the Pb isotopic signature of the atmospheric deposition was used to represent the anthropogenic Pb in the water columns. Lead is usually weakly associated with air particles and can be easily dissolved in water (Foster and Charlesworth, 1996). They are therefore highly reactive and biologically available (Gelinias and Schmit, 1997). The Pb isotopic signature of natural soil is used to represent the Pb derived from natural weathering, erosion, and different processes of transport. In order to examine the significant differences between the  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of the geological materials and aquatic organism samples, a paired sample *t* test was performed. The factors of the paired samples *t* test for the  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of fish, crab, shrimp and shellfish samples compared with those of the geological materials were 5.844 (degree of freedom, *df* = 4), 3.913 (*df* = 4), 8.262 (*df* = 4), and 4.819 (*df* = 2), respectively. Therefore, the Pb isotopic signatures of these biological samples had significant differences (*p* < 0.05)

in comparison with the geological materials, and provided a useful tool for distinguishing the relative contributions from various natural and anthropogenic sources.

As shown in Fig. 2, all of the Pb isotopic ratios of the aquatic organisms ranged between those of the surface sediments and those of the atmospheric depositions. The Pb isotopic ratios of a small number of aquatic organisms were similar to those of the surface sediments. The Pb accumulated in these aquatic organisms might be derived from the sediments. A large proportion of the aquatic organisms had lower  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios than the surface sediments, indicating that most of the aquatic organisms received contributions from anthropogenic Pb.

In general, three groups of aquatic organisms could be categorized on the basis of their Pb isotopic signatures in the present study. The aquatic organisms in Group 1 possessed Pb isotopic compositions similar to those of the air depositions of the PRD region. Fish were the dominant species in this group. The result indicated that some fish had very similar Pb isotopic signatures as the anthropogenic sources. The aquatic organisms in Group 2 had Pb isotope signatures similar to those of the PRE sediments. The Pb isotopic ratios of the aquatic organisms in Group 3 were in between those of the two groups. Most of the Pb isotopic compositions of shrimp, crab, and shellfish belonged to Group 3, indicating that these aquatic organisms accumulated Pb inputs from various sources, with the Pb being derived from anthropogenic sources (e.g., air depositions) and surface sediments, or food sources. Fish generally received more contributions from Pb derived from anthropogenic sources (e.g., air depositions) than shrimp, crab, and shellfish in the PRE, possibly because their dominant habitat is the water and also because of their position in the food chain.

In general, a significant linear relationship between the  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios possibly suggests the binary mixing of two end-members with different isotopic compositions (Farmer et al., 1996; Wong and Li, 2004). The Pb isotopic data in this study do not form a single linear correlation (Fig. 2), suggesting that more than two end-members were involved. The different Pb isotopic signatures were mainly due to the different uptake efficiencies of Pb from various sources by the organisms. The  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios in fish were significantly correlated ( $r^2 = 0.764$ ), and those in shrimp and crab were not correlated ( $r^2 = 0.126$ ). Fish can bioaccumulate particle Pb through their gills (Tao et al., 1999). As fishes are more mobile in water columns, they are likely to be more exposed to weakly soluble and potentially bioavailable Pb originating from anthropogenic atmospheric depositions and/or from the inputs derived from the discharges of wastewater. The present study indicated that fish might be bioassimilating Pb from anthropogenic sources such as atmospheric deposits suspended in water columns. Previous studies have also reported that the highest trace metal concentrations in fish were found in their gills (Eisler, 1979; Wang and Fisher, 1996; Barata et al., 2002). This issue can be further investigated in the future by examining the differences in the Pb isotopic composition in the gills, livers, stomachs and fleshes of fish.

#### **4. Conclusions**

High concentrations of trace metals were generally found in shellfish, while the highest concentrations of Pb were found in the fish of the Pearl River Estuary. The highly comparable concentrations of Cd and Cu in shellfish, crab, and shrimp were

partly attributed to their consumption of detritus materials. The differences in the patterns of accumulation of Pb in these aquatic organisms were mainly attributed to the solubility of the metal in an aquatic environment. In addition, the differences in the feeding habitats of these organisms also affected their physiological responses to different trace metals. The significantly elevated concentrations of Pb and low  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios in fish compared with other organism samples could be attributed to the bioaccumulation of Pb from weakly soluble and potentially bioavailable Pb originating from anthropogenic sources and a wide variety of food sources.

### **Acknowledgements**

The work described here was supported by the Chinese Academy of Sciences (CAS Innovation Program ZKCX-2-212), the Research Grants Council of the Hong Kong SAR Government (PolyU 5057/99E and PolyU 5147/03E) and the Area of Excellence Scheme under the University Grants Committee of the Hong Kong Special Administration Region, China (Project No. AoE/P-04/2004).

### **References**

- Allen, P. 1994., Accumulation profiles of lead and the influence of cadmium and mercury in *Oreochromis aureus* during chronic exposure. *Toxicological and Environmental Chemistry* 44, 101-112.
- Barata, C., Markich, S.J., Baird, D.J., Soares, A.M.V.M., 2002. The relative importance of water and food as cadmium sources to *Daphnia magna* Straus. *Aquatic Toxicology* 61, 143-154.
- Blackmore, G., Morton, B., Huang, Z.G., 1998. Heavy metals in *Balanus amphitrite* and *Tetraclita squamosa* (Crustacea: Cirripedia) collected from the coastal waters of Xiamen, China. *Maine Pollution Bulletin* 36, 32-40.



- Bollhöfer, A., Rosman, K.J.R., 2000. Isotopic source signatures for atmospheric lead: The Southern Hemisphere. *Geochimica et Cosmochimica Acta* 64, 3251-3262.
- Bollhöfer, A., Rosman, K.J.R., 2001. Isotopic source signatures for atmospheric lead: The Northern Hemisphere. *Geochimica et Cosmochimica Acta* 65, 1727-1740.
- Bryan, G.W., 1979. Bioaccumulation of marine pollutants. *Philosophical Transactions of the Royal Society London Series B* 286, 483-505.
- Canadian Food Inspection Agency, 2004. Canadian guidelines for chemical contaminants and toxins in fish and fish products. Canadian Food Inspection Agency, Canada. Available (31 March 2004) <http://www.inspection.gc.ca/english/anima/fispoi/guide/chme.shtml>
- Chen, J., 1995. Study on the environmental protection countermeasures of Pearl River coast zones, in: Wang, C.K., Chu, K.H., Chen, Q.C., Ma, X.L. (Eds.), *Environmental Research in Pearl River and Coastal Areas*. Guangdong Higher Education Press, Guangzhou, pp. 153-161.
- Chen, J.S., Zhou, J.Y., 1992. *Heavy Metal Research in the Aquatic Environmental of China*, Environmental Science Press of China, China.
- Chien, L.C., Hung, T.C., Choang, K.Y., Yeh, C.Y., Meng, P.J., Shieh, M.J., Han, B.C., 2002. Daily intake of TBT, Cu, Zn, Cd and As for fishermen in Taiwan. *The Science of the Total Environment* 285, 177-185.
- Eisler, R., 1979. Copper accumulation in coastal and marine biota, in: Nriagu, J.O. (Ed.), *Copper in the Environment Part 1: Ecological Cycling*. A Wiley-Interscience Publication, New York.
- Fang, Z.Q., Cheung, R.Y.H., Wong, M.H., 2001. Heavy metal concentrations in edible bivalves and gastropods available in major markets of the Pearl River Delta. *Journal of Environmental Sciences* 13, 210-217.
- Farmer, J.G., Eades, L.J., Mackenzie, A.B., Kirika, A., Bailey-Watts, T.E., 1996. Stable lead isotope record of lead pollution in Loch Lomond sediments since 1630 A.D. *Environmental Science and Technology* 30, 3080-3083.
- Fu, Y.Y., Yin, J.Q., Chen, Q.C., Huang, L.M., Wong, C.K., 1995. Distribution and seasonality of marine zooplankton in the Pearl River Estuary, in: Wang, C.K., Chu, K.H., Chen, Q.C., Ma, X.L. (Eds.), *Environmental Research in Pearl River and Coastal Areas*. Guangdong Higher Education Press, Guangzhou, pp. 25-33.
- Foster, I.D.L., Charlesworth, S.M., 1996. Heavy metals in the hydrological cycle: trends and explanation. *Hydrological Processes* 10, 227-261.
- Gelinas, Y., Schmit, J.P., 1997. Extending the use of the stable lead isotope ratios as a tracer in bioavailability studies. *Environmental Science and Technology* 31, 1968-1972.
- Goodwin, T.H., Young, A.R., Holmes, M.G.R., Old, G.H., Hewitt, N., Leeks, G.J.L., Packman, J.C., Smith, B.P.G., 2003. The temporal and spatial variability of sediment transport and yield within the Bredford Beck catchment, West Yorkshire. *The Science of the Total Environment* 314-316, 475-494.
- Green, N.W., Knutzen, J., 2003. Organohalogenes and metals in marine fish and mussels and some relationships to biological variables at reference localities in Norway. *Marine Pollution Bulletin* 46, 362-377.

- Hare, L., 1992. Aquatic insects and trace metals: bioavailability, bioaccumulation and toxicity. *Critical Reviews of Toxicology* 22, 327-369.
- Hinrichs, J., Dellwig, O., Brumsack, H.J., 2002. Lead in sediments and suspended particulate matter of the German Bight: natural versus anthropogenic origin. *Applied Geochemistry* 17, 621-632.
- Ho, K.C., Hui, K.C.C., 2001. Chemical contamination of the East River (Dongjiang) and its implication on sustainable development in the Pearl River Delta. *Environment International* 26, 303-308.
- Hoven, H.M., Gaudette, H.E., Short, F.T., 1999. Isotope ratios of  $^{206}\text{Pb}/^{207}\text{Pb}$  in eelgrass, *Zostera marina*, indicate sources of Pb in an estuary. *Marine Environmental Research* 48, 377-387.
- Ip, C.C.M., Li, X.D., Zhang, G., Farmer, J.G., Wai, O.W.H., Li, Y.S., 2004. Over one hundred years of trace metal fluxes in the sediment of the Pearl River Estuary, South China. *Environmental Pollution* 132, 157-172.
- Jacobson, M.C., Charlson, R.J., Rodhe, H., Orians, G.H., 2000. *Earth system science: from biogeochemical cycles to global change*, Academic Press, San Diego, California.
- Jickells, T., 1995. Atmospheric inputs of metals and nutrients to the oceans: their magnitude and effects. *Marine Chemistry* 48, 199-214.
- Karadede, H., Unlu, E. 2000., Concentrations of some heavy metals in water, sediment and fish species from the Ataturk Dam Lake (Euphratees), Turkey. *Chemosphere* 41, 1371-1376.
- Kneip, T.J., Lauer, G.J., 1973. Trace metal concentration factors in aquatic ecosystems, in: Ahuja, S., Cohen, E.M., Kneip, J.L., Lambert, J.K., Sweig, G. (Eds.), *Chemical Analysis of the Environment and Other Modern Techniques*. Plenum Press, New York, pp. 43-62.
- Kurkjian, R., Flegel, A.R., 2003. Isotopic evidence of the persistent dominance of blood lead concentration by previous gasoline lead emissions in Yerevan, Armenia. *Environmental Research* 93, 308-315
- Labonne, M., Othman, D.B., Luck, J.M., 2001. Pb isotopes in mussels as tracers of metal sources and water movements in a lagoon (Thau Basin, S. France). *Chemical Geology* 181, 181-191.
- Li, R., Feng, S., Jiang, W.P., 1997. *Industrial Development of Hong Kong and Pearl River Delta: Opportunities and Strategies*. The Hong Kong Polytechnic University: Department of Manufacturing Engineering and The Hong Kong Association for the Advancement of Science and Technology Ltd., Hong Kong.
- Li, X.D., Thornton, I., 1992. Multi-element contamination in soil and plant in the old mining area. *Applied Geochemistry* S2, 51-56
- Li, X.D., Wai, O., Li, Y.S., Coles, B., Ramsey, M.H., Thornton, I., 2000a. Heavy metal distribution in sediment profiles of the Pearl River estuary, South China. *Applied Geochemistry* 15, 567-581.
- Li, X.D., Shen, Z.G., Wai, O., Li, Y.S., 2000b. Chemical partitioning of heavy metal contaminants in sediments of the Pearl River Estuary. *Chemical Speciation and Bioavailability* 12, 17-25.

- Li, X.D., Shen, Z.G., Wai, O., Li, Y.S., 2001. Chemical forms of Pb, Zn and Cu in the sediment profiles of the Pearl River Estuary. *Marine Pollution Bulletin* 42, 215-223.
- Lin, S., Hsieh, I.J., Huang, K.M., Wang, C.H., 2002. Influence of the Yangtze River and grain size on the spatial variations of heavy metals and organic carbon in the East China Sea continental shelf sediments. *Chemical Geology* 182, 377-394.
- Liu, W.X., Li, X.D., Shen, Z.G., Wang, D.C., Wai, O.W.H., Li, Y.S., 2003. Multivariate statistical study of heavy metal enrichment in sediments of the Pearl River Estuary. *Environmental Pollution* 121, 377-388.
- Manahan, S.E., 2000. *Environmental Chemistry*. Lewis Publishers, Boca Raton.
- Neff, J.M., 2002. *Bioaccumulation in Marine Organisms: Effect of Contaminants from Oil Well Produced Water*. Elsevier, Amsterdam, Boston.
- Parsons, E.C.M., 1998. Trace metal pollution in Hong Kong: implications for the health of Hong Kong's Indo-Pacific hump-backed dolphins (*Sousa chinensis*). *The Science of the Total Environment* 214, 175-184.
- Rabinowitz, M.B., 1995. Stable isotopes of lead for source identification. *Clinical Toxicology* 33, 649-655.
- Reuer, M.K., Weiss, D.J., 2002. Anthropogenic lead dynamics in the terrestrial and marine environment. *Philosophical Transactions of the Royal Society of London Series A* 360, 2889-2904.
- Ritson, P.I., Bouse, R.M., Flegal, A.R., Luoma, S.N., 1999. Stable lead isotopic analyses of historic and contemporary lead contamination of San Francisco Bay estuary. *Marine Chemistry* 64, 71-83.
- Roy, I., Hare, L., 1999. Relative importance of water and food as cadmium sources to the predatory insect *Sialius celata* (Megaloptera). *Canadian Journal of Fisheries and Aquatic Sciences* 56, 1143-1149.
- Semlali, R.M., Oort, F.V., Denaix, L., Loubet, M., 2001. Estimating distributions of endogenous and exogenous Pb in soils by using Pb isotopic ratios. *Environmental Science and Technology* 35, 4180-4188.
- Spencer, K., Shafer, D.J., Gauldie, R.W., DeCarlo, E.H., 2000. Stable lead isotope ratios from distinct anthropogenic sources in fish otoliths: a potential nursery ground stock marker. *Comparative Biochemistry and Physiology Part A* 127, 273-284.
- Stokes, P.M., 1979. Copper accumulations in freshwater biota, in: Nriagu, J.O. (Ed.), *Copper in the Environment Part 1: Ecological Cycling*, A Wiley-Interscience Publication, New York.
- Sugden, C.L., Frammer, J.G., MacKenzie, A.B., 1993. Isotopic ratios of lead in contemporary environmental material from Scotland. *Environmental Geochemistry and Health* 15, 59-65.
- Tam, S.Y.K., Mok, C.S., 1991. Metallic contamination in oyster and other seafood in Hong Kong. *Food Additives and Contaminants* 8, 333-342.
- Tao, S., Liu, R., Dawson, J., Cao, J., Li, B., 1999. Uptake of particulate lead via the gills of fish (*Carassius auratus*). *Archives of Environmental Contamination and Toxicology* 37, 352-357.

- USEPA., 1999. Determination of Metals in Ambient Particulate Matter Using Inductively Coupled Plasma Mass Spectrometry (ICP/MS) – EPA/625/R-96/010a. Centre for Environmental Research Information, Office of Research and Development, US Environmental Protection Agency, Cincinnati.
- Usero, J., Izquierdo, C., Morillo, J., Gracia, I., 2003. Heavy metals in fish (*Solea vulgaris*, *Anguilla aguilla* and *Liza aura*) from salt marshes on the southern Atlantic coast of Spain. *Environmental International* 169, 1-8.
- Wang, C.K., Chu, K.H., Chen, Q.C., Ma, X.L., 1995. *Environmental Research in Pearl River and Coastal Areas*. Guangdong Higher Education Press, Guangzhou.
- Wang, W.X., Fisher, N.S., 1996. Assimilation of trace metals and carbon by the mussel *Mytilus edulis*: effects of food composition. *Limnology and Oceanography* 41, 197-207.
- Wei, T.L., Yang, W.L., Lai, Z.N., Zhang, Q., Liu, M., 2002. Residues of heavy metals in economic aquatic animal muscles in Pearl River estuary, South China. *Journal of Fishery Sciences of China* 9, 172-176.
- WHO., 1996. *Trace Elements in Human Nutrition and Health*. World Health Organization, Geneva.
- Wong, S.C., Li, X.D., Zhang, G., Qi, S.H., Min, Y.S., 2002. Heavy metals in agricultural soils of the Pearl River Delta, South China. *Environmental Pollution* 119, 33-44.
- Wong, C.S.C., Li, X.D., Zhang, G., Qi, S.H., Peng, X.Z., 2003. Atmospheric deposition of heavy metals in the Pearl River Delta, China. *Atmospheric Environment* 37, 767-776.
- Wong, C.S.C., Li, X.D., 2004. Pb contamination and isotopic composition of urban soils in Hong Kong. *The Science of the Total Environment* 319, 185-195.
- Zheng, J.L., 1992. The study of heavy metals in the Pearl River Estuary, in: Chen, J.S., Zhou, J.Y. (Eds.), *Heavy Metal Research in the Aquatic Environment of China*. Environmental Science Press of China, China, pp. 369-387
- Zhu, B.Q., 1995. The mapping of geochemical provinces in China based on Pb isotopes. *Journal of Geochemical Exploration* 55, 171-181.
- Zhu, B.Q., Chen, Y.W., Peng, J.H., 2001. Lead isotope geochemistry of the urban environment in the Pearl River Delta. *Applied Geochemistry* 16, 409-417.

**TABLE 1. Aquatic organisms analysed in this study**

Groups of aquatic organisms	Common name	Scientific name of species	No. of samples	Sampling locations	Sample ID
Fish	Chinese herring	<i>Iisha elongata</i>	2	A, F	AY1, FY3
	flat head fish	<i>Platycephalus indicus</i>	3	A, F, G	AY2, FY5, GY7
	ray-finned fish	<i>Odontamblyopus rubicundus</i>	2	A, C	AY4, CY7
	ponyfish	<i>Leiognathus bin</i>	5	A, B, E, F, G	AY5, BY5, EY2, FY4, GY6
	white flower croaker	<i>Nibea albiflora</i>	1	B	BY2
	common mullet	<i>Mugil cephalus</i>	3	B, C, F	BY3, CY3, FY7
	zebra sole	<i>Zebrias zebra</i>	1	B	BY4
	large scaled tongue sole	<i>Cynoglossus macrolepidotus</i>	2	C, D	CY2, DY8
	croaker	<i>Collichthys lucidus</i>	5	C, D, E, F, G	CY5, DY2, EY1, FY2, GY1
	golden sardine	<i>Sardinella aurita</i>	1	C	CY6
	white sea bass	<i>Lates calcarifer</i>	1	D	DY3
	hilsa herring	<i>Macrura reeuesii</i>	1	D	DY4
	-	<i>Collicchthys gunther</i>	2	D, F	DY7, FY6
	sea horse	<i>Syngnathus linnaeus</i>	1	F	FY1
	spinefoot	<i>Siganus oramin</i>	1	G	GY8
	-	<i>Ambassidae, siganus forskal</i>	1	G	GY4
-	-	3	C, D, G	CY4, DY6, GY2	
Shrimp	Mantis shrimp	<i>Dictyosquilla foveolata</i>	7	A to G	AX1, BX2, CX1, DX2, EX1, FX1, GX1
	sand prawn	<i>Metapenaeus ensis</i>	6	A to F	AX2, BX1, CX2, DX1, EX2, FX2
Crab	redspot swimming crab	<i>Portunus pelagicus</i>	7	A to G	AP1, BP1, CP1, DP1, EP1, FP1, GP1
Shellfish	-	<i>Scapharca subcrenata</i>	1	A	AB1
	-	<i>Turritella bacillum keener</i>	1	B	BB2
	-	<i>Murex ttrapa</i>	1	B	BB4

**TABLE 2. Summary of trace metal concentrations (mg/kg, wet weight) in different sub-groups of aquatic organisms collected in the PRE**

Metals	Concentrations (mg/kg)	Fish (n=35)	Crab (n=7)	Shrimp (n=13)	Shellfish (n=3)
Cd	Mean ± standard derivation (S.D.)	0.0409 ± 0.0289	0.795 ± 0.506	0.835 ± 0.637	0.725 ± 0.305
	Median	0.0306	0.871	0.851	0.791
	Median of absolute deviations (mad)	0.0165	0.322	0.516	0.267
	Range	0.01 ~ 0.13	0.2 ~ 1.61	0.04 ~ 2.10	0.39 ~ 0.99
Co	Mean ± S.D.	0.100 ± 0.101	0.128 ± 0.065	0.0775 ± 0.0372	1.51 ± 2.44
	Median	0.0595	0.132	0.0583	0.105
	mad	0.0546	0.0299	0.0300	1.42
	Range	0.02 ~ 0.48	0.05 ~ 0.26	0.03 ~ 0.53	0.09 ~ 4.33
Cr	Mean ± S.D.	0.667 ± 0.756	0.411 ± 0.065	0.201 ± 0.131	1.17 ± 0.86
	Median	0.381	0.403	0.152	1.07
	mad	0.347	0.146	0.07512	0.809
	Range	0.11 ~ 4.27	0.14 ~ 0.76	0.08 ~ 0.53	0.37 ~ 2.08
Cu	Mean ± S.D.	1.81 ± 1.74	26.1 ± 24.4	28.0 ± 11.0	28.7 ± 42.6
	Median	0.381	24.4	27.8	6.08
	mad	1.01	6.89	3.38	26.4
	Range	0.15 ~ 7.55	16.3 ~ 41.8	15.2 ~ 56.2	2.28 ~ 77.8
Ni	Mean ± S.D.	0.653 ± 0.550	0.616 ± 0.359	0.560 ± 0.220	10.9 ± 17.4
	Median	0.428	0.529	0.493	0.890
	mad	0.309	0.124	0.182	10.1
	Range	0.17 ~ 2.08	0.26 ~ 1.39	0.26 ~ 0.99	0.73 ~ 31.0
Pb	Mean ± S.D.	2.20 ± 6.02	0.177 ± 0.062	0.135 ± 0.064	0.424 ± 0.234
	Median	0.405	0.176	0.103	0.298
	mad	1.86	0.0299	0.0525	0.144
	Range	0.09 ~ 30.7	0.09 ~ 0.29	0.04 ~ 0.23	0.28 ~ 0.69
V	Mean ± S.D.	0.616 ± 0.451	0.315 ± 0.089	0.252 ± 0.102	0.967 ± 1.44
	Median	0.428	0.289	0.268	0.158
	mad	0.308	0.0631	0.0821	0.852
	Range	0.15 ~ 1.93	0.17 ~ 0.42	0.07 ~ 0.41	0.12 ~ 2.63
Zn	Mean ± S.D.	18.4 ± 6.25	16.3 ± 2.92	15.8 ± 2.81	41.2 ± 39.0
	Median	18.8	17.7	16.0	19.1
	mad	4.53	1.68	1.85	23.0
	Range	8.78 ~ 30.26	12.2 ~ 19.9	11.0 ~ 20.0	18.3 ~ 86.3

**TABLE 3. Trace element concentrations in some aquatic organisms (mg/kg, wet weight) in other regions in China, reference concentrations from Norway, and the assessment guidelines in China**

Commodity (city/ country)	Cd	Cr	Cu	Ni	Pb	Zn
Shellfish (Hong Kong) <sup>a</sup>	0.49	0.21	-	-	0.254	-
Shellfish (Yangtze River Estuary), 1982 – 1983 <sup>b</sup>	0.42	-	14.9	-	2.08	37.8
Crab (Hong Kong) <sup>a</sup>	0.58	<0.05	-	-	0.04	-
Shrimp (Hong Kong) <sup>a</sup>	0.12	<0.05	-	-	0.08	-
Shrimp (Zhanjiang Harbour Bay), 1990 – 1994 <sup>b</sup>	0.04	-	1.56	0.11	0.42	13.48
Shrimp (PRE), 2000 <sup>b</sup>	0.04	0.15	1.28	0.27	0.50	2.60
Marine fish (Hong Kong) <sup>a</sup>	<0.02	<0.05	-	-	0.03	-
Fresh-water fish (Hong Kong) <sup>a</sup>	<0.02	<0.05	-	-	0.03	-
Fish (Yangtze River Estuary), 1982 – 1983 <sup>b</sup>	0.14	-	2.29	-	1.68	18.3
Fish (Yellow River Estuary), 1984 <sup>b</sup>	0.13	-	0.31	-	0.81	12.0
Fish (Zhanjiang Harbour Bay), 1990 – 1994 <sup>b</sup>	0.08	-	0.68	0.09	0.67	13.1
Fish (Guangdong Coastal waters), 1986 – 1988 <sup>b</sup>	0.03	-	0.77	0.14	0.22	6.26
Fish (PRE), 2000 <sup>b</sup>	0.03	0.16	0.18	0.28	0.51	1.01
Reference values:						
Fish, <i>Cod</i> (Norway), 2003 <sup>d</sup> (n=1184)	0.10	-	7.5	-	0.03	23.6
Shellfish, <i>Blue Mussels</i> (Norway), 2003 <sup>d</sup> (n=291)	0.25	-	1.30	-	0.26	20.29
Assessment Guidelines:						
Assessment Standard in China <sup>b</sup>	0.05	1.00	5.00	-	1.00	-
Action level for fish in Canada <sup>e</sup>	-	-	-	-	0.5	-
<sup>a</sup> Tam and Mok, 1991						
<sup>b</sup> Wei et al., 2002						
<sup>c</sup> Fang et al., 2001						
<sup>d</sup> Green and Knutzen, 2003						
<sup>e</sup> Canadian Food Agency, 2004						

**TABLE 4. The means ( $\pm$  standard derivations) and ranges of  $^{206}\text{Pb}/^{207}\text{Pb}$  and  $^{208}\text{Pb}/^{207}\text{Pb}$  ratios in the aquatic organisms collected from the PRE**

Commodity		$^{206}\text{Pb}/^{207}\text{Pb}$	$^{206}\text{Pb}/^{207}\text{Pb}$
Fish (n = 35)	Mean	$1.1789 \pm 0.0017$	$2.4662 \pm 0.0123$
	Range	1.1610 ~ 1.1933	2.4383 ~ 2.4889
Shrimp (n = 13)	Mean	$1.1796 \pm 0.0022$	$2.4842 \pm 0.0066$
	Range	1.1715 ~ 1.1908	2.4754 ~ 2.4942
Crab (n = 7)	Mean	$1.1808 \pm 0.0021$	$2.4766 \pm 0.0049$
	Range	1.1730 ~ 1.1875	2.4691 ~ 2.4861
Shellfish (n = 3)	Mean	$1.1826 \pm 0.0034$	$2.4748 \pm 0.0111$
	Range	1.1791 ~ 1.1894	2.4645 ~ 2.4865



## List of Figure Captions

Fig. 1. The location of sampling sites in the Pearl River Estuary (PRE)

Fig. 2. The correlation between the  $^{208}\text{Pb}/^{207}\text{Pb}$  and  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of aquatic organisms in the PRE




Fig. 1. The location of sampling sites in the Pearl River Estuary (PRE)




Fig. 2. The correlation between the  $^{208}\text{Pb}/^{207}\text{Pb}$  and  $^{206}\text{Pb}/^{207}\text{Pb}$  ratios of aquatic organisms in the PRE

<sup>a</sup> Wong et al., 2003; <sup>b</sup> Wong et al., 2002.