



Available online at www.sciencedirect.com





Journal of Sport and Health Science 11 (2022) 67-74

Original article

Changes in segment coordination variability and the impacts of the lower limb across running mileages in half marathons: Implications for running injuries

Tony Lin-Wei Chen^{a,b}, Duo Wai-Chi Wong^{a,c}, Yan Wang^{a,c}, Qitao Tan^a, Wing-Kai Lam^{b,d,*}, Ming Zhang^{a,c,*}

^a Department of Biomedical Engineering, Faculty of Engineering, Hong Kong Polytechnic University, Hong Kong SAR 999077, China

^b Li Ning Sports Science Research Center, Li Ning (China) Sports Goods Co. Ltd, Beijing 101111, China

^c Hong Kong Polytechnic University Shenzhen Research Institute, Shenzhen 518057, China

^d Department of Kinesiology, Shenyang Sports Institute, Shenyang 110102, China

Received 19 March 2020; revised 18 May 2020; accepted 10 August 2020 Available online 28 September 2020

2095-2546/© 2022 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license. (http://creativecommons.org/licenses/by-nc-nd/4.0/)

Abstract

Background: Segment coordination variability (CV) is a movement pattern associated with running-related injuries. It can also be adversely affected by a prolonged run. However, research on this topic is currently limited. The purpose of this study was to investigate the effects of a prolonged run on segment CV and vertical loading rates during a treadmill half marathon.

Methods: Fifteen healthy runners ran a half marathon on an instrumental treadmill in a biomechanical laboratory. Synchronized kinematic and kinetic data were collected every 2 km (from 2 km until 20 km), and the data were processed by musculoskeletal modeling. Segment CVs were computed from the angle-angle plots of selected pelvis-thigh, thigh-shank, and shank-rearfoot couplings using a modified vector coding technique. The loading rate of vertical ground reaction force was also calculated. A one-way MANOVA with repeated measures was performed on each of the outcome variables to examine the main effect of running mileage.

Results: Significant effects of running mileage were found on segment CVs ($p \le 0.010$) but not on loading rate (p = 0.881). Notably, during the early stance phase, the CV of pelvis frontal vs. thigh frontal was significantly increased at 20 km compared with the CV at 8 km (g = 0.59, p = 0.022). The CV of shank transverse vs. rearfoot frontal decreased from 2 km to 8 km (g = 0.30, p = 0.020) but then significantly increased at both 18 km (g = 0.05, p < 0.001) and 20 km (g = 0.36, p < 0.001).

Conclusion: At the early stance, runners maintained stable CVs on the sagittal plane, which could explain the unchanged loading rate throughout the half marathon. However, increased CVs on the frontal/transverse plane may be an early sign of fatigue and indicative of possible injury risk. Further studies are necessary for conclusive statements in this regard.

Keywords: Coupling angle; Endurance running; Motion capture analysis; Musculoskeletal modeling; Vector coding technique

1. Introduction

Participation in marathon running events has increased worldwide by more than 50% between 2009 and 2019.¹ The total number of finishers competing at various marathon events reached 1.2 million in 2019.² Additionally, more runners participated in

Peer review under responsibility of Shanghai University of Sport.

* Corresponding authors.

half-marathon or 10-km races than in full-marathon races.³ Running offers many benefits because it is easily accessible and promotes physical fitness while being preventive against chronic diseases.⁴ However, a long-distance run such as a marathon may not be suitable for everyone, particularly for those who have chronic cardiovascular diseases and are vulnerable to joint problems.⁴ It is thought that every runner possesses his or her own "preferred running pattern"⁵ that is energetically efficient and better attenuates the impact to the lower limb. Nevertheless, maintaining this movement pattern during an endurance run can be challenging.⁶ Changes in movement patterns can occur as

https://doi.org/10.1016/j.jshs.2020.09.006

Cite this article: Chen TL-W, Wong DW-C, Wang Y, Tan Q, Lam W-K, Zhang M. Changes in segment coordination variability and the impacts of the lower limb across running mileages in half marathons: Implications for running injuries. *J Sport Health Sci* 2022;11:67–74.

E-mail addresses: gilbertlam@li-ning.com.cn (W.-K. Lam), ming. zhang@polyu.edu.hk (M. Zhang).

mileage accumulates, which compromises force absorption^{7,8} and, thus, relates to overuse injury.^{6,9}

To date, discrete kinematic variables have been used predominantly to identify the attributes of movement patterns in various sports.¹⁰ However, existing studies investigating lower-limb joint or segment angulation in endurance runs have reported inconsistent outcomes.^{11–16} Indeed, joint or segment kinematics in isolation neither reflect effectively how the underlying segments are coordinated to produce resultant angular positions,¹⁷ nor do they provide a holistic view of the role of running-induced exertion in altering movement patterns.¹⁸ Prolonged physical activities have been found to disrupt neuromuscular synergies and inter-segment coordination spanning the joints, which is reflected by changes in the coordination pattern and its stride-to-stride variances over time.¹⁹

Coordination patterns can measure the timing and magnitude of relative motions between 2 body parts in any given task.²⁰ Coordination variability (CV) quantifies the degrees of fluctuation in the coordination pattern.²⁰ In recent decades, the CV of lower-limb couplings between adjacent joints or segments has received increasing research interest due to its potential link to injury. Although the underlying mechanism is not fully understood, changes in CVs during physical activities have been seen commonly in injured persons.²¹ Compared to healthy subjects, individuals with symptoms or injury histories tended to deviate from the "normal" CV range, which either increased^{18,22-25} or decreased^{23,26-28} in coordinate couplings that were hypothesized to be associated with the injured or painful regions. An example of this is knee pain and ankle instability. The changed CVs were considered to be an unstable compensatory movement strategy to avoid load on sensitized or painful regions.^{29,30} However, for increased CVs, another explanation is that the injured group may have decreased neuromuscular control, leading to poorly regulated movement coordination.^{21,31} We found this viewpoint intriguing because reduced neuromuscular control was also seen in healthy but fatigued people undergoing endurance runs,³²⁻³⁵ which likely exacerbated loading on the lower limb and increased injury risks.^{8,36,37} Accordingly, it was anticipated that runners would increase their lower-limb CVs across different time points in an endurance run such as a half marathon. Previous relevant research was scarce, so a study was needed to provide supporting evidence for this hypothesis.

Hence, this study aimed to investigate the CVs of segment couplings of the lower limb and ground reaction loading for healthy runners at various distance checkpoints during a half marathon. The segment couplings were analyzed using angle-angle plots, and their coordination variabilities were quantified by a modified vector coding technique.³⁸ As mentioned, it was hypothesized that the CVs of the selected segment couplings and the loading rates of the vertical ground reaction force would increase as mileage accumulated during the half marathon.

2. Methods

2.1. Participants

A total of 15 recreational runners (13 males, 2 females, age = 26.85 ± 6.02 years, height = 1.75 ± 0.08 m,

weight = 64.50 \pm 9.26 kg; mean \pm SD) were recruited from the local community by using flyers and posters. Of the runners. 13 were rearfoot strikers (foot strike index: 6.18%-25.74%), and the other 2 were midfoot strikers (foot strike index: 39.88%-46.98%). The foot strike index was calculated by averaging 10 strides from each runner's running trials at the 2-km checkpoint in the half marathon.³⁹ The participants had finished at least 3 marathon races annually for more than 2 years and reported no lower-limb injuries at the time of experiment entry based on consensual standards.⁴⁰ The sample-size calculation showed that 15 participants ensured a statistical power of 0.8, assuming a medium effect size and a significance level of 5%. The participants were fully informed of the research procedure, and they signed a written consent form before the study's commencement. The study was reviewed and approved by the Human Subjects Ethics Sub-Committee of the Hong Kong Polytechnic University (reference #HSEARS20170327001).

2.2. Experiment protocol

All running trials were conducted on a treadmill instrumented with 2 force platforms (Bertec, Columbus, OH, USA). The participants were given sufficient time to warm up and acclimate themselves to the research protocol. They were asked to use their own running shoes for the tests and to identify a comfortable running speed before the commencement of data collection. Each runner ran on the treadmill at a starting speed that they normally adopted for daily training. An investigator operated the instrument on a remote computer and finetuned the treadmill speed based on the runner's feedback. The procedure was repeated until runners found a comfortable speed at which they could be confident about finishing a half marathon on the treadmill.⁴¹

To facilitate motion capture, a total of 28 markers were attached to the following anatomical landmarks:⁴² acromioclavicular joints, between clavicles, seventh cervical, posterior/anterior iliac spines, lateral femoral epicondyles, lateral malleoli, calcaneal tuberosity, the heads of the first and fifth metatarsals, and the tip of the toe. Two additional markers were placed on the upper and lower thirds of the side of the thigh and shank segments, respectively. The marker set ensured that at least 3 markers were assigned to each segment of interest to enable 3-dimensional motion tracking and were also accommodated to the OpenSim environment. All markers were secured with skin-medical tape (Kinesio Tex Classic; Kinesio, Albuquerque, NM, USA) to avoid drop-off while running. Markers of the foot region were placed directly on the shoes.⁴³

Data collection began with a static calibration trial that was later used to scale an established musculoskeletal model. The participants then ran on the treadmill at their preferred speed for a half marathon. Kinematic and kinetic data were recorded for 10 s at every 2 km. To exclude the initial effects, data collection was started at 2 km and ended at the 20-km checkpoint. An optical-based motion-capture system with 10 cameras (T20; Vicon Motion Systems, Oxford, UK) and 2 force plates were synchronized to measure the marker trajectories and ground reaction forces at sampling rates of 250 Hz and 1600 Hz, respectively.

2.3. Musculoskeletal model

In the study, segment angles of interest were calculated using an established musculoskeletal model.⁴² The model featured 12 rigid body segments, 23 degrees of freedom, and 92 musculotendinous units. It had a ball-and-socket joint at the hip and permitted hinge joints at the knee, ankle, subtalar, and metatarsophalangeal. The raw kinematic data were first lowpass Butterworth filtered at 8 Hz⁴⁴ and put into the OpenSim platform (Version 4.0; OpenSim, National Centre for Simulation in Rehabilitation Research, Stanford, CA, USA). The generic model was scaled to the participant's geometry using the data from the static trials. Inverse kinematics⁴⁵ were implemented on the data of running trials to calculate the joint coordinates that best reproduced the movement patterns. Segment angles of the pelvis, thigh, shank, and rearfoot were expressed as the orientation of the segmental reference frame (fixed to the segment) with respect to the global coordinate system.³⁸ The segmental reference frames were aligned with the global frame at a neutral position, and their locations were defined previously.⁴⁶ Data for segment angles were normalized to the standard standing position.47

2.4. Data reduction and outcome measures

For each running gait cycle, data for segment angles were truncated to the stance phase and normalized to 101 time points.⁴⁸ Inter-segmental coordination was calculated from angle-angle plots of the couplings of interest, with the proximal segment comprising the abscissa and the distal segment comprising the ordinate.²⁸ The coupling angle (γ_i) was defined

as the angle with respect to the right horizontal axis created by the vector connecting the 2 consecutive time points (Eq. 1).⁴⁷ A modified vector coding technique³⁸ was employed in this study:

$$\gamma_i = \tan^{-1} \left(\frac{\theta_{D(i+1)} - \theta_{Di}}{\theta_{P(i+1)} - \theta_{Pi}} \right) \cdot \frac{180}{\pi} \qquad \qquad \text{Eq. (1)}$$

where *i* is the *i*th point of the stance phase. θ_D and θ_P represent the angles of the distal and proximal segments. Segment couplings of interest (Fig. 1) included pelvis frontal plane *vs.* thigh frontal plane, pelvis sagittal plane *vs.* thigh sagittal plane, thigh sagittal plane *vs.* shank sagittal plane, shank sagittal plane *vs.* rearfoot sagittal plane, and shank transverse plane *vs.* rearfoot frontal plane. These couplings were selected because they were closely related to the lower-limb loading response and injury development during running (e.g., the couplings of pelvis frontal *vs.* thigh frontal for iliotibial band syndrome and thigh transverse *vs.* rearfoot frontal for rearfoot overpronation).^{13,49,50}

Using circular statistics,⁵¹ coordination variability (CV_i) was calculated as the circular SD of the coupling angle at the *i*th point of the stance phase across 10 strides of data (Eqs. 2–5). This circular standard deviation was averaged across the following stance phases: early (first 24% of the stance phase), mid (25%–50% of the stance phase), and terminal stance (51%–100% of the stance phase).⁵²

$$\overline{x}_i = \frac{1}{n} \sum_{i=1}^n \cos \gamma_i \qquad \qquad \text{Eq. (2)}$$

$$\overline{y}_i = \frac{1}{n} \sum_{i=1}^n \sin \gamma_i \qquad \qquad \text{Eq. (3)}$$



Fig. 1. Segment couplings of interest for investigating coordination variability: (A) pelvis frontal vs. thigh frontal; (B) pelvis sagittal vs. thigh sagittal; (C) thigh sagittal vs. shank sagittal; (D) shank sagittal vs. rearfoot sagittal; (E) shank transverse vs. rearfoot frontal.

$$\overline{\gamma}_i = \tan^{-1}\left(\frac{\overline{y}_i}{\overline{x}_i}\right) \cdot \frac{180}{\pi}$$
 Eq. (4)

$$CV_i = \sqrt{2\left(1 - \sqrt{\overline{x_i}^2 + \overline{y_i}^2}\right)} \cdot \frac{180}{\pi} \qquad \text{Eq. (5)}$$

where *n* is the *n*th stride of the data, and \overline{x}_i and \overline{y}_i are the horizontal and vertical components, respectively, of the average coupling angle at the *i*th point of the stance phase.

The ground reaction force of each stride was filtered using a fourth-order 0-lag Butterworth filter with a lowpass frequency of 50 Hz.³⁹ The vertical average loading rate (VALR) and vertical instantaneous loading rate (VILR) of initial contact were computed using a previously established method.³⁹ The 20% and 80% points of the impact peak were identified in the curve of the vertical ground reaction force. VALR refers to the slope of the line through the 2 points, and VILR refers to the maximum slope of the lines connecting successive data points within the same region.

In a previous study, both segment CVs and loading rates were averaged across 10 strides at each distance checkpoint (2-20 km, at 2-km intervals). A total of 10 strides comprises the sufficient number of gait cycles needed to predict accurately coordination variability.⁴⁸ Computations of all outcome variables were accomplished using a custom MATLAB code (MathWorks, Natick, MA, USA).

2.5. Statistical analysis

The outcome variables were assessed for skewness prior to statistical analysis. Appropriate log data transform was performed if the skewness was found to be greater than $\pm 1.0^{53}$ Loading rate and segment CVs were compared among the distance checkpoints using one-way multivariate analyses of variances (MANO-VAs) repeated measures. The 2 measures of vertical loading rates and all measures of CVs at the early, mid, and terminal stance phases were 4 separate entries in the statistic model. Equality of variances was tested for each comparison, and the *p* value was adjusted by the Greenhouse-Geisser method when the equality could not be assumed. Any differences following each MAN-OVA were then established using univariate statistics. If a significant main effect was found, a post hoc analysis was conducted using the least significant difference method. The p value for multiple pairwise comparisons was corrected by the Benjamini-Hochberg method,⁵⁴ which controls for type I errors and false discovery rates while having higher statistical power than other options.⁵⁵ Effect sizes were indicated by the Hedges'g for each significant pair⁵⁶ and interpreted using the Cohen convention⁵⁷ of negligible (<0.20), small (0.20-0.49), medium (0.50-0.79), and large (≥ 0.80) values. All statistics were generated using SPSS software (Version 16.0; SPSS Inc., Chicago, IL, USA) at an α level of p < 0.05.

3. Results

All participants successfully completed the half marathon with an average running speed of 11.37 \pm 1.49 km/h. VALR

and VILR were similar among the distance checkpoints (F = 0.623, p = 0.881), while segment CVs gradually increased as mileage accumulated. MANOVA reported significant between-subject effects in the early-stance phase (F = 1.776, p = 0.002). Significant changes in CVs, as indicated by univariate tests, were found in pelvis frontal vs. thigh frontal (F = 2.653, p = 0.008), thigh sagittal vs. shank sagittal (F=2.915, p=0.004), shank sagittal vs. rearfoot sagittal (F = 2.556, p = 0.010), and shank transverse vs. rearfoot frontal (F=6.035, p < 0.001) planes at the early stance (Table 1). After correction of significant level, post hoc tests (Fig. 2A) showed that the CV of pelvis frontal vs. thigh frontal was higher at 20 km than at 8 km (g = 0.59, p = 0.022). Compared to the earlier distance checkpoint (2 km), the CV of shank transverse vs. rearfoot frontal was significantly reduced at 8 km (g = 0.30, p = 0.020) but increased at 18 km (g = 0.05, p< 0.001) and at 20 km (g = 0.36, p < 0.001). The CV of shank transverse vs. rearfoot frontal was also higher at 20 km than at 4 km (g = 0.66, p = 0.025) and at 8 km (g = 0.63, p = 0.036) (Fig. 2E).

4. Discussion

In this study, we quantified differences in segment CVs and vertical loading rate according to mileages during a half marathon to indicate coordination adaptation and its potential injury risk. We examined multiple-plane coordination variabilities with the hypothesis that, under a controlled condition at the runner's preferred speed, the segment CVs and loading rate would increase as the half marathon progressed. In partial support of this hypothesis, the participants generally produced higher segment CVs across all couplings of interest and stance phases at the late stage of prolonged running, while VALR and VILR did not change based on running mileages. CVs of the pelvis frontal vs. femur frontal during the early stance were significantly increased at the 20-km checkpoint, compared to those at the 8-km checkpoint. CVs of shank transverse vs. rearfoot frontal during early stance were higher at the 20-km checkpoint than those at the 2-km, 4-km, and 8-km checkpoints.

Runners in our study had decreases in their CVs (2.75%-19.83%) at the early stages of the running trials (within the first 8 km). Because the participants were asymptomatic at the time of the experiments, we speculated that they adopted a conservative coordination strategy to retain stability and balance^{58,59} in response to a non-habitual running environment.⁴⁴ We may have not assigned the participants sufficient warm-up time; there were no measures or standards of complete adaptations in the study. It is plausible that the reduced CVs from the 2-km to 8-km checkpoints reflected the runners' actual coordination under a research condition, though this possibility did not affect the trend in our results.

Following these early-stage adjustments, the runners regained their CV levels across the couplings of interest and produced larger CVs as they progressed to the end of the half marathon. However, statistical significance existed only in the early stance for coordination on the frontal/transverse planes.

Table 1 Vertical loading	rate and coordin	ation variability o	of segment coupli	ngs of interest acr	oss the distance	checknoints of a	half marathon (n	nean + SD).					Coordi
Mileage	2 km	4 km	6 km	8 km	10 km	12 km	14 km	16 km	18 km	20 km	F	d	nation
Vertical loading 1	ate of ground read	tion force (BW/s)											n va
Average	50.23 ± 19.46	48.45 ± 15.21	52.03 ± 18.66	53.55 ± 17.96	48.76 ± 17.31	52.18 ± 21.60	49.82 ± 16.76	51.06 ± 20.68	49.93 ± 18.84	48.99 ± 19.82	0.892	0.470	aria 090 [.] 0
Instantaneous	75.22 ± 24.79	74.61 ± 21.48	78.61 ± 23.53	79.68 ± 22.37	77.13 ± 23.47	78.58 ± 27.04	76.75 ± 22.32	77.36 ± 22.82	76.56 ± 23.05	74.36 ± 21.66	0.710	0.601	abi 0.048
Pelvis frontal vs.	thigh frontal (°)												lit
Early stance	10.39 ± 5.92	10.90 ± 6.81	10.10 ± 5.28	9.87 ± 5.84	11.66 ± 7.22	11.67 ± 7.38	12.97 ± 7.61	11.59 ± 8.15	13.59 ± 8.91	14.76 ± 9.41	2.653	0.008*	0.159 ii 0.159
Mid stance	3.66 ± 1.14	2.43 ± 1.35	3.18 ± 1.43	3.23 ± 1.15	3.65 ± 2.60	3.41 ± 0.82	3.85 ± 1.79	4.14 ± 2.48	3.99 ± 1.71	4.15 ± 1.38	1.144	0.344	0.076 u u
Terminal stance	11.24 ± 7.87	11.11 ± 7.19	10.66 ± 8.13	11.82 ± 8.71	11.31 ± 7.12	10.80 ± 7.39	11.87 ± 8.94	12.30 ± 9.63	12.27 ± 7.84	13.62 ± 8.77	2.199	0.026	0.136 0.136
Pelvis sagittal vs.	thigh sagittal (°)												fn
Early stance	20.75 ± 8.53	20.82 ± 9.25	18.26 ± 7.50	17.67 ± 6.21	17.93 ± 6.92	18.07 ± 6.40	20.02 ± 6.98	18.65 ± 5.40	19.69 ± 7.60	20.09 ± 6.13	1.216	0.291	nar 080.0
Mid stance	2.91 ± 0.88	2.76 ± 0.91	2.83 ± 1.00	3.08 ± 1.17	3.32 ± 0.97	2.98 ± 0.73	3.20 ± 0.79	3.33 ± 1.06	3.35 ± 1.36	4.04 ± 1.78	1.601	0.122	0.103 0.103
Terminal stance	3.66 ± 1.14	3.43 ± 1.35	3.18 ± 1.43	3.23 ± 1.15	3.65 ± 2.60	3.41 ± 0.82	3.85 ± 1.79	4.14 ± 2.48	3.99 ± 1.71	4.15 ± 1.38	1.725	060.0	0.110 0.110
Thigh sagittal vs.	shank sagittal (°)												ler
Early stance	11.24 ± 7.86	11.11 ± 7.19	10.66 ± 8.13	11.81 ± 8.71	11.30 ± 7.12	10.80 ± 7.39	11.87 ± 8.93	12.30 ± 9.63	12.27 ± 7.84	13.62 ± 8.77	2.915	0.004*	0.172 5
Mid stance	3.23 ± 1.30	2.69 ± 0.61	2.84 ± 0.84	2.88 ± 0.84	2.94 ± 0.84	2.87 ± 0.79	2.93 ± 0.87	2.90 ± 0.95	3.45 ± 1.22	3.45 ± 1.22	1.600	0.122	0.103
Terminal stance	3.43 ± 1.15	3.05 ± 0.84	2.75 ± 0.96	3.02 ± 1.03	2.92 ± 1.14	2.95 ± 0.88	3.16 ± 1.41	3.24 ± 1.32	3.50 ± 1.06	3.75 ± 1.56	2.726	0.006	0.163
Shank sagittal vs.	rearfoot sagittal ((,											
Early stance	21.84 ± 7.33	19.65 ± 8.07	19.27 ± 6.31	19.87 ± 7.05	23.06 ± 6.39	20.92 ± 7.48	22.73 ± 7.11	21.95 ± 7.29	20.88 ± 6.56	24.49 ± 9.77	2.556	0.010*	0.154
Mid stance	3.43 ± 1.15	3.05 ± 0.84	2.75 ± 0.96	3.02 ± 1.03	2.92 ± 1.14	2.95 ± 0.88	3.16 ± 1.41	3.24 ± 1.32	3.50 ± 1.06	3.75 ± 1.56	1.258	0.266	0.082
Terminal stance	1.92 ± 0.55	1.76 ± 0.43	1.75 ± 0.54	1.67 ± 0.36	1.66 ± 0.38	1.70 ± 0.48	1.78 ± 0.40	2.03 ± 1.05	2.05 ± 0.43	2.13 ± 0.80	2.368	0.017	0.145
Shank transverse	vs. rearfoot fronts	u(°)											
Early stance	7.27 ± 2.61	6.37 ± 2.53	7.06 ± 2.72	6.45 ± 2.58	7.45 ± 2.61	7.01 ± 2.65	7.26 ± 1.82	6.97 ± 2.42	7.41 ± 2.35	8.39 ± 3.22	6.035	$< 0.001^{*}$	0.301
Mid stance	5.36 ± 1.73	5.35 ± 2.45	5.40 ± 2.15	6.26 ± 2.64	6.15 ± 2.70	5.33 ± 1.83	6.20 ± 2.09	6.64 ± 3.49	6.40 ± 3.07	7.35 ± 2.93	0.829	0.492	0.056
Terminal stance	23.19 ± 12.34	21.06 ± 10.28	21.01 ± 11.22	21.43 ± 11.05	21.09 ± 10.83	19.97 ± 8.76	21.44 ± 10.74	22.08 ± 12.87	23.06 ± 10.50	23.12 ± 12.16	1.309	0.278	0.086
* Significant di	fference reported	in univariate test	ts (p values are in	boldface type) fo	llowing significa	int MANOVA re	sults.						

Our results were in line with those of other studies using discrete kinematic data, which showed changed angulation between adjacent lower-limb segments in marathon finishers.^{12,16} Nevertheless, many studies^{18,26,27} reporting segment coordination found small to no effects of running time or mileage on CVs for healthy runners. Maclean et al.²⁶ even reported slightly reduced CVs in a healthy control group during a 30-min run. It is important to note that the participants in these studies finished an apparently shorter distance (approximately 4-5 km) than those in the present study (more than 21 km). Limited running distance may not expose runners to adequate fatigue effects that would trigger detectable changes in coordination patterns for experienced marathoners.⁶⁰

During the early-stance phase of running gait, segment coordination on the sagittal plane is the major component of the lower-limb kinematic chain for loading response.⁶¹ Following the heel strike, adjacent segments of the lower limb usually rotate in opposite directions 62,63 to yield negative work for cushioning the landing. An increased CV is thought to indicate changes in this coordinative state and affect impact attenuation.²⁸ Our participants did not produce significant alterations in CVs on the sagittal plane during the half marathon; this corresponded with the unchanged vertical loading rate. CVs on the frontal/transverse plane also aid in force absorption at the early stance, though its increments in our study seemed to have limited effects on the loading results. Segment coordination on the sagittal plane is regulated by powerful muscle groups, such as the gluteus, quadriceps, and calf muscles.⁶⁴ In contrast, muscles driving coordination on the frontal/transverse planes are smaller in volume⁶⁴ and are more prone to exertion when resisting external loading. This subsequently interrupts neuromuscular recruitment in the lower limb and introduces variances in coordination patterns. Based on our results, we speculate that the increased frontal/ transverse-plane CVs may be the precursor sign of fatigue. The loss of coordination stability could extend to the sagittal plane if the participants were given more running time. Runners undertaking a full marathon or longer run may produce more outcomes that can verify this conjecture.

Although largely speculative, it appears that there is a "normal" range of segment CV in healthy individuals,²¹ which leads to optimal load distribution within the lower limb.⁶⁵ Going beyond this range at early stance could not only cause the ground force to be applied in extreme magnitude²¹ but also could poorly engage the soft tissues.⁴⁴ Increased CVs on the frontal/transverse plane may not be problematic because running is not heavily reliant on lateral/spinning coordination stability. However, if increased CVs on the sagittal plane occur as anticipated, they would induce unaccustomed strains on the internal structures, such as the meniscus and anterior cruciate ligament,²¹ and exacerbate the injury risk.

Abbreviations: BW = body weight; ES = effect size; MANOVA = multivariate analyses of variances.

Running involves a linear and cyclic movement that favors a stable coordination pattern,^{21,44,66} which, in turn, challenges runners' muscular endurance. Training programs for endurance runners should focus on improving the sustainability of their neuromuscular control over a "normal" level of segment CVs. Running-specific strengthening programs are



Fig. 2. Changes in coordination variabilities of interest (early stance phase) over the course of the half marathon: (A) pelvis frontal vs. thigh frontal; (B) pelvis sagittal vs. thigh sagittal (C) thigh sagittal vs. shank sagittal; (D) shank sagittal vs. rearfoot sagittal; (E) shank transverse vs. rearfoot frontal. * p < 0.05, ** p < 0.01 in post hoc pairwise comparison. Scale of the y-coordinate: 1 unit = 0.5° .

increasingly used to supplement regular running regimens for an injury-free running career.⁶⁷ In light of the results of our study, this training strategy could be particularly helpful for the hip and ankle joints, which possess more mobility on the frontal/transverse plane.⁶⁸ Although a longitudinal study is needed in the future to understand how interventions in training affect the running coordination and injury profile, coaches and sports professionals could use the information given in our study to develop optimal training regimes for individual runners.

Our study had a few limitations. First, the data of the running trials were collected on a treadmill, and the calculated CVs could differ from those of over-ground running.⁶⁹ We considered this limitation negligible because changes in the variables were attributed mainly to increasing running mileages on the same baseline condition. Furthermore, because the running speed was controlled in our research protocol, it could have influenced the coordination profile.⁷⁰ Runners are known to change their running speeds at various stages of an actual marathon or half marathon race.^{71,72} It is unclear how the varying spacing strategy affected the runners' coordination patterns; this warrants further investigation. Future studies should apply segmental inertia sensors instead of a laboratory-based motion-capture system to assess running movements across different mileages in actual half marathon races. Second, the effect sizes reported in our study were medium to low and may have resulted primarily from the large individual variances in running coordination. This issue may be better addressed by a larger scale study targeting a runner sub-group. Our study reported CV values that generally agreed with those in the literature.^{18,26,44} However, CV-quantifying methods could be slightly modified from time to time to make the outcomes less comparable among various studies. Third, we tended to link our CV results with running-related injuries.

However, it is unknown whether the increased CVs reached a pathological threshold.⁶⁵ Given the cross-sectional design of our study, it was impossible to determine the injury risk using the current research protocol.

5. Conclusion

Runners who finished a half marathon showed increased segment CV of the lower limb. These changes appeared to have greater impacts on the pelvis-thigh and shank-rearfoot couplings during the early stance phase. Vertical loading rates to the lower limb were not affected by running mileages. The results from our study suggest that, at the early stance, the runners' coordination patterns were stable on the sagittal plane but more variable on the frontal/transverse plane as the half marathon progressed, which may be related to fatigue onset. Additional studies should be carried out in this regard before our conclusions can be confirmed.

Acknowledgments

This work was supported by General Research Funds granted by the Hong Kong Research Grant Council (#PolyU152065/17E) and by funding from the National Natural Science Foundation of China (#11732015).

Authors' contributions

TLC participated in the research design, data collection, data analysis, result interpretation, initial manuscript draft, revision, and submission preparation; DWW conducted data analysis and manuscript revision; YW and CT conducted data collection and manuscript revision; WKL and MZ conceived the study, supervised data collection, managed research activities, and revised the manuscript. All authors have read and approved the final version of the manuscript, and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

References

- 1. Andersen JJ. *The state of running 2019. Athletic shoe reviews*. Available at: https://runrepeat.com/state-of-running. [accessed 03.03.2020].
- Andersen JJ. Marathon statistics 2019 worldwide (research). Athletic shoe reviews. Available at: https://runrepeat.com/research-marathon-per formance-across-nations. [accessed 03.03.2020].
- Knechtle B, Nikolaidis PT, Zingg MA, Rosemann T, Rüst CA. Half-marathoners are younger and slower than marathoners. *SpringerPlus* 2016;5:76. doi:10.1186/s40064-016-1704-9.
- Lee DC, Brellenthin AG, Thompson PD, Sui X, Lee IM, Lavie CJ. Running as a key lifestyle medicine for longevity. *Prog Cardiovasc Dis* 2017;60:45–55.
- Nigg BM. The role of impact forces and foot pronation: A new paradigm. *Clin J Sport Med* 2001;11:2–9.
- Benson LC, O'Connor KM. The effect of exertion on joint kinematics and kinetics during running using a waveform analysis approach. *J Appl Biomech* 2015;31:250–7.

- Hardin EC, van den Bogert AJ, Hamill J. Kinematic adaptations during running: effects of footwear, surface, and duration. *Med Sci Sports Exerc* 2004;36:838–44.
- Mizrahi J, Verbitsky O, Isakov E, Daily D. Effect of fatigue on leg kinematics and impact acceleration in long distance running. *Hum Mov Sci* 2000;19:139–51.
- **9.** Benson L. *The effect of exertion on intra-limb joint coordination variability during running using a waveform analysis approach*. Milwaukee, WI: University of Wisconsin-Milwaukee; 2013. [Master's thesis].
- Bartlett R. Introduction to sports biomechanics: Analysing human movement patterns. Abingdon, UK: Routledge; 2014.p.360.
- Maas E, De Bie J, Vanfleteren R, Hoogkamer W, Vanwanseele B. Novice runners show greater changes in kinematics with fatigue compared with competitive runners. *Sports Biomech* 2018;17:350–60.
- Reenalda J, Maartens E, Homan L, Buurke JHJ. Continuous three dimensional analysis of running mechanics during a marathon by means of inertial magnetic measurement units to objectify changes in running mechanics. *J Biomech* 2016;49:3362–7.
- Dierks TA, Manal KT, Hamill J, Davis I. Lower extremity kinematics in runners with patellofemoral pain during a prolonged run. *Med Sci Sports Exerc* 2011;43:693–700.
- 14. Derrick TR, Dereu D, McLean SP. Impacts and kinematic adjustments during an exhaustive run. *Med Sci Sports Exerc* 2002;**34**:998–1002.
- Dierks TA, Davis IS, Hamill J. The effects of running in an exerted state on lower extremity kinematics and joint timing. *J Biomech* 2010;43:2993–8.
- Chan-Roper M, Hunter I, Myrer J, Eggett D, Seeley M. Kinematic changes during a marathon for fast and slow runners. J Sports Sci Med 2012;11:77–82.
- Wilson C, Simpson SE, van Emmerik REA, Hamill J. Coordination variability and skill development in expert triple jumpers. Sports Biomech 2008;7:2–9.
- Hafer JF, Brown AM, Boyer KA. Exertion and pain do not alter coordination variability in runners with iliotibial band syndrome. *Clin Biomech* (*Bristol, Avon*) 2017;47:73–8.
- Ferber R, Pohl MB. Changes in joint coupling and variability during walking following tibialis posterior muscle fatigue. *J Foot Ankle Res* 2011;4:6. doi:10.1186/1757-1146-4-6.
- 20. Trezise J, Bartlett R, Bussey M. Coordination variability changes with fatigue in sprinters. *Int J Sports Sci Coach* 2011;6:357–63.
- Baida SR, Gore SJ, Franklyn-Miller AD, Moran KA. Does the amount of lower extremity movement variability differ between injured and uninjured populations? A systematic review. *Scand J Med Sci Sports* 2018;28:1320–38.
- Pollard CD, Stearns KM, Hayes AT, Heiderscheit BC. Altered lower extremity movement variability in female soccer players during side-step cutting after anterior cruciate ligament reconstruction. *Am J Sports Med* 2015;43:460–5.
- Gribbin TC, Slater LV, Herb CC, et al. Differences in hip-knee joint coupling during gait after anterior cruciate ligament reconstruction. *Clin Biomech (Bristol, Avon)* 2016;32:64–71.
- 24. Herb CC, Chinn L, Dicharry J, McKeon PO, Hart JM, Hertel J. Shankrearfoot joint coupling with chronic ankle instability. *J Appl Biomech* 2014;**30**:366–72.
- Cunningham TJ, Mullineaux DR, Noehren B, Shapiro R, Uhl TL. Coupling angle variability in healthy and patellofemoral pain runners. *Clin Biomech (Bristol, Avon)* 2014;29:317–22.
- MacLean CL, van Emmerik R, Hamill J. Influence of custom foot orthotic intervention on lower extremity intralimb coupling during a 30-minute run. J Appl Biomech 2010;26:390–9.
- Miller RH, Meardon SA, Derrick TR, Gillette JC. Continuous relative phase variability during an exhaustive run in runners with a history of iliotibial band syndrome. *J Appl Biomech* 2008;24:262–70.
- Heiderscheit BC, Hamill J, van Emmerik REA. Variability of stride characteristics and joint coordination among individuals with unilateral patellofemoral pain. J Appl Biomech 2002;18:110–21.
- Tsao H, Galea MP, Hodges PW. Reorganization of the motor cortex is associated with postural control deficits in recurrent low back pain. *Brain* 2008;131:2161–71.
- **30.** Hodges PW, Tucker K. Moving differently in pain: A new theory to explain the adaptation to pain. *Pain* 2011;**152**(Suppl. 3):S90–8.

- **31.** Schmidt RA. Motor schema theory after 27 years: Reflections and implications for a new theory. *Res Q Exerc Sport* 2003;74:366–75.
- Saugy J, Place N, Millet GY, Degache F, Schena F, Millet GP. Alterations of neuromuscular function after the world's most challenging mountain ultramarathon. *PLoS One* 2013;8:e65596. doi:10.1371/journal.pone.0065596.
- Paavolainen L, Nummela A, Rusko H, Häkkinen K. Neuromuscular characteristics and fatigue during 10 km running. *Int J Sports Med* 1999;20:516–21.
- Ross EZ, Goodall S, Stevens A, Harris I. Time course of neuromuscular changes during running in well-trained subjects. *Med Sci Sports Exerc* 2010;42:1184–90.
- García-Manso JM, Rodríguez-Ruiz D, Rodríguez-Matoso D, de Saa Y, Sarmiento S, Quiroga M. Assessment of muscle fatigue after an ultra-endurance triathlon using tensiomyography (TMG). J Sports Sci 2011;29:619–25.
- **36.** Mizrahi J, Verbitsky O, Isakov E. Fatigue-related loading imbalance on the shank in running: A possible factor in stress fractures. *Ann Biomed Eng* 2000;**28**:463–9.
- Voloshin AS, Mizrahi J, Verbitsky O, Isakov E. Dynamic loading on the human musculoskeletal system—effect of fatigue. *Clin Biomech (Bristol, Avon)* 1998;13:515–20.
- Needham R, Naemi R, Chockalingam N. Quantifying lumbar-pelvis coordination during gait using a modified vector coding technique. *J Biomech* 2014;47:1020-6.
- **39.** Chen TL, An WW, Chan ZYS, Au IPH, Zhang ZH, Cheung RTH. Immediate effects of modified landing pattern on a probabilistic tibial stress fracture model in runners. *Clin Biomech (Bristol, Avon)* 2016;**33**:49–54.
- 40. Yamato TP, Saragiotto BT, Lopes AD. A consensus definition of runningrelated injury in recreational runners: A modified Delphi approach. *J Orthop Sports Phys Ther* 2015;45:375–80.
- Orendurff MS, Kobayashi T, Tulchin-Francis K, et al. A little bit faster: Lower extremity joint kinematics and kinetics as recreational runners achieve faster speeds. *J Biomech* 2018;71:167–75.
- John CT, Anderson FC, Higginson JS, Delp SL. Stabilisation of walking by intrinsic muscle properties revealed in a three-dimensional muscledriven simulation. *Comput Meth Biomech Biomed Engin* 2013;16:451–62.
- Altman AR, Davis IS. A kinematic method for footstrike pattern detection in barefoot and shod runners. *Gait Posture* 2012;35:298–300.
- Hafer JF, Peacock J, Zernicke RF, Agresta CE. Segment coordination variability differs by years of running experience. *Med Sci Sports Exerc* 2019;51:1438–43.
- 45. Schellenberg F, Taylor WR, Jonkers I, Lorenzetti S. Robustness of kinematic weighting and scaling concepts for musculoskeletal simulation. *Comput Methods Biomech Biomed Engin* 2017;20:720–9.
- 46. Delp SL, Loan JP, Hoy MG, Zajac FE, Topp EL, Rosen JM. An interactive graphics-based model of the lower extremity to study orthopaedic surgical procedures. *IEEE Trans Biomed Eng* 1990;37:757–67.
- Chang R, Van Emmerik R, Hamill J. Quantifying rearfoot-forefoot coordination in human walking. J Biomech 2008;41:3101–5.
- Hafer JF, Boyer KA. Variability of segment coordination using a vector coding technique: reliability analysis for treadmill walking and running. *Gait Posture* 2017;51:222–7.
- 49. Foch E, Reinbolt JA, Zhang S, Fitzhugh EC, Milner CE. Associations between iliotibial band injury status and running biomechanics in women. *Gait Posture* 2015;41:706–10.
- Rodrigues P, TenBroek T, Hamill J. Runners with anterior knee pain use a greater percentage of their available pronation range of motion. *J Appl Biomech* 2013;29:141–6.

- Batschelet E. Circular statistics in biology. London: Academic Press; 1981. p.371.
- 52. Novacheck TF. The biomechanics of running. Gait Posture 1998;7:77-95.
- Bland JM, Altman DG. Statistics notes. Logarithms. *BMJ* 1996;**312**:700. doi:10.1136/bmj.312.7032.700.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. J R Stat Soc Series B Stat Methodol 1995;57:289–300.
- Williams VSL, Jones LV, Tukey JW. Controlling error in multiple comparisons, with examples from state-to-state differences in educational achievement. J Educ Behav Stat 1999;24:42–69.
- Hedges LV. Distribution theory for Glass's estimator of effect size and related estimators. *J Educ Stat* 1981;6:107–28.
- Cohen J. Statistical power analysis for the behavioral sciences. Hillsdale, NJ: Routledge; 1988.
- Wong DW, Lam WK, Lee WC. Gait asymmetry and variability in older adults during long-distance walking: implications for gait instability. *Clin Biomech (Bristol, Avon)* 2019;72:37–43.
- Wong DW, Lam WK, Yeung LF, Lee WC. Does long-distance walking improve or deteriorate walking stability of transtibial amputees? *Clin Biomech (Bristol, Avon)* 2015;30:867–73.
- Glover B, Glover SF. The competitive runner's handbook: The bestselling guide to running 5Ks through marathons. London: Penguin Books; 1999.
- **61.** Svoboda Z, Janura M, Kutilek P, Janurova E. Relationships between movements of the lower limb joints and the pelvis in open and closed kinematic chains during a gait cycle. *J Hum Kinet* 2016;**51**: 37–43.
- Hafer JF, Boyer KA. Age related differences in segment coordination and its variability during gait. *Gait Posture* 2018;62:92–8.
- Hafer JF, Freedman Silvernail J, Hillstrom HJ, Boyer KA. Changes in coordination and its variability with an increase in running cadence. *J Sports Sci* 2016;34:1388–95.
- 64. Vishram S. *Textbook of anatomy: Abdomen and lower limb.* Vol II, 2nd ed. Chennai: Elsevier; 2014.
- Hamill J, Palmer C, Van Emmerik RE. Coordinative variability and overuse injury. *Sports Med Arthrosc Rehabil Ther Technol* 2012;4:45. doi:10.1186/1758-2555-4-45.
- Erdmann WS, Lipinska P. Kinematics of marathon running tactics. *Hum Mov Sci* 2013;6:1379–92.
- Middelkoop MV, Kolkman J, Ochten JV, Bierma-Zeinstra SMA, Koes B. Prevalence and incidence of lower extremity injuries in male marathon runners. *Scand J Med Sci Sports* 2008;18:140–4.
- Medina McKeon JM, Hoch MC. The ankle-joint complex: A kinesiologic approach to lateral ankle sprains. J Athl Train 2019;54:589–602.
- Wheat JS, Baltzopoulos V, Milner CE, Bartlett RM, Tsaopoulos D. Coordination variability during overground, treadmill and treadmill-ondemand running. XXIII ISBS Annual Conference. Beijing. 2005;7:81–4.
- Bailey JP, Freedman Silvernail J, Dufek JS, Navalta J, Mercer JA. Effects of treadmill running velocity on lower extremity coordination variability in healthy runners. *Hum Mov Sci* 2018;61:144–50.
- Renfree A, St Clair Gibson A. Influence of different performance levels on pacing strategy during the Women's World Championship marathon race. *Int J Sports Physiol Perform* 2013;8:279–85.
- 72. Haney Jr. TA, Mercer JA. A description of variability of pacing in marathon distance running. *Int J Exerc Sci* 2011;4:133–40.