

DOI: 10.1002/adsc.201((will be filled in by the editorial staff))

Rhodium-Catalyzed Oxidative Cycloaddition of *N*-tert-Butoxycarbonyl Hydrazones with Alkynes for the Synthesis of Functionalized Pyrroles via C(sp³)-H Bond Functionalization

Chun-Ming Chan, Zhongyuan Zhou and Wing-Yiu Yu*

State Key Laboratory for Chirosciences and Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong
 Fax : (+852) 2364 9932; email: wing-yiu.yu@polyu.edu.hk

Received: ((will be filled in by the editorial staff))



Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/adsc.201#####>. ((Please delete if not appropriate))

Abstract. A rhodium(III)-catalyzed cycloaddition of *N*-tert-butoxycarbonyl (*N*-Boc) hydrazones with internal alkynes was developed. The reaction features a regioselective α -imino alkyl C(sp³)-H bond functionalization resulting in selective formation of highly functionalized *NH*-free pyrroles. Our studies showed that utilizing the *N*-tert-butoxycarbonyl (*N*-Boc) as the oxidizing directing group is critical for achieving the observed pyrrole formation versus the isoquinoline formation. To account for the pyrrole formation, we hypothesi-

zed a prior tautomerization of the *N*-Boc hydrazones to enamines should occur, followed by regioselective C(sp²)-H cleavage to form a putative five-membered rhodacycle. Subsequent coupling of the rhodacycle with the alkynes should afford the pyrrole products.

Keywords: cycloaddition; C-H activation; nitrogen heterocycles; alkynes; rhodium

Introduction



Transition metal-catalyzed oxidative C-H bond cycloaddition reactions with alkynes^[1,2] and carbenoids^[3] are attracting current attention for developing atom-economical synthesis of medicinally valuable heterocycles. The versatility of the transition metal-catalysis approach is exemplified by the recent advance in indoles and isoquinolines synthesis involving the Cp*Rh(III)-catalyzed oxidative C(sp²)-H cycloaddition with *N*-oxyenamine as substrates.^[1b,e] With the N-O moieties as oxidizing directing groups, the cycloadditions with alkynes and carbenoids can operate under mild conditions without external oxidants and exhibit high regio- and chemoselectivity.^[1-3]

Pyrroles are privileged heterocyclic scaffolds present in many pharmaceutical products and functional materials.^[4] Analogous to the indoles and isoquinolines synthesis, metal-catalyzed oxidative C-H bond cycloaddition for pyrroles synthesis has also received considerable interest.^[5] In 2010, Fagnou^[6a] and Glorius^[6b] first reported independently the *N*-acetyl pyrrole synthesis by [Cp*RhCl₂]₂-catalyzed C(sp²)-H bond cycloaddition of *N*-acetyl enamides with alkynes (Scheme 1a). Later, the research groups of Wang,^[7a] Ackermann^[7b] and Liu^[7c] successfully developed the analogous cycloaddition reactions based on ruthenium catalysis. For other transition metal approaches, Loh's^[8a] and Guan's^[8b] groups reported the Pd-catalyzed pyrrole synthesis, whereas

Examples involving sp² C-H cycloaddition



Examples involving sp³ C-H cycloaddition



Scheme 1. Selected examples for pyrroles synthesis by metal-catalyzed cycloaddition.

Zhang and co-workers also developed the related cobalt-catalyzed cycloaddition reaction.^[9]

Currently, extensive effort is directed to developing Rh-catalyzed regioselective C(sp³)-H bond functionalization for C-C and C-X (X = N, B, Si) bonds formation.^[10] However, examples for pyrrole

synthesis by catalytic regioselective C(sp³)-H bond functionalization are sparse in the literature.^[11] In 2010, Glorius and co-workers pioneered the Cp*Rh(III)-catalyzed oxidative cycloaddition of enamides with alkynes to form pyrroles via C(sp³)-H bond activation, and Cu(II) salt was employed as co-oxidant (Scheme 1b).^[6b] In 2014, Zeng and co-workers reported the Pd-catalyzed C(sp³)-H bond oxidative cycloaddition of *N*-(2-pyridyl) ketoimines with internal alkynes to form *N*-(2-pyridyl)pyrroles.^[11b] Later in 2015, they also developed the Rh-catalyzed cyclization of α -imino C(sp³)-H bonds with α -acyl diazocarbonyl compounds to form the corresponding pyrroles (Scheme 1c).^[11a] In this report, we disclose the Cp*Rh(III)-catalyzed oxidative cycloaddition of *N*-Boc hydrazones with internal alkynes to afford *NH*-free pyrroles via regioselective α -imino C(sp³)-H bond cycloaddition. This reaction features the N-N bond cleavage for effective product formation and catalyst turnovers without the need for external oxidants.


Results and Discussion

To begin, we reacted 4-methyl-acetophenone *N*-Boc hydrazone (**1a**, 0.4 mmol) and diphenylacetylene (**2a**, 0.2 mmol) with [Cp*RhCl₂]₂ (2.5 mol %), CsOAc (25 mol %) and AcOH (3.0 equiv) in MeCN (2.5 mL) at 120 °C for 16 h, and pyrrole **3aa** was obtained in 68% yield. We were gratified that isoquinoline **4aa** was formed in <10% yield (Supporting information, Table S3).^[12] However, when the same protocol was applied to 4-bromo-acetophenone *N*-Boc hydrazone **1f**, no pyrrole or isoquinoline were obtained. After several trials, we found that doubling the solvent volume to 5 mL led to effective formation of pyrrole **3fa** in 60% yield (Supporting information, S6).^[12]

To further optimize the reaction, we first examined the reaction temperature. Apparently, performing the reaction at lower temperature (e.g. 60 °C) led to **3aa** formation in trace quantity (Table 1, *c.f.* entry 1 and 2). The presence of CsOAc, AcOH and [Cp*RhCl₂]₂ is essential, and absence of any one component resulted in ineffective transformation (entries 3-5). Apart from AcOH, pivalic acid and benzoic acid are equally effective for promoting the pyrrole formation (entries 6-7). By examining several acid-base combinations, employing Na₂CO₃ (25 mol %) instead of CsOAc was found to give **3aa** in 90% yield (entry 9). Common organic solvents other than MeCN gave slightly lower product yields (entries 10-13).

It is known that employing oxidizing directing groups is a favourable approach for designing transition metal-catalyzed C-H bond cycloaddition reactions.^[13] As shown by Hartwig,^[14a] Fagnou^[2r] and Glorius,^[14b] the catalytic cross coupling reactions of oximes and benzohydroxamic acids with alkynes afforded indoles and isoquinolones respectively in excellent selectivity. It is generally accepted that cleavage of the weak N-O σ -bond facilitates the product formation (*presumably via a sequence of*

Table 1. Reaction optimization.^[a]





Entry	Solvent (5.0 mL)	Acid (3.0 equiv)	Base (25 mol %)	Yield (%) ^[b]
1	MeCN	AcOH	CsOAc	70
2 ^[c]	MeCN	AcOH	CsOAc	<5
3	MeCN	AcOH	-	0
4	MeCN	-	CsOAc	0
5 ^[d]	MeCN	AcOH	CsOAc	0
6	MeCN	PivOH	CsOAc	89
7	MeCN	PhCO ₂ H	CsOAc	88
8	MeCN	AcOH	NaOAc	80
9	MeCN	AcOH	Na₂CO₃	90
10	dioxane	AcOH	Na ₂ CO ₃	40
11	toluene	AcOH	Na ₂ CO ₃	77
12	DCE	AcOH	Na ₂ CO ₃	42
13	THF	AcOH	Na ₂ CO ₃	78

^[a] Reaction conditions: **1a** (0.4 mmol), **2a** (0.2 mmol), and [Cp*RhCl₂]₂ (2.5 mol %) at 120 °C for 16 h. ^[b] NMR yields.

^[c] Reaction was performed at 60 °C. ^[d] Reaction was performed without [Cp*RhCl₂]₂.

Table 2. Effect of leaving groups (LG).^[a,b]




Entry	Leaving groups (LG)	Yield (%) ^[b] 3aa:4aa
1		90:<5
2		0:75
3		0:37
4		0:0
5		0:0

^[a] Reaction conditions: **1** (0.4 mmol), **2a** (0.2 mmol), [Cp*RhCl₂]₂ (2.5 mol %), Na₂CO₃ (25 mol %) and AcOH (3.0 equiv) in MeCN (5.0 mL) at 120 °C for 16 h. ^[b] Isolated yields.

oxidative addition and reductive elimination) and catalyst turnovers. Recently, Glorius^[2c] and co-workers reported a catalytic oxidative C(sp²)-H bond cycloaddition of some Boc-protected hydrazines with alkynes to form indoles, and the N-N bond was found to be an effective oxidizing directing group. In this work, we have examined a series of hydrazones bearing different terminal leaving groups for the catalytic cycloaddition with alkynes (Table 2). With *N*-Boc as a leaving group, facile cycloaddition of 4-methyl-acetophenone *N*-Boc hydrazone **1a** with diphenylacetylene **2a** furnished pyrrole **3aa** in 90%

Table 3. Scope of *N*-Boc hydrazones.^[a,b]



^[a] Reaction conditions: **1** (0.4 mmol) and **2a** (0.2 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (2.5 mol %), Na_2CO_3 (25 mol %) and AcOH (3.0 equiv) in MeCN (5.0 mL) at 120 °C for 16 h. ^[b] Isolated yields. ^[c] Reaction conditions: **1s** (0.8 mmol) and **2a** (0.2 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (2.5 mol %), Na_2CO_3 (25 mol %) and AcOH (3.0 equiv) in MeCN (5.0 mL) at 120 °C for 2 h.





Figure 1. Molecular structure of 5-(4-(methylthio)phenyl)-2,3-diphenyl-1 *H*-pyrrole (**3ea**).

yield with isoquinolines **4aa** being formed in <5% yield. Importantly, the analogous reactions with 4-methyl-acetophenone hydrazone bearing NHAc and OMe leaving groups produced isoquinoline **4aa** exclusively without any pyrrole formation. Hydrazones with picolinamide and NH₂ leaving group are ineffective substrates; neither pyrroles nor isoquinolines were obtained.

With these optimized conditions in hand, reactions of various para-substituted acetophenone *N*-Boc hydrazones with diphenylacetylene **2a** were examined, and the corresponding pyrroles **3aa-3ga** were obtained in 62-90% yields (Table 3). The molecular structure of the pyrrole product **3ea** has been confirmed by X-ray crystallography (Figure 1).^[15] In these cases,

Table 4. Scope of alkynes.^[a,b]



^[a] Reaction conditions: **1a** (0.4 mmol) and **2** (0.2 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (2.5 mol %), Na_2CO_3 (25 mol %) and AcOH (3.0 equiv) in MeCN (5.0 mL) at 120 °C for 16 h. ^[b] Isolated yields. ^[c] NMR yield, isoquinoline **4af**: 60% yield.

isoquinoline formation resulted from the ortho C(sp²)-H bond activation was insignificant (Supporting Information, Table S1).^[12] For acetophenone *N*-Boc hydrazones bearing substituents located at other positions of the aryl ring, successful cycloadditions were also achieved to give pyrroles **3ha-3oa** in 42-73% yields. As expected, the cycloaddition of a naphthyl-substituted acetophenone *N*-Boc hydrazone furnished pyrrole **3pa** in 72% yield.

For the Rh(III)-catalyzed C(sp³)-H bond functionalization, examples concerning the functionalization of primary C(sp³)-H bonds are well documented.^[10] However, functionalizations of the analogous secondary C(sp³)-H bond are less common.^[16] In this work, when an ethylphenyl ketone-derived *N*-Boc hydrazone was subjected to the Rh-catalyzed cycloaddition reaction with diphenylacetylene, effective functionalization of the secondary α -imino C(sp³)-H bond was achieved to give pyrrole **3qa** in 75% yield. Nonetheless, the analogous reaction with the substrate bearing an *n*-propyl imino substituent was sluggish; isoquinoline **4va** was obtained in 23% yield, with the desired pyrrole **3va** being formed in <5% yield.^[12] *N*-Boc hydrazone containing a seven-membered carbocyclic ketone moiety was found to undergo effective cycloaddition to produce the tricyclic pyrrole **3ra** in 42% yield with <10% isoquinoline formation. Yet, the analogous cycloaddition of the *N*-Boc hydrazone containing a six-membered carbocyclic ketone afforded isoquinoline **4ua** in 90% yield.^[12] In this work, when a *N*-Boc hydrazone-derived from acetone was employed as substrate, effective C(sp³)-H bond cycloaddition was achieved to give the α -methylpyrrole **3sa** in 32% yield.


With 4-methyl-acetophenone *N*-Boc hydrazone **1a** as substrate, the alkyne scope has also been examined (Table 4). The reaction with 1-phenyl-1-propyne **2b** under the Rh-catalyzed conditions furnished pyrrole **3ab** as a single regioisomer. Similar results were obtained for the analogous reactions with 1-phenyl-1-butyne **2c**, 1-phenyl-1-pentyne **2d** and 1-phenyl-1-

hexyne **2e**. When 4-octyne **2f** was employed as the coupling partner, isoquinoline **4af** became a major product (60%), and the desired pyrrole **3af** was formed in <20% yield. This finding implies that the transition state for successful pyrrole formation should be rather sensitive to steric factors. Having such hypothesis in mind, we anticipated that employing a bulkier substituent such as trimethylsilyl (TMS) group on the alkyne would hinder the pyrrole formation. Interestingly, the reaction of phenyl silylacetylene **2g** with **1a** gave the desilylated 2,5-diarylpyrrole **3ag** in 55% yield with a minor isoquinoline formation (<5%). Effective cycloadditions were also achieved with some substituted diphenylacetylenes, and pyrroles **3ah-3aj** were obtained in 53-75% yields.


The catalytic pyrrole formation developed in this work features the regioselective α -imino $C(sp^3)$ -H bond functionalization versus the aryl $C(sp^2)$ -H bond functionalization (isoquinoline formation). As shown earlier, replacing the *N*-Boc group of the substrate with an NHAc or OMe group resulted in exclusive isoquinoline formation via the aryl $C(sp^2)$ -H bond functionalization. Presumably, successful pyrrole formation is associated with the *N*-Boc group to direct the $C(sp^3)$ -H bond functionalization by the Rh(III) complex. To understand the observed selectivity for the α -imino $C(sp^3)$ -H bond functionalization, the cycloaddition of the deuterated *N*-Boc hydrazone **1t** with diphenylacetylene was performed under our optimized conditions. The expected **3ta** was obtained in 90% yield *without any loss of deuterium* on the phenyl ring (Scheme 2a). This result implies that the *N*-Boc group should play a key role in directing the kinetically competitive $C(sp^3)$ -H bond functionalization.

Moreover, subjecting 4-methyl-acetophenone *N*-Boc hydrazone **1a** to the Rh(III)/AcOD/CD₃CN system at 120 °C *in the absence of diphenylacetylene* for 20 min, 50% deuterium incorporation was detected at the α -imino methyl and the *N*-Boc groups (Scheme 2b). This result may suggest that the imine-enamine tautomerization could have been promoted by the Rh(III) catalyst coordinated to the *N*-Boc group.

A plausible reaction mechanism is depicted in Scheme 3. The reaction should be initiated by the coordination of the enamine isomer of the *N*-Boc hydrazone **1** to the Rh(III) centre,^[17] followed by the cleavage of the $C(sp^2)$ -H bond to form rhodacycle **A**. The complex **A** is presumably stabilized by the Boc group.^[18] After alkyne insertion, the 7-membered intermediate **B** would undergo the *N*-ligand shift to give the 6-membered rhodacycle **C**.^[19] Reductive elimination via oxidative addition of the *N*-*N* bond to the Rh(III) should produce species **D**. Protonation by acetic acid should furnish the desired pyrrole product. An alternative route involving a Rh(V) nitrene intermediate **E** by acylamino migration is also possible.^[20] Subsequent carbon-migration to the nitrene ligand on species **E** should produce species **D**.



Scheme 2. H/D exchange experiments.



Scheme 3. Plausible reaction mechanism.

Conclusion

In summary, we have developed a Rh(III)-catalyzed oxidative cycloaddition of *N*-Boc hydrazones with internal alkynes. Our study shows that the *N*-Boc group is essential for the selective pyrrole formation. The reaction provides a direct route to highly functionalized pyrroles via direct $C(sp^3)$ -H bond functionalization. This reaction offers an expedient access to *NH*-free pyrroles, which are key structural scaffolds of many pharmaceutical products and functional materials.

Experimental Section

General: All the solvents and reagents were obtained from commercial sources and used without purification unless stated otherwise. All glasswares were dried overnight at 150 °C prior to use. All the Rh-catalyzed reactions were performed without specific protection from air or moisture. Thin layer chromatography was performed on silica gel plates. Flash column chromatography was performed on a silica gel (Merck, 230-400 mesh) column. ¹H and ¹³C NMR spectra were recorded on a Brüker DPX-400 MHz spectrometer. The chemical shift (δ) values are

given in ppm and are referenced to residual solvent peaks; carbon multiplicities were determined by DEPT-135 experiments. Coupling constants (J) were reported in hertz (Hz). The NMR yield values were determined with dibromomethane (6.98 μL) as internal standard, which shows a singlet signal at δ_{H} 4.9 ppm in CD_2Cl_2 . Mass spectra and high resolution mass spectra (HRMS) were obtained on a VG MICROMASS Fison VG platform, a Finnigan Model Mat 95 ST instrument, or a Brüker APEX 47e FT-ICR mass spectrometer. X-ray crystallographic study was performed by a Brüker CCD area detector diffractometer.

General Procedure for Rh(III)-Catalyzed Cycloaddition of *N*-Boc hydrazones with Alkynes for the Synthesis of Pyrrole 3

A 10 mL-sealed tube equipped with a magnetic stirrer was charged with $[\text{Rh}(\text{Cp}^*)\text{Cl}_2]_2$ (3.2 mg, 2.5 mol %), Na_2CO_3 (5.3 mg, 25 mol %), *N*-Boc hydrazone **1** (0.4 mmol) and MeCN (5.0 mL). Alkyne **2** (0.2 mmol) and AcOH (34.3 μL , 0.6 mmol) were added sequentially. The mixture was then stirred and heated at 120 °C for 16 h. After cooling to room temperature, the crude mixture was filtered through Celite® and concentrated under reduced pressure. The residue was then purified by flash column chromatography (hexanes / ethyl acetate, 9:1, v / v) to give the desired product **3**.

3ea was isolated as a pale yellow solid (50 mg, 73% yield), mp 185 – 186 °C. (R_f = 0.6, hexanes / ethyl acetate 85:15, v / v). The solid was redissolved in a minimum amount of DCM; diffusion with hexane gave a pale yellow solid. ^1H NMR (400 MHz, CD_2Cl_2) δ 2.55 (s, 3H), 6.72 (d, J = 2.8 Hz, 1H), 7.23-7.27 (m, 1H), 7.30-7.45 (m, 11H), 7.54 (d, J = 8.4 Hz, 2H), 8.58 ppm (s, 1H); ^{13}C NMR (100 MHz, CD_2Cl_2) δ 136.68 (C), 136.42 (C), 133.05 (C), 131.74 (C), 129.28 (C), 129.10 (C), 128.68 (CH), 128.32 (CH), 127.50 (CH), 127.13 (CH), 126.99 (CH), 125.94 (CH), 125.74 (C), 124.09 (CH), 123.77 (C), 108.27 (CH), 15.72 ppm (CH_3); HRMS (ESI) m/z calcd for $\text{C}_{23}\text{H}_{19}\text{NS}^+$: 341.1223, found 341.1233.

General Procedure for Preparation of *N*-Boc hydrazones 1

A 100 mL-round-bottom flask equipped with a magnetic stirrer was charged with arylketone **6** (5 mmol), acetic acid (29 μL , 0.5 mmol) and EtOH (30.0 mL). The mixture was stirred and heated under reflux for 15 minutes. Then *tert*-butyl carbazate (0.990 g, 7.5 mmol) was added to the mixture and subjected to reflux. Upon completion as indicated by TLC, the crude mixture was concentrated under reduced pressure. The solid remained was then filtered and recrystallized with a minimum amount of hot DCM.

Desired product **1** was used without further purification.

General Procedure for Preparation of 1s

A 100 mL-round-bottom flask equipped with a magnetic stirrer was charged with acetone (40 mL), acetic acid (0.2 mL) and *tert*-butyl carbazate (5 g, 37.8 mmol). The mixture was subjected to reflux for 1h under a nitrogen atmosphere. The crude mixture was cooled and the resulting suspension was filtered off. The filtrate was concentrated under reduced pressure. The residue solid was then recrystallized with a minimum amount of hot DCM. Desired product **1s** was used without further purification.

1s was obtained as a white solid (6.37 g, 98% yield), mp 103 - 104 °C. ^1H NMR (400 MHz, CDCl_3) δ 1.39 (s, 9H), 1.73 (s, 3H), 1.90 (s, 3H), 7.58 ppm (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 153.02 (C), 149.87 (C), 80.56 (C), 28.16 (CH_3), 25.23 (CH_3), 16.05 ppm (CH_3).

Acknowledgements

We thank The Hong Kong Research Grants Council (PolyU5032/13P, C5023/14G) for financial support.

References

- [1] Reviews on transition metal-catalyzed oxidative C(sp²)–H annulations with alkynes, see: a) M. Gulías, J. L. Mascareñas, *Angew. Chem. Int. Ed.* **2016**, *55*, 11000; b) H. Huang, X. Ji, W. Wu, H. Jiang, *Chem. Soc. Rev.* **2015**, *44*, 1155, and the referenes cited therein; c) R. Chinchilla, C. Nájera, *Chem. Rev.* **2014**, *114*, 1783; d) L. Ackermann, *Acc. Chem. Res.* **2014**, *47*, 281; e) A. A. Tabolin, S. L. Ioffe, *Chem. Rev.* **2014**, 5426.
- [2] Selected examples on transition metal-catalyzed oxidative C(sp²)–H annulations for indoles synthesis, see: a) B. Li, H. Xu, H. Wang, B. Wang, *ACS Catal.* **2016**, *6*, 3856; b) Q. Lu, S. Vásquez-Céspedes, T. Gensch, F. Glorius, *ACS Catal.* **2016**, *6*, 2352; c) A. Lerchen, S. Vásquez-Céspedes, F. Glorius, *Angew. Chem. Int. Ed.* **2016**, *55*, 3208; d) W.-J. Chen, Z. Lin, *Organometallics* **2015**, *34*, 309; e) G. Zhang, H. Yu, G. Qin, H. Huang, *Chem. Commun.* **2014**, *50*, 4331; f) L. Zheng, R. Hua, *Chem. Eur. J.* **2014**, *20*, 2352; g) K. Muralirajan, C.-H. Cheng, *Adv. Synth. Catal.* **2014**, *356*, 1571; h) D. Zhao, Z. Shi, F. Glorius, *Angew. Chem. Int. Ed.* **2013**, *52*, 12426; i) B. Liu, C. Song, C. Sun, S. Zhou, J. Zhu, *J. Am. Chem. Soc.* **2013**, *135*, 16625; j) C. Wang, Y. huang, *Org. Lett.* **2013**, *15*, 5294; k) D. R. Stuart, P. Alsabeh, M. Kuhn, K. Fagnou, *J. Am. Chem. Soc.* **2010**, *132*, 18326; l) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess, K. Fagnou, *J. Am. Chem. Soc.* **2008**, *130*, 16474; m) R. C. Larock, E. K. Yum, M. D. Refvik, *J. Org. Chem.* **1998**, *63*, 7652; n) R. C. Larock, E. K. Yum, *J. Am. Chem. Soc.* **1991**, *113*, 6689. For isoquinolines synthesis, see: o) S. Zhang, D. Huang, G. Xu, S. Cao, R.

- Wang, S. Peng, J. Sun, *Org. Biomol. Chem.* **2015**, *13*, 7920; p) P. C. Too, Y.-F. Wang, S. Chiba, *Org. Lett.* **2010**, *12*, 5688. For pyridines synthesis, see: q) P. C. Too, T. Noji, Y. J. Lim, X. Li, S. Chiba, *Synlett.* **2011**, *19*, 2789. For isoquinolones synthesis, see: r) N. Guimond, C. Gouliaras, K. Fagnou, *J. Am. Chem. Soc.* **2010**, *132*, 6908.
- [3] Selected examples on the transition metal-catalyzed carbenoids cross coupling reactions involving C(sp²)-H functionalizations, see: a) J. Li, M. Tang, L. Zang, X. Zhang, Z. Zhang, L. Ackermann, *Org. Lett.* **2016**, *18*, 2742; b) Y.-S. Lu, W.-Y. Yu, *Org. Lett.* **2016**, *18*, 1350; c) G.-D. Tang, C.-L. Pan, X. Li, *Org. Chem. Front.* **2016**, *3*, 87; d) F.-N. Ng, Y.-F. Lau, Z. Zhou, W.-Y. Yu, *Org. Lett.* **2015**, *17*, 1676; e) D. Zhao, J. H. Kim, L. Stegemann, C. A. Strassert, F. Glorius, *Angew. Chem. Int. Ed.* **2015**, *54*, 4508; f) H.-W. Lam, K.-Y. Man, W.-W. Chan, Z. Zhou, W.-Y. Yu, *Org. Biomol. Chem.* **2014**, *12*, 4112; g) Y. Xia, Z. Liu, R. Ge, F. Ye, M. Hossain, Y. Zhang, J. Wang, *J. Am. Chem. Soc.* **2014**, *136*, 3013; h) T. K. Hyster, K. E. Ruhl, T. Rovis, *J. Am. Chem. Soc.* **2013**, *135*, 5364; i) W.-W. Chan, S.-F. Lo, Z. Zhou, W.-Y. Yu, *J. Am. Chem. Soc.* **2012**, *134*, 13565.
- [4] a) M. Baumann, I. R. Baxendale, S. V. Ley, N. Nikbin, *Beilstein J. Org. Chem.* **2011**, *7*, 442; b) J.-S. Lee, N.-Y. Kang, Y. K. Kim, A. Samanta, S. Feng, H. K. Kim, M. Vendrell, J. H. Park, Y.-T. Chang, *J. Am. Chem. Soc.* **2009**, *131*, 10077; c) H. Fan, J. Peng, M. T. Hamann, J.-F. Hu, *Chem. Rev.* **2008**, *108*, 264; d) A. Loudet, K. Burgess, *Chem. Rev.* **2007**, *107*, 4891; e) A. Grube, M. Köck, *Org. Lett.* **2006**, *8*, 4675; f) Z. S. Yoon, J. H. Kwon, M.-K. Koh, S. B. Noh, J. L. Sessler, J. T. Lee, D. Seidel, A. Aguilar, S. Shimizu, M. Suzuki, A. Osuka, D. Kim, *J. Am. Chem. Soc.* **2006**, *128*, 14128; g) A. Fürstner, *Angew. Chem. Int. Ed.* **2003**, *42*, 3582; h) M. Fujita, Y. Nakao, S. Matsunaga, M. Seiki, Y. Itoh, J. Yamashita, R. W. M. van Soest, N. Fusetani, *J. Am. Chem. Soc.* **2003**, *125*, 15700.
- [5] a) V. Estévez, M. Villacampa, J. C. Menéndez, *Chem. Soc. Rev.* **2014**, *43*, 4633; b) B. M. Trost, G. A. Doherty, *J. Am. Chem. Soc.* **2000**, *122*, 3801; c) A. Alberola, A. G. Ortega, M. L. Sádaba, C. Sañuda, *Tetrahedron* **1999**, *55*, 6555; d) A. Hantzsch, *Ber. Dtsch. Chem. Ges.* **1890**, *23*, 1474; e) C. Paal, *Ber. Dtsch. Chem. Ges.* **1885**, *18*, 1474.
- [6] a) M. P. Huestis, L. Chan, D. R. Stuart, K. Fagnou, *Angew. Chem. Int. Ed.* **2011**, *50*, 1338; b) S. Rakshit, F. W. Patureau, F. Glorius, *J. Am. Chem. Soc.* **2010**, *132*, 9585.
- [7] a) B. Li, N. Wang, Y. Liang, S. Xu, B. Wang, *Org. Lett.* **2013**, *15*, 136; b) L. Wang, L. Ackermann, *Org. Lett.* **2013**, *15*, 176; c) K. Murugan, S.-T. Liu, *Tetrahedron. Lett.* **2013**, *54*, 2608.
- [8] a) Y.-H. Xu, T. He, Q.-C. Zhang, T.-P. Loh, *Chem. Commun.* **2014**, *50*, 2784; b) M.-N. Zhao, Z.-H. Ren, Y.-Y. Wang, Z.-H. Guan, *Org. Lett.* **2014**, *16*, 608.
- [9] W. Yu, W. Zhang, Y. Liu, Y. Zhou, Z. Liu, Y. Zhang, *RSC Adv.* **2016**, *6*, 24768.
- [10] Selected examples on the rhodium-catalyzed cross coupling reactions involving C(sp³)-H functionalizations, for C(sp³)-H cycloaddition, see: a) A. Archambeau, T. Rovis, *Angew. Chem. Int. Ed.* **2015**, *54*, 13337; b) B. Zhou, Z. Chen, Y. Yang, W. Ai, H. Tang, Y. Wu, W. Zhu, Y. Li, *Angew. Chem. Int. Ed.* **2015**, *54*, 12121; c) C.-Z. Luo, P. Gandeepan, Y.-C. Wu, C.-H. Tsai, C.-H. Cheng, *ACS Catal.* **2015**, *5*, 4837; d) X. Tan, B. Liu, X. Li, B. Li, S. Xu, H. Song, B. Wang, *J. Am. Chem. Soc.* **2012**, *134*, 16163. For C(sp³)-H amidation, see: e) X.-H. Hu, X.-F. Yang, T.-P. Loh, *ACS Catal.* **2016**, *6*, 5930; f) X. Huang and J. You, *Chem. Lett.* **2015**, *44*, 1685; g) X. Huang, Y. Wang, J. Lan, J. You, *Angew. Chem. Int. Ed.* **2015**, *54*, 9404; h) H. Wang, G. Tang, X. Li, *Angew. Chem. Int. Ed.* **2015**, *54*, 13049; i) N. Wang, R. Li, L. Li, S. Xu, H. Song, B. Wang, *J. Org. Chem.* **2014**, *79*, 5379. For C(sp³)-H alkenylation, see: j) X. Huang, M. Wang, L. Wang, J. Wang, *J. Organomet. Chem.* **2015**, *787*, 1; k) J. Jiang, R. Ramozzi, K. Morokuma, *Chem. Eur. J.* **2015**, *21*, 11158; l) B. Liu, T. Zhou, B. Li, S. Xu, H. Song, B. Wang, *Angew. Chem. Int. Ed.* **2014**, *53*, 4191. For C(sp³)-H arylation, see: m) X. Wang, D.-G. Yu, F. Glorius, *Angew. Chem. Int. Ed.* **2015**, *54*, 10280. For C(sp³)-H silylation, see: n) Y. Kuninobu, T. Nakahara, H. Takeshima, K. Takai, *Org. Lett.* **2013**, *15*, 426. For C(sp³)-H borylation, see: o) S. Kawamorita, T. Miyazaki, T. Iwai, H. Ohmiya, M. Sawamura, *J. Am. Chem. Soc.* **2012**, *134*, 12924. For C(sp³)-H allylic amination, see: p) T. Cochet, V. Bellosta, D. Roche, J.-Y. Ortholand, A. Greiner, J. Cossy, *Chem. Commun.* **2012**, *48*, 10745.
- [11] Selected examples on transition metal -catalyzed pyrrole synthesis involving C(sp³)-H functionalizations, see: a) X. Chen, Y. Xie, X. Xiao, G. Li, Y. Deng, H. Jiang, W. Zeng, *Chem. Commun.* **2015**, *51*, 15328; b) Y. Xie, T. Chen, S. Fu, X.-S. Li, Y. Deng, H. Jiang, W. Zeng, *Chem. Commun.* **2014**, *50*, 10699; c) L. Meng, K. Wu, C. Liu, A. Lei, *Chem. Commun.* **2013**, *49*, 5853.
- [12] See Supporting Information for details.
- [13] a) S. Zhou, J. Wang, F. Zhang, C. Song, J. Zhu, *Org. Lett.* **2016**, *18*, 2427; b) S. Yu, S. Liu, Y. Lan, B. Wan, X. Li, *J. Am. Chem. Soc.* **2015**, *137*, 1623; c) S.-C. Chuang, P. Gandeepan, C.-H. Cheng, *Org. Lett.* **2013**, *15*, 5750; d) Y. Li, S. Liu, Z. Qi, X. Qi, X. Li, Y. Lan, *Chem. Eur. J.* **2015**, *21*, 10131; e) H. Wang, F. Glorius, *Angew. Chem. Int. Ed.* **2012**, *51*, 7318; f) J. Willwacher, S. Rakshit, F. Glorius, *Org. Biomol. Chem.* **2011**, *9*, 4736. For a review on oxidizing directing groups, see: g) J. Mo, L. Wang, Y. Liu, X. Cui, *Synthesis* **2015**, *47*, 439.
- [14] a) Y. Tan, J. F. Hartwig, *J. Am. Chem. Soc.* **2010**, *132*, 3676; b) S. Rakshit, C. Grohmann, T. Besset, F. Glorius, *J. Am. Chem. Soc.* **2011**, *133*, 2350.
- [15] The crystals of **3ea** were obtained from DCM:hexane (1:4) upon slow solvent diffusion. CCDC 1495321 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

- [16] Selected examples on the palladium-catalyzed cross coupling reactions involving secondary C(sp³)-H functionalizations, see: a) S. Ye, W. Yang, T. Coon, D. Fanning, T. Neubert, D. Stamos, J.-Q. Yu, *Chem. Eur. J.* **2016**, *22*, 4748. b) F.-L. Zhang, K. Hong, T.-J. Li, H. Park, J.-Q. Yu, *Science* **2016**, *351*, 252.
- [17] N. Wang, B. Li, H. Song, S. Xu, B. Wang, *Chem. Eur. J.* **2013**, *19*, 358.
- [18] D. Zhao, S. Vásquez-Céspedes, F. Glorius, *Angew. Chem. Int. Ed.* **2015**, *54*, 1657.
- [19] a) B. Sun, T. Yoshino, M. Kanai, S. Matsunaga, *Angew. Chem. Int. Ed.* **2015**, *54*, 12968; b) H. Wang, J. Koeller, W. Liu, L. Ackermann, *Chem. Eur. J.* **2015**, *21*, 15525; c) M. Sen, D. Kalsi, B. Sundararaju, *Chem. Eur. J.* **2015**, *21*, 15529.
- [20] a) L. Xu, Q. Zhu, G. Huang, B. Cheng, Y. Xia, *J. Org. Chem.* **2012**, *77*, 3017; b) D. Zhao, Z. Shi, F. Glorius, *Angew. Chem. Int. Ed.* **2013**, *52*, 12426.

Rhodium-Catalyzed Oxidative Cycloaddition of *N*-*tert*-Butoxycarbonyl Hydrazones with Alkynes for the Synthesis of Functionalized Pyrroles via C(sp³)-H Bond Functionalization

Adv. Synth. Catal. **Year**, *Volume*, Page – Page

Chun-Ming Chan, Zhongyuan Zhou and Wing-Yiu Yu*

