


This document is the Accepted Manuscript version of a Published Work that appeared in final form in Organic Letters, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <https://doi.org/10.1021/acs.orglett.6b00283>.

Cp*Rh(III)-Catalyzed Cross Coupling of Alkyltrifluoroborate with α -Diazomalonates for C(sp³)-C(sp³) Bond Formation

Yin-Suo Lu and Wing-Yiu Yu*

State Key Laboratory of Chirosciences and Department of Applied Biology and Chemical Technology, The Hong Kong Polytechnic University, Hung Hum, Kowloon, Hong Kong



ABSTRACT: A Cp*Rh(III)-catalyzed cross coupling of alkyltrifluoroborate with α -diazomalonates was developed; the C(sp³)-C(sp³) bonds coupled products were formed in up to 97% yields. The reaction tolerates some useful functional groups, including ketone, ester, amide, ether, sulfonyl, and thiophene. Electrospray ionization mass spectrometry (ESI-MS) analysis revealed the formation of a distinct molecular species corresponding to σ -alkylrhodium(III) complexes. The successful diazo coupling reaction may be attributed to coordination of the amide group that promotes stability of the alkylrhodium(III) complex through the formation of a five-membered metallacycle.


The use of diazo compounds for transition metal-catalyzed cross coupling reactions has recently received much attention.¹ Pioneered by van Vrenken,² Barluenga³ and Wang⁴, arylpalladium(II) complexes have been extensively investigated for diazo coupling reactions for synthesis of aryl-substituted alkenes.⁵⁻⁸ The diazo coupling reaction is believed to involve reactive arylpalladium-carbene complexes, and migratory carbene insertion would afford an σ -alkylpalladium(II) complex characterized by a quaternary stereocenter. Owing to the high propensity of the d⁸ σ -alkylpalladium(II) to undergo β -hydride elimination, further functionalization of the σ -alkyl complexes has met limited success.⁹

Our recent findings revealed that organorhodium-mediated diazo cross coupling reactions would offer a unique point of entry to synthesis of functionalized sp³ carbon centers.^{6c-f} For instance, arylrhodium(III) complexes of 1,2,3,4,5-pentamethylcyclopentadienyl (Cp*) ligand were found to couple with diazomalonates to form σ -alkylrhodium(III) complexes by migratory carbene insertion. With benzoquinoline as the substrate for the diazomalonate coupling reaction, the σ -alkylrhodium(III) complex has been structurally characterized.^{6d} While protonolysis of the σ -alkylrhodium(III) would lead to C-H bond formation,^{6d} the analogous alkyl complexes derived from benzohydroxamic acids would react by intramolecular N-O bond cleavage to effect C-N bond formation.^{6e} Recently, we achieved a [Cp*Rh(III)]-catalyzed cascade arylation-chlorination of diazomalonates for the synthesis of α -aryl- α -chlorocarbonyl compounds.^{6f} The cascade reaction should occur via migratory carbene insertion of the arylrhodium(III) to form a 1,3-diketone complex, followed by chlorination with *N*-chlorosuccinimide.

Despite the facile C(aryl)-C(sp³) bond formation by the diazo cross coupling reactions, the analogous diazo reactions with σ -alkylmetal complexes for C(sp³)-C(sp³) bond formation are rare.^{1,10-11} Recently, Szabó and co-workers reported a copper-catalyzed cross coupling of substituted allylboronic acids with diazoketones for effective C(sp³)-C(sp³) bond formations.¹² Nevertheless, σ -alkyl-transition metal complexes bearing β -hydrogen are reactive toward β -hydride elimination, and the development of broadly applicable catalytic alkyl-alkyl cross coupling reactions remains a formidable challenge. Here we report a successful [Cp*Rh(III)]-catalyzed cross coupling reaction of α -diazomalonates with primary alkyltrifluoroborates bearing β -hydrogens; up to 97% C(sp³)-C(sp³) coupled alkanes were obtained.

At the outset, potassium alkyltrifluoroborate **1a** was employed as the model substrate for the Rh-catalyzed diazo coupling reaction.¹³ **1a** was prepared by copper(I)-catalyzed β -boration of α,β -unsaturated carbonyl compounds according to a literature method.¹⁴ When **1a** (0.2 mmol) was treated with diazomalonate **2a** (0.2 mmol), [Cp*RhCl₂]₂ (2.5 mol%), AgOAc (15 mol%) in MeOH (1.5 mL) at 60 °C for 20 h,^{6d} the desired C(sp³)-C(sp³) coupled product was formed in 96% isolated yield (entry 1, Table 1). In the absence of AgOAc, the “**1a** + **2a**” coupling reaction still afforded **3a** in 85% isolated yield (entry 2). Changing solvent to aprotic solvents, such as THF, dioxane, toluene, DCE and CH₃CN [combined with water in a ratio of 20:1 (v/v)] resulted in lower product yields (51-85% NMR yield; entries 3-7).

Table 1. Reaction Optimization and Substrate Scope Studies^a




entry	n	X	solvent	yield (%)
1	1	NMe ₂ (1a)	MeOH	96 ^b
2	1	NMe ₂ (1a)	MeOH	85 ^c (93 ^d)
3	1	NMe ₂ (1a)	THF/H ₂ O (20:1)	85 ^d
4	1	NMe ₂ (1a)	dioxane/H ₂ O (20:1)	61 ^d
5	1	NMe ₂ (1a)	toluene/H ₂ O (20:1)	80 ^d
6	1	NMe ₂ (1a)	DCE/H ₂ O (20:1)	80 ^d
7	1	NMe ₂ (1a)	CH ₃ CN/H ₂ O (20:1)	51 ^d
8	1	N(OMe)Me (1b)	MeOH	71 ^b
9	1	Et (1c)	MeOH	5 ^d
10	1	<i>t</i> BuO (1d)	MeOH	8 ^d
11	2	NMe ₂ (1e)	MeOH	6 ^d
12	3	NMe ₂ (1f)	MeOH	<3 ^d

^a Reaction conditions: potassium alkyltrifluoroborate **1** (0.2 mmol), **2a** (0.2 mmol), [Cp*RhCl₂]₂ (2.5 mol %), AgOAc (15 mol %), solvent (1.5 mL) at 60 °C for 20 h under N₂ atmosphere. ^b isolated yield. ^c isolated yield without AgOAc as additive. ^d detected by ¹H NMR spectroscopy using 0.1 mmol dibromomethane as internal standard.

The structural requirements of the alkyltrifluoroborate for effective diazo coupling reactions have been examined. First, when **1b** containing a Weinreb amide moiety was employed as substrate, the diazomalonate coupling afforded **3ba** in 71% isolated yield (entry 8, Table 1). Notably, the related keto (**1c**) and ester (**1d**) derivatives were poor substrates for the diazo coupling reaction with <10% yield of the desired products being formed based on ¹H NMR analysis of the reaction mixture (entries 9-10). Interestingly, side products dimethylmalonate (**4a**) and malonate dimer **4b** were detected.¹⁵ Moreover, with *N,N*-dimethylamido moiety as the terminal group, the effect of chain length has also been studied. While the coupling of **1a** with diazomalonate **2a** afforded **3aa** in 96% yield, the analogous reactions of the alkyltrifluoroborates with a longer carbon chain [n = 2 (**1e**); n = 3 (**1f**)]¹⁶ gave negligible product yields. Notably, dimethyl malonate **4a** was found to be the major side product (entries 11-12, see Supporting Information for details). To account for the above findings, coordination of the amide group should promote stability of the putative alkylrhodium(III) complex through formation a five-membered metallacycle.¹⁷

We next turned to examine the diazomalonate substrate scope (Scheme 1). The coupling reaction using amide substrate **1a** always gave better yields than the substrates bearing Weinreb amide (**1b**).¹⁸ Changing the ester group from methyl to ethyl, benzyl, and *tert*-butyl were tolerated in the standard condition, and the corresponding products were obtained in excellent (**3aa-3ad**) and good (**3ba-3bd**) yields. Similar to our earlier findings,^{6f} both disubstituted (**3af**, **3bf**, **3ak** and **3al**) and tri-substituted C=C bonds (**3ag**, **3bg**, **3ah** and **3aj**) were tolerated under the Rh^{III}-catalyzed conditions. Assuming the alkylrhodium(III) intermediate, this finding suggested that migratory carbene insertion is kinetically more competitive than intramolecular cyclopropanation. The Rh(III)-catalyzed reaction is also compatible with functional groups such as ether, sulfonyl and thiophene, and the corresponding products (**3aj**, **3am** and **3an**) were formed in 85%, 85% and 33% yield, respectively.

Scheme 1. Substrate Scope study of diazomalonate^{a,b}



^a Reaction conditions: potassium alkyltrifluoroborate **1a-b** (0.2 mmol), **2a** (0.2 mmol), [Cp*RhCl₂]₂ (2.5 mol %), AgOAc (15 mol %), MeOH (1.5 mL) at 60 °C for 20 h under N₂ atmosphere. ^b isolated yield

For the reactions of some amide-containing diazo compounds, lower coupling yields (**3an-3aq**, 31-50%) were encountered for **2n**, **2o** and **2p** each containing an *N*-Ar group. Notably, 70% cross-coupled product was obtained for **2q** bearing a dihydroindole moiety.

It is plausible that the diazo coupling reaction is initiated by transmetalation with the alkyltrifluoroborate to form a σ -alkylrhodium(III) complex **A**. In this work, the reaction of **1a** with [Cp*Rh(OAc)₂] was analyzed by ESI-MS technique. As shown in Figure 1, a prominent peak at *m/z* = 338.0984 was observed (full scan). More detailed analysis of the peak composition revealed two distinct molecular ion species with maximum *m/z* peak intensity at 338.0984 and 336.0828. These ion species are assigned respectively as the σ -alkylrhodium(III) complex **5a** and π -acrylamide-rhodium(III) complex **5b** based on isotope simulation studies.¹⁹

Scheme 2 depicts a plausible mechanism for the diazo coupling reaction. The alkylrhodium(III) complex **A** generated by transmetalation from the organoborate would then react with diazo carbonyl compound, presumably via formation of rhodium-carbene **B**. Subsequent migratory carbene insertion would afford alkylrhodium complex **C**. Yet, a concerted alkyl migration-dinitrogen elimination pathway cannot be ruled out. Complex **C** should undergo protonolysis with MeOH to furnish the alkane product.

ASSOCIATED CONTENT

Supporting Information

Detailed experimental procedures, analytical data, and copies of NMR spectra of the products (PDF). These materials are available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* E-mail: wing-yiu.yu@polyu.edu.hk.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank The Hong Kong Research Grants Council (PolyU5032/13P, C5023/14G) for financial support. Ms. Melody Y. M. Wong (The Polytechnic University of Hong Kong) is thanked for technical assistance involving ESI-MS analysis.

REFERENCES

- (1) For reviews, see: (a) Barluenga, J.; Valdés, C. *Angew. Chem., Int. Ed.* **2011**, *50*, 7486. (b) Shao, Z.; Zhang, H. *Chem. Soc. Rev.* **2012**, *41*, 560. (c) Xia, Y.; Zhang, Y.; Wang, J. *ACS Catal.* **2013**, *3*, 2586. (d) Xiao, Q.; Zhang, Y.; Wang, J. *Acc. Chem. Res.* **2013**, *46*, 236. (e) Zhang, Y.; Wang, J. *Eur. J. Org. Chem.* **2011**, 1015. (f) Ford, A.; Miel, H.; Ring, A.; Slattery, C. N.; Maguire, A. R.; McKevey, M. A. *Chem. Rev.* **2015**, *115*, 9981. (g) Doyle, M. P.; Duffy, R.; Ratnikov, M.; Zhou, L. *Chem. Rev.* **2010**, *110*, 704.
- (2) (a) Greenman, K. L.; Carter, D. S.; Van Vranken, D. L. *Tetrahedron* **2001**, *57*, 5219. (b) Khanna, A.; Maung, C.; Johnson, K. R.; Luong, T. T.; Van Vranken, D. L. *Org. Lett.* **2012**, *14*, 3233. (c) Kudirka, R.; Van Vranken, D. L. *J. Org. Chem.* **2008**, *73*, 3585. (d) Gutman, E. S.; Arredondo, V.; Van Vranken, D. L. *Org. Lett.* **2014**, *16*, 5498.
- (3) (a) Barluenga, J.; Moriel, P.; Valdés, C.; Aznar, F. *Angew. Chem., Int. Ed.* **2007**, *46*, 5587. (b) Barluenga, J.; Florentino, L.; Aznar, F.; Valdés, C. *Org. Lett.* **2011**, *13*, 510. (c) Barluenga, J.; Tomas-Gamasa, M.; Aznar, F.; Valdés, C. *Chem.-Eur. J.* **2010**, *16*, 12801.
- (4) (a) Wang, X.; Xu, Y.; Deng, Y.; Zhou, Y.; Feng, J.; Ji, G.; Zhang, Y.; Wang, J. *Chem.-Eur. J.* **2014**, *20*, 961. (b) Xia, Y.; Xia, Y.; Liu, Z.; Zhang, Y.; Wang, J. *J. Org. Chem.* **2014**, *79*, 7711. (c) Zhou, Y.; Ye, F.; Wang, X.; Xu, S.; Zhang, Y.; Wang, J. *J. Org. Chem.* **2015**, *80*, 6109.
- (5) Other selected examples for the Pd-catalyzed carbenoid cross coupling reaction: (a) Chen, Z.-S.; Duan, X.-H.; Zhou, P.-X.; Ali, S.; Luo, J.-Y.; Liang, Y.-M. *Angew. Chem., Int. Ed.* **2012**, *51*, 1370. (b) Zhou, P.-X.; Luo, J.-Y.; Zhao, L.-B.; Ye, Y.-Y.; Liang, Y.-M. *Chem. Commun.* **2013**, *49*, 3254. (c) Zhou, P.-X.; Zhou, Z.-Z.; Chen, Z.-S.; Ye, Y.-Y.; Zhao, L.-B.; Yang, Y.-F.; Xia, X.-F.; Luo, J.-Y.; Liang, Y.-M. *Chem. Commun.* **2013**, *49*, 561. Also see ref 6a-b.
- (6) For our previous studies on carbenoid cross-coupling reactions, [Pd]: (a) Tsoi, Y.-T.; Zhou, Z.; Chan, A. S. C.; Yu, W.-Y. *Org. Lett.* **2010**, *12*, 4506. (b) Yu, W.-F.; Tsoi, Y.-T.; Zhou, Z.; Chan, A. S. C. *Org. Lett.* **2009**, *11*, 469. [Rh]: (c) Tsoi, Y.-T.; Zhou, Z.; Yu, W.-Y. *Org. Lett.* **2011**, *13*, 5370. (d) Chan, W.-W.; Lo, S.-F.; Zhou, Z.; Yu, W.-Y. *J. Am. Chem. Soc.* **2012**, *134*, 13565. (e) Lam, H.-W.; Man, K.-Y.; Chan, W.-W.; Zhou, Z.; Yu, W.-Y. *Org. Biomol. Chem.* **2014**, *12*, 4112. (f) Ng, F.-N.; Lau, Y.-F.; Zhou, Z.; Yu, W.-Y. *Org. Lett.* **2015**, *17*, 1676.
- (7) Other selected examples for the Rh-catalyzed carbenoid cross-coupling reaction: [Rh]: (a) Liu, Z.; Xia, Y.; Feng, S.; Wang, S.; Qiu, D.; Zhang, Y.; Wang, J. *Aust. J. Chem.* **2015**, *68*, 1379. (b) Xia, Y.; Liu, Z.; Feng, S.; Ye, F.; Zhang, Y.; Wang, J. *Org. Lett.* **2015**, *17*, 956. (c) Ghorai, J.; Anbarasan, P. *J. Org. Chem.* **2015**, *80*, 3455. [Rh^{III}]: (d) Chen, X.; Hu, X.; Bai, S.; Deng, Y.; Jiang, H.; Zeng, W. *Org. Lett.* **2016**, *18*, 192. (e) Dateer, R. B.; Chang, S. *Org. Lett.* **2016**, *18*, 68. (f)





Figure 1. ESI-MS analysis of the reaction of $[\text{Cp}^*\text{Rh}(\text{OAc})_2]$ with **1a** in MeCN

Scheme 2. Plausible Reaction Mechanism



In summary, the first Rh(III)-catalyzed cross coupling reaction of alkyltrifluoroborate with α -diazomalonates for $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$ bond formation is developed. The present strategy brings about $\text{C}(\text{sp}^3)\text{-C}(\text{sp}^3)$ bond formation in moderate to excellent yields under an operationally simple and mild reaction conditions. Importantly, the chelation of the carbonyl oxygen to the Rh center through five-membered metallacycle should stabilize the alkylmetal intermediate to avoid spontaneous β -hydride elimination. Furthermore, The Rh-coupling reaction tolerates various functional groups such as ketone, ester, amide, ether, sulfonyl, thiophene, etc. Further exploration of this reaction for multi-component reaction is under investigation.

Son, J.-Y.; Kim, S.; Jeon, W. H.; Lee, P. H. *Org. Lett.* **2015**, *17*, 2518. (g) Yu, S.; Liu, S.; Lan, Y.; Wan, B.; Li, X. *J. Am. Chem. Soc.* **2015**, *137*, 1623. (h) Jeong, J.; Patel, P.; Hwang, H.; Chang, S. *Org. Lett.* **2014**, *16*, 4598. (i) Shi, Z.; Koester, D. C.; Bouladakis-Arapinis, M.; Glorius, F. *J. Am. Chem. Soc.* **2013**, *135*, 12204. (j) Yu, X.; Yu, S.; Xiao, J.; Wan, B.; Li, X. *J. Org. Chem.* **2013**, *78*, 5444. Also see ref 6c-f.

(8) Selected examples for the Fe-, Cu-, Ni-, Au-, Ir- and Co-catalyzed carbenoid cross-coupling reactions: [Fe]: (a) Prabharasuth, R.; Van Vranken, D. L. *J. Org. Chem.* **2001**, *66*, 5256. [Cu]: (b) Xiao, Q.; Xia, Y.; Li, H.; Zhang, Y.; Wang, J. *Angew. Chem., Int. Ed.* **2011**, *50*, 1114. [Ni]: (c) Ni, Y.; Montgomery, J. *J. Am. Chem. Soc.* **2006**, *128*, 2609. (d) Yu, Z.; Ma, B.; Chen, M.; Wu, H.-H.; Liu, L.; Zhang, J. *J. Am. Chem. Soc.* **2014**, *136*, 6904. [Au] (e) Yu, Z.; Ma, B.; Chen, M.; Wu, H.-H.; Liu, L.; Zhang, J. *J. Am. Chem. Soc.* **2014**, *136*, 6904. [Ir]: (f) Xia, Y.; Liu, Z.; Feng, S.; Zhang, Y.; Wang, J. *J. Org. Chem.* **2015**, *80*, 223. (g) Phatake, R. S.; Patel, P.; Ramana, C. V. *Org. Lett.* **2016**, *18*, 292. [Co]: (h) Zhao, D.; Kim, J. H.; Stegemann, L.; Strassert, C. A.; Glorius, F. *Angew. Chem., Int. Ed.* **2015**, *54*, 4508. (i) Liu, X.-G.; Zhang, S.-S.; Wu, J.-Q.; Li, Q.; Wang, H. *Tetrahedron Lett.* **2015**, *56*, 4093.

(9) For successful examples of further functionalization of the σ -alkylpalladium complex: (a) Zhou, L.; Ye, F.; Zhang, Y.; Wang, J. *J. Am. Chem. Soc.* **2010**, *132*, 13590. (b) Kudirka, R.; Devine, S. K. J.; Adams, C. S.; Van Vranken, D. L. *Angew. Chem., Int. Ed.* **2009**, *48*, 3677. (c) Devine, S. K. J.; Van Vranken, D. L. *Org. Lett.* **2008**, *10*, 1909. (d) Devine, S. K. J.; Van Vranken, D. L. *Org. Lett.* **2007**, *9*, 2047.

(10) For reviews of $C(sp^3)-C(sp^3)$ bond formation: (a) Sigman, M. S. *Chem. Rev.* **2011**, *111*, 1417. (b) Han, F.-S. *Chem. Soc. Rev.* **2013**, *42*, 5270. (c) Kambe, N.; Iwasaki, T.; Terao, J. *Chem. Soc. Rev.* **2011**, *40*, 4937. (d) Rudolph, A.; Lautens, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 2656. (e) Frisch, A. C.; Beller, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 674.


(11) Selected examples of $C(sp^3)-C(sp^3)$ bond formation by alkyl halide/pseudohalide Suzuki-Miyaura cross-coupling reaction: (a) Ishiyama, S.; Abe, S.; Miyaura, N.; Suzuki, A. *Chem. Lett.* **1992**, *21*, 691. (b) Netherton, M. R.; Dai, C.; Neuschütz, K.; Fu, G. C. *J. Am. Chem. Soc.* **2001**, *123*, 10099. (c) Kirchhoff, J. H.; Dai, C.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 1945. (d) Netherton, M. R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 3910. (e) Kirchhoff, J. H.; Netherton, M. R.; Hills, I. D.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 13662. (f) Peh, G.-R.; Kantchev, E. A. B.; Er, J.-C.; Ying, J. Y. *Chem. Eur. J.* **2010**, *16*, 4010. (g) Saito, B.; Fu, G. C. *J. Am. Chem. Soc.* **2007**, *129*, 9602. (h) Saito, B.; Fu, G. C. *J. Am. Chem. Soc.* **2008**, *130*, 6694. (i) Lu, Z.; Wilsily, A.; Fu, G. C. *J. Am. Chem. Soc.* **2011**, *133*, 8154.

(12) Das, A.; Wang, D.; Belhomme, M.-C.; Szabó, K. J. *Org. Lett.* **2015**, *17*, 4754.

(13) For reviews of potassium organotrifluoroborates: (a) Darses, S.; Genet, J.-P.; *Chem. Rev.* **2008**, *108*, 288. (b) Molander, G. A.; Ellis, A. *Acc. Chem. Res.* **2007**, *40*, 275. (c) Stefani, H. A.; Cellia, R.; Vieira, A. *Tetrahedron* **2007**, *63*, 3623. (d) Molander, G. A.; Figueroa, R. *Aldrichim. Acta* **2005**, *38*, 49.

(14) (a) Mun, S.; Lee, J.-E.; Yun, J. *Org. Lett.* **2006**, *8*, 4887. (b) Molander, G. A.; Petrillo, D. E. *Org. Lett.* **2008**, *10*, 1795. (c) Molander, G. A.; Jean-Gérard, L. *J. Org. Chem.* **2009**, *74*, 5446. (d) Sandrock, D. L.; Jean-Gérard, L.; Chen, C.-y.; Dreher, S. D.; Molander, G. A. *J. Am. Chem. Soc.* **2010**, *132*, 17108.

(15) The following mechanism is proposed to account for the formation of compounds **4a** and **4b**:



(16) For preparation of amides **6a** and **6b** from their corresponding acids: (a) Lam, J. K.; Pham, H. V.; Houk, K. N.; Vanderval, C. D. *J. Am. Chem. Soc.* **2013**, *135*, 17585. For Rhodium-catalyzed hydroboration reaction: (b) Hadebe, S. W.; Robinson, R. S. *Eur. J. Org. Chem.* **2006**, 4898. For conversion to the potassium organotrifluoroborate salt see ref 14.

(17) Lee, J. C. H.; Sun, H.-Y.; Hall, D. G. *J. Org. Chem.* **2015**, *80*, 7134. For similar works also see ref 14b-d.

(18) Nigst, T. A.; Antipova, A.; Mayr, H. *J. Org. Chem.* **2012**, *77*, 8142.

(19) A mechanism is proposed to account for the formation of complexes **5a** and **5b**.

