

GLOBAL STABILIZATION OF THE FULL ATTRACTION-REPULSION KELLER-SEGEL SYSTEM

HAI-YANG JIN

Department of Mathematics
South China University of Technology
Guangzhou 510640, China

ZHI-AN WANG*

Department of Applied Mathematics
Hong Kong Polytechnic University
Hung Hom, Hong Kong, China

*To Professor Wei-Ming Ni on the occasion of his 70th birthday,
with our best wishes*

ABSTRACT. We are concerned with the following full Attraction-Repulsion Keller-Segel (ARKS) system

$$\begin{cases} u_t = \Delta u - \nabla \cdot (\chi u \nabla v) + \nabla \cdot (\xi u \nabla w), & x \in \Omega, t > 0, \\ v_t = D_1 \Delta v + \alpha u - \beta v, & x \in \Omega, t > 0, \\ w_t = D_2 \Delta w + \gamma u - \delta w, & x \in \Omega, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), w(x, 0) = w_0(x) & x \in \Omega, \end{cases} \quad (*)$$

in a bounded domain $\Omega \subset \mathbb{R}^2$ with smooth boundary subject to homogeneous Neumann boundary conditions. By constructing an appropriate Lyapunov functions, we establish the boundedness and asymptotical behavior of solutions to the system (*) with large initial data $(u_0, v_0, w_0) \in [W^{1,\infty}(\Omega)]^3$. Precisely, we show that if the parameters satisfy $\frac{\xi\gamma}{\chi\alpha} \geq \max\left\{\frac{D_1}{D_2}, \frac{D_2}{D_1}, \frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$ for all positive parameters $D_1, D_2, \chi, \xi, \alpha, \beta, \gamma$ and δ , the system (*) has a unique global classical solution (u, v, w) , which converges to the constant steady state $(\bar{u}_0, \frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\delta}\bar{u}_0)$ as $t \rightarrow +\infty$, where $\bar{u}_0 = \frac{1}{|\Omega|} \int_{\Omega} u_0 dx$. Furthermore, the decay rate is exponential if $\frac{\xi\gamma}{\chi\alpha} > \max\left\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$. This paper provides the first results on the full ARKS system with unequal chemical diffusion rates (i.e. $D_1 \neq D_2$) in multi-dimensions.

1. Introduction. To describe the aggregation of *Microglia* in the central nervous system in Alzheimer's disease due to the interaction of chemoattractant (i.e. β -amyloid) and chemorepellent (i.e. TNF- α), Luca et al. [21] proposed the following

2010 *Mathematics Subject Classification.* Primary: 35A01, 35B40, 35B44, 35K57, 35Q92, 92C17.

Key words and phrases. Chemotaxis, attraction-repulsion, global stability, exponential decay rate.

The research of H.Y. Jin was supported by the NSF of China No. 11871226, and the Fundamental Research Funds for the Central Universities. The research of Z.A. Wang acknowledges partial supports from an internal grant No. ZZHY in Hong Kong Polytechnic University and from NSFC grant 11571086.

* Corresponding author: Zhi-An Wang.

attraction-repulsion chemotaxis system

$$\begin{cases} u_t = \Delta u - \nabla \cdot (\chi u \nabla v) + \nabla \cdot (\xi u \nabla w), & x \in \Omega, t > 0, \\ \tau_1 v_t = D_1 \Delta v + \alpha u - \beta v, & x \in \Omega, t > 0, \\ \tau_2 w_t = D_2 \Delta w + \gamma u - \delta w, & x \in \Omega, t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = \frac{\partial w}{\partial \nu} = 0, & x \in \partial\Omega, t > 0, \\ u(x, 0) = u_0(x), \tau_1 v(x, 0) = \tau_1 v_0(x), \tau_2 w(x, 0) = \tau_2 w_0(x), & x \in \Omega, \end{cases} \quad (1)$$

where Ω is a bounded domain in \mathbb{R}^n with smooth boundary $\partial\Omega$ and ν denotes the outward normal vector of $\partial\Omega$. The density of *Microglia* cells is denoted by $u(x, t)$, while $v(x, t)$ and $w(x, t)$ denote the concentration of chemoattractant and chemorepellent, respectively. The model (1) can also be regarded as a particularized model proposed in [23] to model the quorum sensing effect in chemotaxis.

When $\xi = 0$, the variable w can be decoupled from the system (1), where the variables u and v satisfy the classical attractive Keller-Segel (KS) system

$$\begin{cases} u_t = \Delta u - \chi \nabla \cdot (u \nabla v), & x \in \Omega, t > 0, \\ v_t = D_1 \Delta v + \alpha u - \beta v, & x \in \Omega, t > 0. \end{cases} \quad (2)$$

The KS model (2) has been extensively studied in the past four decades in various perspectives and massive results are available (cf. survey articles [6, 1] and references therein). One of the mostly studied topics for the KS model (2) is the boundedness and blowup of solutions in two or higher dimensions [22, 30, 7] based on the following Lyapunov function:

$$\mathcal{E}_1(u, v) = \int_{\Omega} u \ln u - \chi \int_{\Omega} uv + \frac{\beta\chi}{2\alpha} \int_{\Omega} v^2 + \frac{\chi D_1}{2\alpha} \int_{\Omega} |\nabla v|^2.$$

If $\chi = 0$, the variable v can be decoupled and (u, w) satisfies the following repulsive Keller-Segel model

$$\begin{cases} u_t = \Delta u + \xi \nabla \cdot (u \nabla w), & x \in \Omega, t > 0, \\ w_t = D_2 \Delta w + \gamma u - \delta w, & x \in \Omega, t > 0. \end{cases} \quad (3)$$

Compared to the attractive KS model (2), the results on the repulsive KS model (3) are much less. The global existence of classical solutions in two dimensions and weak solutions in three or four dimensions were established in [4] based on the following Lyapunov function

$$\mathcal{E}_2(u, w) = \int_{\Omega} u \ln u + \frac{\tau_2 \xi}{2\gamma} \int_{\Omega} |\nabla w|^2$$

which is difference from the one for the attractive KS model. A further investigation on the repulsive KS model was made in [26].

Roughly speaking, the attraction-repulsion Keller-Segel model (1) can be regarded as a superposition of the attractive and repulsive KS models. Hence one may expect the ARKS model should behave more or less the same as the attractive or repulsive models. However it is not straightforward to justify this suspicion due to the interaction between attraction and repulsion. In particular, as we recalled above, the understanding of the attractive and repulsive KS models heavily rely on the finding of Lyapunov functions. Therefore to have a comprehensive understanding for the ARKS model, finding appropriate Lyapunov function is indispensable. This is by no means an easy work for a strongly coupled cross-diffusion system of PDEs like ARKS model. A sequence of works thus have been stimulated to reveal

the mystery underlying the model gradually. The first such progress was made by Tao and Wang [27] who found that the solution behavior of the ARKS model was essentially determined by the sign of

$$\theta_1 = \xi\gamma - \chi\alpha,$$

which is an index measuring the competition between attraction and repulsion. More precisely, they showed that if $D_1 = D_2 = 1$ and $\tau_1 = \tau_2 = 0$, the ARKS system (1) has a unique classical solution with uniform-in-time bound if $\theta_1 \geq 0$ (i.e. repulsion dominates or cancels attraction) in higher dimensions ($n \geq 2$). The main idea of [27] was a transformation $s = \xi w - \chi v$ which may significantly simplify the system and has become a major source for many of subsequent researches on the ARKS model. For the opposite case $\theta_1 < 0$ (i.e. attraction dominates repulsion), it was shown that the solution of system (5) may blow up in finite time if initial mass is large [5, 14] and exist globally for small initial mass [5] in two dimensions. If $\tau_1 = 1$ and $\tau_2 = 0$, Jin and Wang [11] constructed a Lyapunov function

$$\begin{aligned} \mathcal{E}_3(u, v, w) = & \int_{\Omega} u \ln u - \chi \int_{\Omega} uv + \frac{\beta\chi}{2\alpha} \int_{\Omega} v^2 \\ & + \frac{\chi D_1}{2\alpha} \int_{\Omega} |\nabla v|^2 + \frac{\xi\delta}{2\gamma} \int_{\Omega} w^2 + \frac{\xi D_2}{2\gamma} \int_{\Omega} |\nabla w|^2, \end{aligned}$$

to establish the global existence of uniformly-in-time bounded classical solutions in two dimensions for large initial data if $\theta_1 \geq 0$. Conversely if $\theta_1 < 0$, they showed there exists a critical mass m_* such that the solution blows up if $\int_{\Omega} u_0 > m_*$ and globally exists if $\int_{\Omega} u_0 < m_*$.

If the three equations of the ARKS model (1) are all parabolic (i.e. $\tau_1 = \tau_2 = 1$), it is much harder to study and much less results are available. We recall the known results below. In one dimension, the global existence of classical solutions, non-trivial stationary state, asymptotic behavior and pattern formation of the system (1) have been studied in [10, 19, 20]. In two dimensions, when $D_1 = D_2$, it was shown in [27] that global classical solutions exist for large data if $\beta = \delta$ and for small data if $\beta \neq \delta$ when $\theta_1 \geq 0$ (i.e. repulsion dominates or cancels attraction). Subsequently the global existence of large-data solutions was extended to the case $\beta \neq \delta$ in [8, 18]. Moreover, for $\beta \neq \delta$, when cell mass is small, it was shown that the global classical solution will exponentially converge to the unique constant steady state $(\bar{u}_0, \frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\delta}\bar{u}_0)$ with $\bar{u}_0 = \frac{1}{|\Omega|} \int_{\Omega} u_0$ in [15, 16], which was further elaborated by assuming

$$\bar{u}_0 < \frac{4\beta\delta}{\chi\alpha(\beta - \delta)^2} \text{ and } \xi\gamma > \frac{4\beta\delta(\chi\alpha\bar{u}_0 + 1)}{[4\beta\delta - (\beta - \delta)^2\chi\alpha\bar{u}_0]\bar{u}_0} \tag{4}$$

in [17] wherein the convergence rate was, however, not given. Whether or not the same results holds for large initial data in multi-dimensions still remains unknown. Part of above-mentioned results have been extended to the multi-dimensional whole space in [9, 25]. We should underline that all existing results in two or higher dimensions recalled above for the case $\tau_1 = \tau_2 = 1$ are essentially based on the assumption $D_1 = D_2$ so that the idea of making a change of variable $s = \xi w - \chi v$ introduced in [27] can be employed. To the best of our knowledge, no result for the case $\tau_1 = \tau_2 = 1$ and $D_1 \neq D_2$ has been available to (1) in multi-dimensions to date. It is the purpose of this paper to exploit this challenging case and contribute

some results, where the corresponding ARKS model (1) reads as

$$\begin{cases} u_t = \Delta u - \nabla \cdot (\chi u \nabla v) + \nabla \cdot (\xi u \nabla w), & x \in \Omega, t > 0, \\ v_t = D_1 \Delta v + \alpha u - \beta v, & x \in \Omega, t > 0, \\ w_t = D_2 \Delta w + \gamma u - \delta w, & x \in \Omega, t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = \frac{\partial w}{\partial \nu} = 0, & x \in \partial \Omega, t > 0, \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), w(x, 0) = w_0(x), & x \in \Omega. \end{cases} \quad (5)$$

The main challenge is that when $D_1 \neq D_2$ the conventional approach of using the transformation in [27] is no longer effective and new ideas are desirable. Here we shall construct a Lyapunov functional for (5) which allows us to establish the global boundedness and asymptotic behavior of solutions to (5) in some parameter regimes. Specifically, the following results are obtained in the paper.

Theorem 1.1. *Let Ω be a bounded domain in \mathbb{R}^2 with smooth boundary. Suppose that $0 \leq (u_0, v_0, w_0) \in [W^{1,\infty}(\Omega)]^3$ and the parameters satisfy*

$$\frac{\xi\gamma}{\chi\alpha} \geq \max \left\{ \frac{D_1}{D_2}, \frac{D_2}{D_1}, \frac{\beta}{\delta}, \frac{\delta}{\beta} \right\}. \quad (6)$$

Then the problem (5) has a unique classical solution $(u, v, w) \in [C^0([0, \infty) \times \bar{\Omega}) \cap C^{2,1}((0, \infty) \times \bar{\Omega})]^3$, which satisfies

$$\|(u, v, w)(\cdot, t)\|_{L^\infty} \leq C \quad (7)$$

for some constant $C > 0$ independent of t and

$$\|(u, v, w) - (\bar{u}_0, \frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\delta}\bar{u}_0)\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow +\infty,$$

where $\bar{u}_0 = \frac{1}{|\Omega|} \int_{\Omega} u_0 dx$. Furthermore, if $\frac{\xi\gamma}{\chi\alpha} > \max \left\{ \frac{\beta}{\delta}, \frac{\delta}{\beta} \right\}$, the decay is exponential.

Remark 1. If $D_1 = D_2 = 1$ and $\beta = \delta$, Tao and Wang [27, Proposition 2.6] proved that if $\theta_1 > 0$ the global classical solution (u, v, w) of system (5) exists and exponentially converges to the constant steady state $(\bar{u}_0, \frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\delta}\bar{u}_0)$ as $t \rightarrow \infty$. Hence in this paper, we will, unless otherwise mentioned, focus on the case $D_1 \neq D_2$ or $\beta \neq \delta$ under which the condition (6) implies $\theta_1 > 0$.

Remark 2. The results of Theorem 1.1 hold for all $D_1, D_2, \alpha, \beta, \xi, \gamma > 0$ without any smallness conditions on initial data under the parameter regime given by (6). In the case $D_1 = D_2 = 1$ and $\beta \neq \delta$, the same result was recently obtained in [17] under the essential assumption (4) where the initial cell mass can not be arbitrarily large and parameter regime depends upon the initial data. Hence our results not only improve those of [17], but also cover the case $D_1 \neq D_2$ for which no results have been known so far.

Outline of proof: We first establish the boundedness criterion of solution for system (5) such that the boundedness of $\|u\|_{L^\infty}$ can be reduced to prove the boundedness of $\|u\|_{L^p}$ with $p > \max\{1, \frac{n}{2}\}$. Motivated by the results in [8, 18], we know that the boundedness of $\|u\|_{L^2}$ holds in two dimensions if there exists a constant $c_1 > 0$ such that

$$\|u \ln u\|_{L^1} + \|\nabla v\|_{L^2} + \|\nabla w\|_{L^2} \leq c_1. \quad (8)$$

Hence to show the global existence of classical solutions in two dimensions, we only need to prove (8). When $D_1 = D_2$ and $\theta_1 > 0$, using the transformation $s = \xi w - \chi v$ as in [27], one can derive the following entropy inequality (cf. [18, 8])

$$\frac{d}{dt} \left(\int_{\Omega} u \ln u + \frac{1}{2\theta_1} \int_{\Omega} |\nabla s|^2 \right) + \int_{\Omega} \frac{|\nabla u|^2}{u} + \frac{D_1}{2\theta_1} \int_{\Omega} |\Delta s|^2 + \frac{\delta}{\theta_1} \int_{\Omega} |\nabla s|^2 \leq c_2, \tag{9}$$

which can be used to derive (8) and hence the boundedness of solutions.

However, when $D_1 \neq D_2$, the transformation idea fails to work. Luckily, we are able to find a different Lyapunov function $E(u, v, w)$ (see the definition in (31)) for the system (5) under the condition (6), which satisfies

$$\frac{d}{dt} E(u, v, w) + F(u, v, w) = 0, \tag{10}$$

where $F(u, v, w)$ is defined by (32). We remark that the form of $E(u, v, w)$ is quite different from the one (9) for $D_1 = D_2$. To prove $E(u, v, w)$ is a Lyapunov function, we organize the estimates into a quadratic form which is the new idea developed in the paper. Then using (10), we show that under the condition (6), there exists a constant $c_3 > 0$ such that $\|u \ln u\|_{L^1} + \|\nabla v\|_{L^2} + \|\nabla w\|_{L^2} \leq c_3$ and $\int_0^t \int_{\Omega} \frac{|\nabla u|^2}{u} \leq c_3$ (see Lemma 4.1 for details). The former estimate leads to the boundedness of solutions in two dimensions and the later estimate gives the convergence properties of u . The convergence of v and w can be derived by the parabolic comparison principle. To study the decay rate, we first show that there exists a constant $\mu > 0$ such that

$$E(u, v, w) \leq \mu F(u, v, w),$$

which together with (10) gives $E(u, v, w) \leq E(u_0, v_0, w_0)e^{-\frac{1}{\mu}t}$. Using the definition of $E(u, v, w)$ and noting the fact $\|u - \bar{u}\|_{L^1} \leq 2\bar{u} \int_{\Omega} u \ln \frac{u}{\bar{u}}$ in Lemma 2.1, the exponential decay of $\|u - \bar{u}\|_{L^1}$ under the condition (6) is obtained. Then using the ideas in [27] or [15], we derive the decay rate of $\|u - \bar{u}\|_{L^\infty}$ and hence the exponential decay rate of $\|v - \frac{\alpha}{\beta}\bar{u}_0\|_{L^\infty}$ and $\|w - \frac{\gamma}{\delta}\bar{u}_0\|_{L^\infty}$.

In the end of this section, we remark that Theorem 1.1 only present some first-hand results on the full ARKS model for $D_1 \neq D_2$ under the parameter regime given in (6) and leave out many interesting questions due to technical difficulty. For example, whether the condition (6) is necessary for global existence of solutions and how solutions behave (in particular whether solutions blow up) if the condition (6) fails remain unsolved in our paper. We hope our studies in this paper will provide useful clues to further explore the ARKS model in future.

2. Some basic inequalities. In what follows, without confusion, we shall abbreviate $\int_{\Omega} f dx$ as $\int_{\Omega} f$ for simplicity. Moreover, we shall use $c_i (i = 1, 2, 3, \dots)$ to denote a generic constant which may vary in the context. For reader’s convenience, we present some known inequalities for later use.

Lemma 2.1. *Suppose that $f(x, t)$ is a positive function on $(x, t) \in \Omega \times (0, \infty)$. Defined $\bar{f} = \frac{1}{|\Omega|} \int_{\Omega} f$, then it has that*

$$0 \leq \frac{1}{2\bar{f}} \|f - \bar{f}\|_{L^1}^2 \leq \int_{\Omega} f \ln \frac{f}{\bar{f}} \leq \frac{1}{\bar{f}} \|f - \bar{f}\|_{L^2}^2. \tag{11}$$

Proof. Using the Csiszár-Kullback-Pinsker inequality (see [3]), one has

$$\int_{\Omega} f \ln \frac{f}{\bar{f}} \geq \frac{1}{2\bar{f}} \|f - \bar{f}\|_{L^1}^2. \tag{12}$$

On the other hand, choosing $\psi = \frac{f}{\bar{f}}$ and using the fact that $\psi \ln \psi - \psi + 1 \leq (\psi - 1)^2$ for $\psi \geq 0$, it holds that

$$\int_{\Omega} f \ln \frac{f}{\bar{f}} \leq \bar{f} \int_{\Omega} \left[\frac{f}{\bar{f}} - 1 + \left(\frac{f}{\bar{f}} - 1 \right)^2 \right] = \frac{1}{\bar{f}} \|f - \bar{f}\|_{L^2}^2. \tag{13}$$

Then the combination of (12) and (13) gives (11). □

Lemma 2.2. *Let Ω be a bounded domain in \mathbb{R}^2 with smooth boundary. Then, for any $\varphi \in W^{3,2}(\Omega)$ satisfying $\frac{\partial \varphi}{\partial \nu}|_{\partial \Omega} = 0$, there exists a positive constant C depending only on Ω such that*

$$\|\Delta \varphi\|_{L^3} \leq C(\|\nabla \Delta \varphi\|_{L^2}^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + \|\nabla \varphi\|_{L^2}). \tag{14}$$

Proof. Using Gagliardo-Nirenberg inequality, we have

$$\|\Delta \varphi\|_{L^3} = \|\nabla \cdot \nabla \varphi\|_{L^3} \leq \|D \nabla \varphi\|_{L^3} \leq c_1 \|D^2 \nabla \varphi\|_{L^2}^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + c_2 \|\nabla \varphi\|_{L^2}, \tag{15}$$

where $|D^k \nabla \varphi| = (\sum_{|i|=k} |D^i \nabla \varphi|^2)^{\frac{1}{2}}$ and i is a multi-index of order k . On the other hand, one can check that

$$\|D^2 \nabla \varphi\|_{L^2} \leq c_3 \|\nabla \varphi\|_{H^2}. \tag{16}$$

Moreover, under the homogeneous Neumann boundary condition (i.e., $\frac{\partial \varphi}{\partial \nu}|_{\partial \Omega} = 0$), it follows from [2, Lemma 1] that $\|\nabla \varphi\|_{H^2} \leq c_4 \|\Delta \varphi\|_{H^1}$, which applied to (16) gives

$$\|D^2 \nabla \varphi\|_{L^2} \leq c_3 c_4 \|\Delta \varphi\|_{H^1}. \tag{17}$$

Note that $|\Delta \varphi|^2 = \nabla \cdot (\nabla \varphi \Delta \varphi) - \nabla \varphi \cdot \nabla \Delta \varphi$. Then using the boundary condition $\frac{\partial \varphi}{\partial \nu}|_{\partial \Omega} = 0$ and Hölder inequality, we have

$$\|\Delta \varphi\|_{L^2}^2 = - \int_{\Omega} \nabla \varphi \cdot \nabla \Delta \varphi \leq \|\nabla \varphi\|_{L^2} \|\nabla \Delta \varphi\|_{L^2}. \tag{18}$$

Then substituting (17) into (15), and using (18), one derives

$$\begin{aligned} \|\Delta \varphi\|_{L^3} &\leq c_5 \left(\|\Delta \varphi\|_{H^1}^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + \|\nabla \varphi\|_{L^2} \right) \\ &= c_5 \left(\|\nabla \Delta \varphi\|_{L^2} + \|\Delta \varphi\|_{L^2} \right)^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + c_5 \|\nabla \varphi\|_{L^2} \\ &\leq c_6 \left(\|\nabla \Delta \varphi\|_{L^2} + \|\nabla \Delta \varphi\|_{L^2}^{\frac{1}{2}} \|\nabla \varphi\|_{L^2}^{\frac{1}{2}} \right)^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + c_6 \|\nabla \varphi\|_{L^2} \\ &\leq c_6 \left(2 \|\nabla \Delta \varphi\|_{L^2} + \|\nabla \varphi\|_{L^2} \right)^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + c_6 \|\nabla \varphi\|_{L^2} \\ &\leq c_7 \|\nabla \Delta \varphi\|_{L^2}^{\frac{2}{3}} \|\nabla \varphi\|_{L^2}^{\frac{1}{3}} + c_7 \|\nabla \varphi\|_{L^2} \end{aligned}$$

which yields (14), and hence completes the proof. □

3. Boundedness criterion and Lyapunov function.

3.1. Local existence. The local existence theorem of system (5) can be proved by the fixed point theorem and maximum principle along the same line as in [27]. Hence we only present the results without proof for brevity.

Lemma 3.1. *Let Ω be a bounded domain in $\mathbb{R}^n (n \geq 2)$ with smooth boundary. Suppose that $0 \leq (u_0, v_0, w_0) \in [W^{1,\infty}(\Omega)]^3$. Then there exist a $T_{max} \in (0, \infty]$ such that the system (5) has a unique solution (u, v, w) of nonnegative functions from $[C^0(\bar{\Omega} \times [0, T_{max})) \cap C^{2,1}(\bar{\Omega} \times (0, T_{max}))]^3$. Moreover $u > 0$ in $\Omega \times (0, T_{max})$ and*

$$\text{if } T_{max} < \infty, \text{ then } \|u(\cdot, t)\|_{L^\infty} \rightarrow \infty \text{ as } t \nearrow T_{max}. \tag{19}$$

Furthermore, the cell mass is conservative:

$$\|u(\cdot, t)\|_{L^1} = \|u_0\|_{L^1}. \tag{20}$$

3.2. Boundedness criterion. To extend the local solutions to global ones, we derive a boundedness criterion for the solution of system (5). The idea of our proof is essentially inspired by [1, lemma 3.2] and we present necessary details below for clarity.

Lemma 3.2. *Suppose the conditions in Lemma 3.1 hold. Let (u, v, w) be the solution of system (5) defined on its maximal existence time interval $[0, T_{max})$. If there exist $p > \frac{n}{2}$ and a constant M_0 such that*

$$\sup_{t \in (0, T_{max})} \|u(\cdot, t)\|_{L^p} \leq M_0,$$

then one can find a constant $C > 0$ independent of t such that

$$\|u(\cdot, t)\|_{L^\infty} + \|v(\cdot, t)\|_{W^{1,\infty}} + \|w(\cdot, t)\|_{W^{1,\infty}} \leq C \text{ for all } t \in (0, T_{max}). \tag{21}$$

Furthermore, there exists $\sigma \in (0, 1)$ such that for all $t > 1$

$$\|u\|_{C^{\sigma, \frac{\sigma}{2}}(\bar{\Omega} \times [t, t+1])} \leq C. \tag{22}$$

Proof. Since $\|u(\cdot, t)\|_{L^p} \leq M_0$, then applying the parabolic regularity estimates in [12, Lemma 1] to the second and third equations of system (5) we have

$$\|\nabla v(\cdot, t)\|_{L^r} + \|\nabla w(\cdot, t)\|_{L^r} \leq c_1, \text{ for all } t \in (0, T_{max}) \tag{23}$$

where

$$r \in \begin{cases} [1, \frac{np}{n-p}), & \text{if } p \leq n, \\ [1, \infty], & \text{if } p > n. \end{cases} \tag{24}$$

Without loss of generality, we assume that $\frac{n}{2} < p \leq n$ which yields $\frac{np}{n-p} > n$. Then we can find a constant $r > 0$ with $n < r < \frac{np}{n-p}$ such that (23) holds. Now, for each $T \in (0, T_{max})$, we define

$$M(T) := \sup_{t \in (0, T)} \|u(\cdot, t)\|_{L^\infty}, \tag{25}$$

which is finite due to the local existence results in Lemma 3.1. Next, we will estimate $M(T)$. Fix $t \in (0, T)$ and let $t_0 = (t-1)_+$. Then applying the variation-of-constants formula to the first equation of system (5), we get

$$\begin{aligned} u(\cdot, t) &= e^{(t-t_0)\Delta} u(\cdot, t_0) - \chi \int_{t_0}^t e^{(t-\tau)\Delta} \nabla \cdot (u(\cdot, \tau) \nabla v(\cdot, \tau)) d\tau \\ &\quad + \xi \int_{t_0}^t e^{(t-\tau)\Delta} \nabla \cdot (u(\cdot, \tau) \nabla w(\cdot, \tau)) d\tau \end{aligned}$$

which implies

$$\begin{aligned} \|u(\cdot, t)\|_{L^\infty} &\leq \|e^{(t-t_0)\Delta}u(\cdot, t_0)\|_{L^\infty} + \chi \int_{t_0}^t \|e^{(t-\tau)\Delta}\nabla \cdot (u(\cdot, \tau)\nabla v(\cdot, \tau))\|_{L^\infty} d\tau \\ &\quad + \xi \int_{t_0}^t \|e^{(t-\tau)\Delta}\nabla \cdot (u(\cdot, \tau)\nabla w(\cdot, \tau))\|_{L^\infty} d\tau \\ &= I_1 + I_2 + I_3. \end{aligned} \tag{26}$$

We first estimate the term I_1 . If $t \leq 1$, then $t_0 = 0$ and we can use the maximum principle for the heat equation to obtain

$$I_1 = \|e^{t\Delta}u_0\|_{L^\infty} \leq \|u_0\|_{L^\infty} \leq c_2, \tag{27}$$

whereas in the case $t > 1$ and $t_0 = t - 1$, we use the standard L^p - L^q estimates for $(e^{\tau\Delta})_{\tau \geq 0}$ to derive

$$I_1 \leq c_3 \|u(\cdot, t_0)\|_{L^p} \leq c_3 M_0 = c_4. \tag{28}$$

Moreover, since $r > n$, we can fix a number $q > n$ satisfying $q \in (\frac{r}{r+1}, r)$. Then by the Hölder inequality, interpolation inequality and (25), we can find $\zeta = \frac{r(q-1)+q}{rq} \in (0, 1)$ such that

$$\begin{aligned} \|u(\cdot, \tau)\nabla v(\cdot, \tau)\|_{L^q} &\leq \|u(\cdot, \tau)\|_{L^{\frac{rq}{r-q}}} \|\nabla v(\cdot, \tau)\|_{L^r} \\ &\leq \|u(\cdot, \tau)\|_{L^\infty}^{1-\frac{r-q}{rq}} \|u(\cdot, \tau)\|_{L^1}^{\frac{r-q}{rq}} \|\nabla v(\cdot, \tau)\|_{L^r} \\ &\leq c_5 M^\zeta(T). \end{aligned}$$

Similarly, we have

$$\|u(\cdot, \tau)\nabla w(\cdot, \tau)\|_{L^q} \leq c_6 M^\zeta(T).$$

Since $t - t_0 \leq 1$, we have $\int_{t_0}^t (t-s)^{-\frac{1}{2}-\frac{n}{2q}} ds = \int_0^{t-t_0} \sigma^{-\frac{1}{2}-\frac{n}{2q}} d\sigma \leq \int_0^1 \sigma^{-\frac{1}{2}-\frac{n}{2q}} d\sigma = \frac{2q}{q-n}$ thanks to $q > n$. Then by the smoothing properties of $(e^{\tau\Delta})_{\tau \geq 0}$ (see [29, Lemma 1.3]), we derive

$$\begin{aligned} I_2 + I_3 &\leq c_7 \int_{t_0}^t (t-\tau)^{-\frac{1}{2}-\frac{n}{2q}} (\|u(\cdot, \tau)\nabla v(\cdot, \tau)\|_{L^q} + \|u(\cdot, \tau)\nabla w(\cdot, \tau)\|_{L^q}) d\tau \\ &\leq c_8 M^\zeta(T) \int_{t_0}^t (t-\tau)^{-\frac{1}{2}-\frac{n}{2q}} d\tau \\ &\leq \frac{2qc_8}{q-n} M^\zeta(T) := c_9 M^\zeta(T). \end{aligned} \tag{29}$$

Substituting (27), (28) and (29) into (26), we can find a constant $c_{10} > 0$ such that

$$\|u(\cdot, t)\|_{L^\infty} \leq c_9 M^\zeta(T) + c_{10}, \quad \text{for all } t \in (0, T),$$

which implies

$$M(T) \leq c_9 M^\zeta(T) + c_{10}, \quad \text{for all } T \in (0, T_{max}). \tag{30}$$

Since $0 < \zeta < 1$, from (30) one has

$$M(T) \leq \max \left\{ \left(\frac{c_{10}}{c_9} \right)^{\frac{1}{1-\zeta}}, (2c_9)^{\frac{1}{1-\zeta}} \right\}, \quad \text{for all } T \in (0, T_{max}),$$

which implies $\|u(\cdot, t)\|_{L^\infty} \leq c_{11}$ for all $t \in (0, T_{max})$. Furthermore the combination of (23) and (24) gives (21).

At last, from (21) we know that $\chi u \nabla v$ and $\xi u \nabla w$ are bounded in $L^\infty(\Omega \times (0, \infty))$. Then applying the standard parabolic regularity theory (e.g. see [24, Theorem 1.3] and [28, Lemma 3.2]) and parabolic Schauder theory [13], we immediately obtain the estimate (22). Then the proof of Lemma 3.2 is completed. \square

3.3. Lyapunov function. As mentioned in Remark 1, we consider the case $D_1 \neq D_2$ or $\beta \neq \delta$ which implies that $\theta_1 > 0$ from (6). When $D_1 = D_2$, the boundedness of solutions shown in Theorem 1.1 has been proved in [27] with $\beta = \delta$ and in [18, 8] with $\beta \neq \delta$ by constructing entropy inequality based on an idea of using the transformation $s = \xi w - \chi v$. However this transformation is no longer helpful for the case $D_1 \neq D_2$. Hence, we need to find a new way. Here we achieve our results by constructing a Lyapunov function for the system (5). First, we define

$$E(u, v, w) := \frac{\theta_1}{2\xi\chi} \int_\Omega u \ln \frac{u}{\bar{u}} + \frac{\theta_2}{4\xi\alpha} \int_\Omega |\nabla v|^2 + \frac{\theta_2}{4\gamma\chi} \int_\Omega |\nabla w|^2 - \int_\Omega \nabla w \cdot \nabla v \quad (31)$$

and

$$F(u, v, w) := \frac{\theta_1}{2\xi\chi} \int_\Omega \frac{|\nabla u|^2}{u} + \frac{\theta_2 D_1}{2\xi\alpha} \int_\Omega |\Delta v|^2 + \frac{\theta_2 D_2}{2\gamma\chi} \int_\Omega |\Delta w|^2 + \frac{\theta_2 \beta}{2\xi\alpha} \int_\Omega |\nabla v|^2 + \frac{\theta_2 \delta}{2\gamma\chi} \int_\Omega |\nabla w|^2 - (D_1 + D_2) \int_\Omega \Delta w \Delta v - (\beta + \delta) \int_\Omega \nabla w \cdot \nabla v, \quad (32)$$

where $\theta_1 := \xi\gamma - \chi\alpha$ and $\theta_2 := \xi\gamma + \chi\alpha$. Then, we will show that $E(u, v, w)$ is indeed a Lyapunov function under (6). More precisely, we have the following results.

Lemma 3.3. *Let (u, v, w) be the solution of system (5). Then we have*

$$\frac{d}{dt} E(u, v, w) + F(u, v, w) = 0 \quad (33)$$

where $E(u, v, w)$ and $F(u, v, w)$ are defined by (31) and (32), respectively. Moreover, if (6) holds, then

$$E(u, v, w) \geq 0 \quad \text{and} \quad F(u, v, w) \geq 0 \quad \text{for all } t > 0. \quad (34)$$

Proof. Multiplying the first equation of system (5) by $\ln \frac{u}{\bar{u}}$, we have

$$\frac{d}{dt} \int_\Omega u \ln \frac{u}{\bar{u}} + \int_\Omega \frac{|\nabla u|^2}{u} = \chi \int_\Omega \nabla u \cdot \nabla v - \xi \int_\Omega \nabla u \cdot \nabla w. \quad (35)$$

Similarly, we multiply the second and third equations of system (5) by $-\Delta v$ and $-\Delta w$, respectively, to obtain

$$\frac{1}{2} \frac{d}{dt} \int_\Omega |\nabla v|^2 + D_1 \int_\Omega |\Delta v|^2 + \beta \int_\Omega |\nabla v|^2 = \alpha \int_\Omega \nabla u \cdot \nabla v \quad (36)$$

and

$$\frac{1}{2} \frac{d}{dt} \int_\Omega |\nabla w|^2 + D_2 \int_\Omega |\Delta w|^2 + \delta \int_\Omega |\nabla w|^2 = \gamma \int_\Omega \nabla u \cdot \nabla w. \quad (37)$$

Multiplying (35) by $\frac{\theta_1}{2\xi\chi}$, (36) by $\frac{\theta_2}{2\xi\alpha}$ and (37) by $\frac{\theta_2}{2\gamma\chi}$, and adding them, we end up with

$$\frac{d}{dt} \left(\frac{\theta_1}{2\xi\chi} \int_\Omega u \ln \frac{u}{\bar{u}} + \frac{\theta_2}{4\xi\alpha} \int_\Omega |\nabla v|^2 + \frac{\theta_2}{4\gamma\chi} \int_\Omega |\nabla w|^2 \right) + \frac{\theta_1}{2\xi\chi} \int_\Omega \frac{|\nabla u|^2}{u}$$

$$\begin{aligned}
 & + \frac{\theta_2 D_1}{2\xi\alpha} \int_{\Omega} |\Delta v|^2 + \frac{\theta_2 D_2}{2\gamma\chi} \int_{\Omega} |\Delta w|^2 + \frac{\theta_2 \beta}{2\xi\alpha} \int_{\Omega} |\nabla v|^2 + \frac{\theta_2 \delta}{2\gamma\chi} \int_{\Omega} |\nabla w|^2 \\
 & = \gamma \int_{\Omega} \nabla u \cdot \nabla v + \alpha \int_{\Omega} \nabla u \cdot \nabla w.
 \end{aligned} \tag{38}$$

On the other hand, the second and third equations of system (5) give us that

$$\begin{aligned}
 \gamma \int_{\Omega} \nabla u \cdot \nabla v & = \int_{\Omega} \nabla(w_t + \delta w - D_2 \Delta w) \cdot \nabla v \\
 & = \int_{\Omega} \nabla w_t \cdot \nabla v + \delta \int_{\Omega} \nabla w \cdot \nabla v - D_2 \int_{\Omega} \nabla(\Delta w) \cdot \nabla v \\
 & = \frac{d}{dt} \int_{\Omega} \nabla w \cdot \nabla v + \int_{\Omega} \Delta w v_t + \delta \int_{\Omega} \nabla w \cdot \nabla v + D_2 \int_{\Omega} \Delta w \Delta v \\
 & = \frac{d}{dt} \int_{\Omega} \nabla w \cdot \nabla v + \int_{\Omega} \Delta w (D_1 \Delta v + \alpha u - \beta v) \\
 & \quad + \delta \int_{\Omega} \nabla w \cdot \nabla v + D_2 \int_{\Omega} \Delta w \Delta v \\
 & = \frac{d}{dt} \int_{\Omega} \nabla w \cdot \nabla v + (D_1 + D_2) \int_{\Omega} \Delta w \Delta v \\
 & \quad + (\beta + \delta) \int_{\Omega} \nabla w \cdot \nabla v - \alpha \int_{\Omega} \nabla u \cdot \nabla w
 \end{aligned}$$

which yields

$$\begin{aligned}
 \gamma \int_{\Omega} \nabla u \cdot \nabla v + \alpha \int_{\Omega} \nabla u \cdot \nabla w & = \frac{d}{dt} \int_{\Omega} \nabla w \cdot \nabla v + (D_1 + D_2) \int_{\Omega} \Delta w \Delta v \\
 & \quad + (\beta + \delta) \int_{\Omega} \nabla w \cdot \nabla v.
 \end{aligned} \tag{39}$$

The combination of (38) and (39) gives (33).

Next, we will show the nonnegative of $E(u, v, w)$ and $F(u, v, w)$ under (6). First, we rewrite $E(u, v, w)$ in (31) as

$$E(u, v, w) = \frac{\theta_1}{2\xi\chi} \int_{\Omega} u \ln \frac{u}{\bar{u}} + \int_{\Omega} \Theta_1^T A_1 \Theta_1$$

where Θ_1^T denotes the transpose of Θ_1 and

$$\Theta_1 = \begin{bmatrix} \nabla v \\ \nabla w \end{bmatrix} \quad \text{and} \quad A_1 = \begin{bmatrix} \frac{\theta_2}{4\xi\alpha} & -\frac{1}{2} \\ -\frac{1}{2} & \frac{\theta_2}{4\gamma\chi} \end{bmatrix}.$$

Since $\theta_1 > 0$, one has $\theta_2^2 > \theta_2^2 - \theta_1^2 = 4\xi\gamma\chi\alpha$. This implies the matrix A_1 is positive definite and hence there exists a constant $c_1 > 0$ such that

$$E(u, v, w) \geq \frac{\theta_1}{2\xi\chi} \int_{\Omega} u \ln \frac{u}{\bar{u}} + c_1 \int_{\Omega} (|\nabla v|^2 + |\nabla w|^2) \geq 0, \tag{40}$$

where we have used the fact $\int_{\Omega} u \ln \frac{u}{\bar{u}} \geq 0$ from Lemma 2.1. Similarly, we rewrite $F(u, v, w)$ as

$$F(u, v, w) = \frac{\theta_1}{2\xi\chi} \int_{\Omega} \frac{|\nabla u|^2}{u} + \int_{\Omega} \Theta_2^T A_2 \Theta_2 + \int_{\Omega} \Theta_1^T A_3 \Theta_1, \tag{41}$$

where

$$\Theta_2 = \begin{bmatrix} \Delta v \\ \Delta w \end{bmatrix}, \quad A_2 = \begin{bmatrix} \frac{\theta_2 D_1}{2\xi\alpha} & -\frac{D_1 + D_2}{2} \\ -\frac{D_1 + D_2}{2} & \frac{\theta_2 D_2}{2\gamma\chi} \end{bmatrix} \quad \text{and} \quad A_3 = \begin{bmatrix} \frac{\theta_2 \beta}{2\xi\alpha} & -\frac{\beta + \delta}{2} \\ -\frac{\beta + \delta}{2} & \frac{\theta_2 \delta}{2\gamma\chi} \end{bmatrix}. \tag{42}$$

Clearly, the matrix A_2 is nonnegative definite if

$$\theta_2^2 D_1 D_2 - \frac{(D_1 + D_2)^2 (\theta_2^2 - \theta_1^2)}{4} \geq 0.$$

Similarly, the matrix A_3 is nonnegative definite under the condition

$$\theta_2^2 \beta \delta - \frac{(\beta + \delta)^2 (\theta_2^2 - \theta_1^2)}{4} \geq 0.$$

Hence, the nonnegativity of the matrices A_2 and A_3 are satisfied simultaneously if

$$\begin{cases} 4\theta_2^2 D_1 D_2 - (D_1 + D_2)^2 (\theta_2^2 - \theta_1^2) \geq 0, \\ 4\theta_2^2 \beta \delta - (\beta + \delta)^2 (\theta_2^2 - \theta_1^2) \geq 0, \end{cases}$$

which is equivalent to

$$\begin{cases} \theta_2^2 (D_1 - D_2)^2 \leq (D_1 + D_2)^2 \theta_1^2, \\ \theta_2^2 (\beta - \delta)^2 \leq (\beta + \delta)^2 \theta_1^2. \end{cases} \tag{43}$$

One can check that (43) holds if $\frac{\xi\gamma}{\chi\alpha} \geq \max\{\frac{D_1}{D_2}, \frac{D_2}{D_1}, \frac{\beta}{\delta}, \frac{\delta}{\beta}\}$ or $\frac{\xi\gamma}{\chi\alpha} \leq \min\{\frac{D_1}{D_2}, \frac{D_2}{D_1}, \frac{\beta}{\delta}, \frac{\delta}{\beta}\}$. However, the latter is impossible due to $\theta_1 > 0$. Hence, if (6) holds, one has $E(u, v, w) \geq 0$ and $F(u, v, w) \geq 0$. The proof of (34) is completed. \square

4. Proof of Theorem 1.1. In this section, we are devoted to proving Theorem 1.1 based on the Lyapunov function constructed in Lemma 3.3.

4.1. Boundedness of solutions. In this subsection, we show the boundedness of solutions for system (5) under the condition (6). First, we give a core lemma concerning the boundedness and asymptotical behavior of solution for system (5) in two dimensions.

Lemma 4.1. *Suppose that $(u_0, v_0, w_0) \in [W^{1,\infty}(\Omega)]^3$ and (6) hold. Then the solution (u, v, w) of system (5) satisfies*

$$\|u \ln u\|_{L^1} + \|\nabla v\|_{L^2} + \|\nabla w\|_{L^2} \leq C \tag{44}$$

and

$$\int_0^t \int_{\Omega} \frac{|\nabla u|^2}{u} \leq C, \tag{45}$$

where $C > 0$ is a constant independent of t .

Proof. The nonnegativity of $E(u, v, w)$ and $F(u, v, w)$ has been proved in Lemma 3.3 under the condition (6). Then integrating (33) and using (40) and (41), along with the nonnegativity of A_2 and A_3 , we have two positive constants c_1, c_2 such that

$$\frac{\theta_1}{2\xi\chi} \int_{\Omega} u \ln \frac{u}{\bar{u}} + c_1 \int_{\Omega} (|\nabla v|^2 + |\nabla w|^2) \leq c_2, \tag{46}$$

which, together with the fact $\int_{\Omega} u \ln \frac{u}{\bar{u}} \geq 0$ from Lemma 2.1, gives

$$\|\nabla v\|_{L^2}^2 + \|\nabla w\|_{L^2}^2 \leq \frac{c_2}{c_1} = c_3. \tag{47}$$

On the other hand, from (46), we directly obtain

$$\frac{\theta_1}{2\xi\chi} \int_{\Omega} u \ln u \leq c_2 + \frac{\theta_1}{2\xi\chi} |\Omega| \bar{u} \ln \bar{u} \leq c_4,$$

which, along with the fact $-u \ln u \leq \frac{1}{e}$ for all $u \geq 0$, gives

$$\int_{\Omega} |u \ln u| \leq \int_{\Omega} \left| u \ln u + \frac{1}{e} - \frac{1}{e} \right| \leq \int_{\Omega} \left(u \ln u + \frac{1}{e} \right) + \int_{\Omega} \frac{1}{e} \leq \frac{2\xi\chi c_4}{\theta_1} + \frac{2|\Omega|}{e}. \tag{48}$$

Then the combination of (47) and (48) gives (44). Hence the proof of this lemma is completed. \square

Lemma 4.2. *Let the assumptions in Lemma 4.1 hold. Then the solution (u, v, w) of system (5) satisfies*

$$\|u(\cdot, t)\|_{L^2} \leq C \tag{49}$$

where the constant $C > 0$ is independent of t .

Proof. Multiplying the first equation of system (5) by u and integrating it by parts, we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Omega} u^2 + \int_{\Omega} |\nabla u|^2 &= \chi \int_{\Omega} u \nabla u \cdot \nabla v - \xi \int_{\Omega} u \nabla u \cdot \nabla w \\ &= -\frac{\chi}{2} \int_{\Omega} u^2 \Delta v + \frac{\xi}{2} \int_{\Omega} u^2 \Delta w \\ &\leq c_1 \|u\|_{L^3}^2 (\|\Delta v\|_{L^3} + \|\Delta w\|_{L^3}). \end{aligned} \tag{50}$$

Noting the fact $\|u \ln u\|_{L^1} \leq c_2$ and $\|u\|_{L^1} \leq c_3$, one can find a small $\varepsilon > 0$ such that

$$\|u\|_{L^3}^2 = (\|u\|_{L^3}^3)^{\frac{2}{3}} \leq (\varepsilon \|\nabla u\|_{L^2}^2 + 1)^{\frac{2}{3}} \leq \varepsilon \|\nabla u\|_{L^2}^{\frac{4}{3}} + c_4, \tag{51}$$

where we have used the following fact (see [22]): when $n = 2$, for any $\varepsilon > 0$, there exists a constant C_ε such that

$$\|u\|_{L^3} \leq \varepsilon \|\nabla u\|_{L^2}^{\frac{2}{3}} \|u \ln u\|_{L^1}^{\frac{1}{3}} + C_\varepsilon (\|u \ln u\|_{L^1} + \|u\|_{L^1}^{\frac{1}{3}}).$$

On the other hand, noting the facts $\frac{\partial v}{\partial \nu} \Big|_{\partial \Omega} = \frac{\partial w}{\partial \nu} \Big|_{\partial \Omega} = 0$ on $\partial \Omega$ and using the boundedness of $\|\nabla v\|_{L^2}$ and $\|\nabla w\|_{L^2}$ (see (44)), from Lemma 2.2, one has

$$\begin{aligned} &\|\Delta v\|_{L^3} + \|\Delta w\|_{L^3} \\ &\leq c_5 (\|\nabla \Delta v\|_{L^2}^{\frac{2}{3}} \|\nabla v\|_{L^2}^{\frac{1}{3}} + \|\nabla v\|_{L^2}) + c_5 (\|\nabla \Delta w\|_{L^2}^{\frac{2}{3}} \|\nabla w\|_{L^2}^{\frac{1}{3}} + \|\nabla w\|_{L^2}) \\ &\leq c_6 (\|\nabla \Delta v\|_{L^2}^{\frac{2}{3}} + \|\nabla \Delta w\|_{L^2}^{\frac{2}{3}} + 1). \end{aligned} \tag{52}$$

Then combining (51) and (52), and using Young's inequality and noting the fact $\varepsilon > 0$ is small, we find a small $\eta > 0$ such that

$$\begin{aligned} &c_1 \|u\|_{L^3}^2 (\|\Delta v\|_{L^3} + \|\Delta w\|_{L^3}) \\ &\leq c_7 \left(\varepsilon \|\nabla u\|_{L^2}^{\frac{4}{3}} + c_4 \right) \left(\|\nabla \Delta v\|_{L^2}^{\frac{2}{3}} + \|\nabla \Delta w\|_{L^2}^{\frac{2}{3}} + 1 \right) \\ &= c_7 \varepsilon \|\nabla u\|_{L^2}^{\frac{4}{3}} \left(\|\nabla \Delta v\|_{L^2}^{\frac{2}{3}} + \|\nabla \Delta w\|_{L^2}^{\frac{2}{3}} \right) + c_7 \varepsilon \|\nabla u\|_{L^2}^{\frac{4}{3}} \\ &\quad + c_1 c_7 \left(\|\nabla \Delta v\|_{L^2}^{\frac{2}{3}} + \|\nabla \Delta w\|_{L^2}^{\frac{2}{3}} \right) + c_1 c_7 \\ &\leq \frac{1}{2} \|\nabla u\|_{L^2}^2 + \eta (\|\nabla \Delta v\|_{L^2}^2 + \|\nabla \Delta w\|_{L^2}^2) + c_8. \end{aligned} \tag{53}$$

Substituting (53) into (50) gives

$$\frac{d}{dt} \int_{\Omega} u^2 + \int_{\Omega} |\nabla u|^2 \leq 2\eta (\|\nabla \Delta v\|_{L^2}^2 + \|\nabla \Delta w\|_{L^2}^2) + c_9. \tag{54}$$

Differentiating the second equation of system (5) once, and multiplying the result by $-\nabla\Delta v$, and then we integrate the product in Ω to obtain

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\Delta v|^2 + D_1 \int_{\Omega} |\nabla\Delta v|^2 + \beta \int_{\Omega} |\Delta v|^2 \\ &= -\alpha \int_{\Omega} \nabla\Delta v \cdot \nabla u \\ &\leq \frac{D_1}{2} \|\nabla\Delta v\|_{L^2}^2 + \frac{\alpha^2}{2D_1} \|\nabla u\|_{L^2}^2, \end{aligned}$$

which yields

$$\frac{d}{dt} \int_{\Omega} |\Delta v|^2 + D_1 \int_{\Omega} |\nabla\Delta v|^2 + 2\beta \int_{\Omega} |\Delta v|^2 \leq \frac{\alpha^2}{D_1} \|\nabla u\|_{L^2}^2. \tag{55}$$

Similarly, we have the following estimates for w :

$$\frac{d}{dt} \int_{\Omega} |\Delta w|^2 + D_2 \int_{\Omega} |\nabla\Delta w|^2 + 2\delta \int_{\Omega} |\Delta w|^2 \leq \frac{\gamma^2}{D_2} \|\nabla u\|_{L^2}^2. \tag{56}$$

Letting $\rho = \frac{\alpha^2 D_2 + \gamma^2 D_1}{D_1 D_2}$, and multiplying (54) by 2ρ , then adding it with (55) and (56), we end up with

$$\begin{aligned} & \frac{d}{dt} (2\rho \|u\|_{L^2}^2 + \|\Delta v\|_{L^2}^2 + \|\Delta w\|_{L^2}^2) + \rho \|\nabla u\|_{L^2}^2 \\ &+ D_1 \|\nabla\Delta v\|_{L^2}^2 + D_2 \|\nabla\Delta w\|_{L^2}^2 + 2\beta \|\Delta v\|_{L^2}^2 + 2\delta \|\Delta w\|_{L^2}^2 \\ &\leq 4\rho\eta \cdot (\|\nabla\Delta v\|_{L^2}^2 + \|\nabla\Delta w\|_{L^2}^2) + c_{10}. \end{aligned} \tag{57}$$

Letting η small such that $4\rho\eta \leq \min\{D_1, D_2\}$, one has

$$\begin{aligned} & \frac{d}{dt} (2\rho \|u\|_{L^2}^2 + \|\Delta v\|_{L^2}^2 + \|\Delta w\|_{L^2}^2) + \rho \|\nabla u\|_{L^2}^2 \\ &+ 2\beta \|\Delta v\|_{L^2}^2 + 2\delta \|\Delta w\|_{L^2}^2 \leq c_{10}. \end{aligned} \tag{58}$$

On the other hand, using the Gagliardo-Nirenberg inequality and (20), we can show that

$$\|u\|_{L^2}^2 \leq c_{11} (\|\nabla u\|_{L^2} \|u\|_{L^1} + \|u\|_{L^1}^2) \leq \frac{1}{2} \|\nabla u\|_{L^2}^2 + c_{12}. \tag{59}$$

Substituting (59) into (58) and letting $y(t) := 2\rho \|u\|_{L^2}^2 + \|\Delta v\|_{L^2}^2 + \|\Delta w\|_{L^2}^2$, we can find two positive constants c_{13} and c_{14} such that

$$y'(t) + c_{13}y(t) \leq c_{14},$$

which, along with Gronwall's inequality gives (49). □

Next, we will show the existence of global classical solutions.

Lemma 4.3. *Suppose that the conditions in Lemma 4.1 hold. Then the problem (5) has a unique global classical solution $(u, v, w) \in [C^0([0, \infty) \times \bar{\Omega}) \cap C^{2,1}((0, \infty) \times \bar{\Omega})]^3$ satisfying (7).*

Proof. From Lemma 4.2, we know that there exists a constant $c_1 > 0$ such that $\|u(\cdot, t)\|_{L^2} \leq c_1$. Noting $n = 2$ and using Lemma 3.2, one has

$$\|u(\cdot, t)\|_{L^\infty} \leq c_2,$$

which together with the local existence results in Lemma 3.1 completes the proof of this lemma. □

4.2. Convergence. In this subsection, we will show the convergence of solutions.

Lemma 4.4. *Let (u, v, w) be the solution of system (5) satisfying (7) and (45). Then one has*

$$\|u(\cdot, t) - \bar{u}_0\|_{L^\infty} \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (60)$$

Proof. The combination of (7) and (45) implies that there exist a constant $c_1 > 0$ such that

$$\int_0^\infty \|\nabla u\|_{L^2}^2 \leq c_1. \quad (61)$$

Noting the conservation of cell mass and using the Poincaré inequality, we will derive

$$\|u(\cdot, t) - \bar{u}_0\|_{L^2}^2 = \|u(\cdot, t) - \bar{u}\|_{L^2}^2 \leq c_2 \|\nabla u\|_{L^2}^2. \quad (62)$$

Combining (61) and (62), one can find a constant $c_3 > 0$ such that

$$\int_0^\infty \|u(\cdot, t) - \bar{u}_0\|_{L^2}^2 \leq c_3. \quad (63)$$

Motivated by the ideas in [28, Lemma 3.10], we next show (63) implies (60). Indeed, if one can show that

$$\|u(\cdot, t) - \bar{u}_0\|_{C^0} \rightarrow 0, \quad \text{as } t \rightarrow \infty, \quad (64)$$

then (60) follows directly. We shall show (64) by the argument of contradiction. Suppose that (64) is wrong, then for some constant $c_4 > 0$, there exist some sequences $(x_j)_{j \in \mathbb{N}} \subset \Omega$ and $(t_j)_{j \in \mathbb{N}} \subset (0, \infty)$ satisfying $t_j \rightarrow \infty$ as $j \rightarrow \infty$ such that

$$|u(x_j, t_j) - \bar{u}_0| \geq c_4, \quad \text{for all } j \in \mathbb{N}.$$

From Lemma 3.2, we know $u - \bar{u}_0$ is uniformly continuous in $\Omega \times (1, \infty)$. Then there exist $r > 0$ and $T_1 > 0$ such that for any $j \in \mathbb{N}$,

$$|u(x, t) - \bar{u}_0| \geq \frac{c_4}{2} \quad \text{for all } x \in B_r(x_j) \cap \Omega \text{ and } t \in (t_j, t_j + T_1). \quad (65)$$

Because of the smoothness of $\partial\Omega$, we can get a constant $c_5 > 0$ such that

$$|B_r(x_j) \cap \Omega| \geq c_5, \quad \text{for all } x_j \in \Omega. \quad (66)$$

Using (65) and (66), for all $j \in \mathbb{N}$, we have

$$\begin{aligned} \int_{t_j}^{t_j+T_1} \int_{\Omega} |u(x, t) - \bar{u}_0|^2 dx dt &\geq \int_{t_j}^{t_j+T_1} \int_{B_r(x_j) \cap \Omega} |u(x, t) - \bar{u}_0|^2 dx dt \\ &\geq \int_{t_j}^{t_j+T_1} |B_r(x_j) \cap \Omega| \cdot \left(\frac{c_4}{2}\right)^2 dt \\ &\geq \frac{c_4^2 c_5 T_1}{4}. \end{aligned} \quad (67)$$

However, by the fact $t_j \rightarrow \infty$ as $j \rightarrow \infty$, we have from (63) that

$$\int_{t_j}^{t_j+T_1} \int_{\Omega} (u(x, t) - \bar{u}_0)^2 dx dt \leq \int_{t_j}^\infty \int_{\Omega} (u(x, t) - \bar{u}_0)^2 dx dt \rightarrow 0, \quad \text{as } j \rightarrow \infty,$$

which contradicts (67). Hence (64) holds by the argument of contradiction. Thus the proof of Lemma 4.4 is completed. \square

Next, we will show the convergence of v and w by the comparison principle.

Lemma 4.5. *Let the conditions in Lemma 4.4 hold. Then it holds that*

$$\|v(\cdot, t) - \frac{\alpha}{\beta}\bar{u}_0\|_{L^\infty} \rightarrow 0, \text{ as } t \rightarrow \infty,$$

and

$$\|w(\cdot, t) - \frac{\gamma}{\delta}\bar{u}_0\|_{L^\infty} \rightarrow 0, \text{ as } t \rightarrow \infty.$$

Proof. Let $\phi(x, t) = v(x, t) - \frac{\alpha}{\beta}\bar{u}_0$. Then from the second equation of (5), one has

$$\begin{cases} \phi_t - D_1\Delta\phi + \beta\phi = \alpha(u - \bar{u}_0), & x \in \Omega, t > 0, \\ \frac{\partial\phi}{\partial\nu} = 0, & x \in \partial\Omega, t > 0, \\ \phi(x, 0) = \phi_0(x) = v_0(x) - \frac{\alpha}{\beta}\bar{u}_0, & x \in \Omega. \end{cases} \tag{68}$$

Let $\phi^*(t)$ be the solution of ODE problem

$$\begin{cases} \phi_t^*(t) + \beta\phi^*(t) = \alpha\|u - \bar{u}_0\|_{L^\infty}, t > 0, \\ \phi^*(0) = \|\phi_0\|_{L^\infty}. \end{cases} \tag{69}$$

The application of the comparison principle show that $\phi^*(t)$ is a super-solution of problem (68) and satisfies

$$\phi(x, t) \leq \phi^*(t) \text{ for all } x \in \Omega, t > 0.$$

Similarly, we can prove that $\phi(x, t) \geq -\phi^*(t)$ for all $x \in \Omega, t > 0$. Hence, one has

$$|\phi(x, t)| \leq \phi^*(t) \text{ for all } x \in \Omega, t > 0. \tag{70}$$

On the other hand, using the fact $\|u(\cdot, t) - \bar{u}_0\|_{L^\infty} \rightarrow 0$ as $t \rightarrow \infty$ and from (69) we have

$$\phi^*(t) \rightarrow 0 \text{ as } t \rightarrow \infty,$$

which combined with (70) gives

$$\|v(\cdot, t) - \frac{\alpha}{\beta}\bar{u}_0\|_{L^\infty} = \|\phi(\cdot, t)\|_{L^\infty} \leq \phi^*(t) \rightarrow 0 \text{ as } t \rightarrow \infty. \tag{71}$$

Similar arguments applied to the third equation of system (5) yield

$$\|w(\cdot, t) - \frac{\gamma}{\delta}\bar{u}_0\|_{L^\infty} \rightarrow 0, \text{ as } t \rightarrow \infty, \tag{72}$$

which completes the proof of Lemma 4.5. □

4.3. Decay rate. It is shown in section 4.2 that $(u, v, w) \rightarrow (\bar{u}_0, \frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\delta}\bar{u}_0)$ as $t \rightarrow \infty$ under the condition (6). Below, we will further show the convergence rate is exponential if $\frac{\xi\gamma}{\chi\alpha} > \max\left\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$.

Lemma 4.6. *Suppose that the conditions in Lemma 4.1 hold. If $\frac{\xi\gamma}{\chi\alpha} > \max\left\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$, then there exist two constants $C > 0$ and $\lambda > 0$ such that*

$$\|u(\cdot, t) - \bar{u}_0\|_{L^1} \leq Ce^{-\lambda t} \text{ for all } t > 0. \tag{73}$$

Proof. The nonnegativity of $E(u, v, w)$ and $F(u, v, w)$ has been proved in Lemma 3.3 under the condition (6). Next, we show that if $\frac{\xi\gamma}{\chi\alpha} > \max\left\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$, there exists a constant $\mu > 0$ which will be chosen later such that

$$E(u, v, w) \leq \mu F(u, v, w). \tag{74}$$

In fact, using the definition of $E(u, v, w)$ and $F(u, v, w)$ in (31) and (32), respectively, we derive that

$$\begin{aligned} \mathcal{D}(u, v, w) &= \mu F(u, v, w) - E(u, v, w) \\ &= \frac{\theta_1}{2\xi\chi} \left(\mu \int_{\Omega} \frac{|\nabla u|^2}{u} - \int_{\Omega} u \ln \frac{u}{\bar{u}} \right) + \mathcal{D}_1(u, v, w) + \mathcal{D}_2(u, v, w) \end{aligned} \tag{75}$$

where

$$\mathcal{D}_1(u, v, w) = \mu \left(\frac{\theta_2 D_1}{2\xi\alpha} \int_{\Omega} |\Delta v|^2 + \frac{\theta_2 D_2}{2\gamma\chi} \int_{\Omega} |\Delta w|^2 - (D_1 + D_2) \int_{\Omega} \Delta w \cdot \Delta v \right)$$

and

$$\begin{aligned} \mathcal{D}_2(u, v, w) &= \frac{\theta_2}{2\xi\alpha} \left(\beta\mu - \frac{1}{2} \right) \int_{\Omega} |\nabla v|^2 + \frac{\theta_2}{2\gamma\chi} \left(\delta\mu - \frac{1}{2} \right) \int_{\Omega} |\nabla w|^2 \\ &\quad + [1 - \mu(\beta + \delta)] \int_{\Omega} \nabla w \cdot \nabla v. \end{aligned}$$

To show the nonnegativity of $\mathcal{D}(u, v, w)$, we first show the nonnegativity of first term on the right hand of (75). From Lemma 2.1, we have

$$\int_{\Omega} u \ln \frac{u}{\bar{u}} \leq \frac{1}{\bar{u}} \|u - \bar{u}\|_{L^2}^2. \tag{76}$$

On the other hand, using (62) and the fact $\|u\|_{L^\infty} \leq c_1$, one derives

$$\frac{1}{\bar{u}} \|u - \bar{u}\|_{L^2}^2 \leq c_2 \|\nabla u\|_{L^2}^2 \leq c_2 \|u\|_{L^\infty} \int_{\Omega} \frac{|\nabla u|^2}{u} \leq c_1 c_2 \int_{\Omega} \frac{|\nabla u|^2}{u},$$

which combined with (76) gives

$$\int_{\Omega} u \ln \frac{u}{\bar{u}} \leq c_1 c_2 \int_{\Omega} \frac{|\nabla u|^2}{u} = \mu_1 \int_{\Omega} \frac{|\nabla u|^2}{u},$$

where $\mu_1 = c_1 c_2$. Hence, we can choose $\mu \geq \mu_1$ such that the first term on the right hand of (75) is nonnegative.

Next, we will show the nonnegativity of $\mathcal{D}_1(u, v, w)$. In fact, we can rewrite $\mathcal{D}_1(u, v, w)$ as

$$\mathcal{D}_1(u, v, w) = \mu \int_{\Omega} \Theta_2^T A_2 \Theta_2,$$

where A_2 and Θ_2 are defined in (42). The condition (6) gives $\frac{\xi\gamma}{\chi\alpha} \geq \max \left\{ \frac{D_1}{D_2}, \frac{D_2}{D_1} \right\}$. Then hence the matrix A_2 is nonnegative definite and hence $\mathcal{D}_1(u, v, w) \geq 0$ for any $\mu > 0$.

Similarly, to show the nonnegativity of $\mathcal{D}_2(u, v, w)$, we rewrite it as

$$\mathcal{D}_2(u, v, w) = \int_{\Omega} \Theta_1^T A_4 \Theta_1,$$

where

$$\Theta_1 = \begin{bmatrix} \nabla v \\ \nabla w \end{bmatrix} \quad \text{and} \quad A_4 = \begin{bmatrix} \frac{\theta_2}{2\xi\alpha}(\beta\mu - \frac{1}{2}) & \frac{1-\mu(\beta+\delta)}{2} \\ \frac{1-\mu(\beta+\delta)}{2} & \frac{\theta_2}{2\gamma\chi}(\delta\mu - \frac{1}{2}) \end{bmatrix}.$$

Using the matrix analysis, we know that A_4 is nonnegative definite if $\mu > \mu_2 := \max \left\{ \frac{1}{2\beta}, \frac{1}{2\delta} \right\}$ and

$$[4\theta_2^2\beta\delta - (\theta_2^2 - \theta_1^2)(\beta + \delta)^2]\mu^2 - 2(\beta + \delta)\theta_1^2\mu + \theta_1^2 \geq 0. \tag{77}$$

Since $\frac{\xi\gamma}{\chi\alpha} > \max\left\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$, one has

$$4\theta_2^2\beta\delta - (\theta_2^2 - \theta_1^2)(\beta + \delta)^2 = 4(\xi\gamma\beta - \chi\alpha\delta)(\xi\gamma\delta - \chi\alpha\beta) > 0.$$

Hence (77) holds if $\frac{\xi\gamma}{\chi\alpha} > \max\left\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\right\}$ and

$$\mu > \mu_3 := \max\left\{\frac{\theta_1}{2(\xi\gamma\delta - \chi\alpha\beta)}, \frac{\theta_1}{2(\xi\gamma\beta - \chi\alpha\delta)}\right\}.$$

Then if $\mu > \max\{\mu_2, \mu_3\}$, the function $\mathcal{D}_2(u, v, w)$ is nonnegative. Hence, choosing $\mu > \max\{\mu_1, \mu_2, \mu_3\}$, the function $\mathcal{D}(u, v, w)$ is nonnegative and (74) holds.

Substituting (74) into (33), we have

$$\frac{d}{dt}E(u, v, w) + \frac{1}{\mu}E(u, v, w) \leq 0,$$

which implies

$$E(u, v, w) \leq c_3 e^{-\frac{1}{\mu}t}. \tag{78}$$

On the other hand, from (40), we have

$$\frac{\theta_1}{2\xi\chi} \int_{\Omega} u \ln \frac{u}{\bar{u}} \leq E(u, v, w),$$

which along with (78) and Lemma 2.1 gives

$$\|u(\cdot, t) - \bar{u}_0\|_{L^1}^2 = \|u(\cdot, t) - \bar{u}\|_{L^1}^2 \leq \frac{4\xi\chi\bar{u}c_3}{\theta_1} e^{-\frac{1}{\mu}t}.$$

This yields (73) and concludes the proof. □

Next, we will derive the decay rate of solutions in L^∞ -norm based on the decay rate of $\|u(\cdot, t) - \bar{u}_0\|_{L^1}$.

Lemma 4.7. *Let (u, v, w) be the global classical solution of system (5). Suppose that there exist two positive constant C, λ such that*

$$\|u(\cdot, t) - \bar{u}_0\|_{L^1} \leq Ce^{-\lambda t}, \tag{79}$$

then the solution (u, v, w) will exponentially decay to $(\bar{u}_0, \frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\delta}\bar{u}_0)$ with L^∞ -norm as $t \rightarrow \infty$.

Proof. With (79) in hand, we can use the Moser-Alikakos iteration procedure as in [27] or the semigroup estimate method in [15] to obtain

$$\|u - \bar{u}_0\|_{L^\infty} \leq c_1 e^{-c_1 t}.$$

Then applying the comparison principle as in [27], one can show that there exists a constant $c_2 > 0$ such that

$$\|v - \frac{\alpha}{\beta}\bar{u}_0\|_{L^\infty} + \|w - \frac{\gamma}{\delta}\bar{u}_0\|_{L^\infty} \leq c_2 e^{-c_2 t}.$$

Then the proof of this lemma is completed. □

4.4. **Proof of Theorem 1.1.** Under the condition (6), we show the boundedness of solution for system (5) with $D_1 \neq D_2$ in Lemma 4.3, which implies there exists a constant $c_1 > 0$ such that $\|u(\cdot, t)\|_{L^\infty} \leq c_1$. Moreover, from Lemma 4.1, one has a constant $c_2 > 0$ such that

$$\int_0^t \int_\Omega \frac{|\nabla u|^2}{u} \leq c_2,$$

which together with the fact $\|u(\cdot, t)\|_{L^\infty} \leq c_1$ implies $\|u - \bar{u}_0\|_{L^\infty} \rightarrow 0$ as $t \rightarrow \infty$ as shown in Lemma 4.4. Then using the comparison principle for parabolic equations, from the second and third equations of system (5), we show that the solution (v, w) converges to $(\frac{\alpha}{\beta}\bar{u}_0, \frac{\gamma}{\beta}\bar{u}_0)$ as $t \rightarrow \infty$ in Lemma 4.5. Moreover, if $\xi\gamma > \chi\alpha \max\{\frac{\beta}{\delta}, \frac{\delta}{\beta}\}$, then using Lemma 4.6, we can obtain

$$\|u(\cdot, t) - \bar{u}_0\|_{L^1} \leq c_3 e^{-\lambda t},$$

which, along with Lemma 4.7, gives the exponential decay rate as shown in Theorem 1.1. Then Theorem 1.1 is proved.

Acknowledgments. The authors would like to thank the referee for his/her comments and suggestions on the improvement of the paper. The second author is particularly grateful to Professor Wei-Ming Ni for his very kind support and constant encouragement in his career pursue.

REFERENCES

- [1] N. Bellomo, A. Bellouquid, Y. Tao and M. Winkler, [Towards a mathematical theory of Keller-Segel models of pattern formation in biological tissues](#), *Math. Models Methods Appl. Sci.*, **25** (2015), 1663–1763.
- [2] J. P. Bourguignon and H. Brezis, [Remarks on Euler Equation](#), *J. Functional Analysis*, **15** (1974), 341–363.
- [3] J. A. Carrillo, A. Juöngle, P. A. Markowich, G. Toscani and A. Unterreiter, [Entropy dissipation methods for degenerate parabolic problems and generalized Sobolev inequalities](#), *Monatsh. Math.*, **133** (2001), 1–82.
- [4] T. Cieślak, Ph. Laurencot and C. Morales-Rodrigo, [Global existence and convergence to steady states in a chemorepulsion system](#), In *Parabolic and Navier-Stokes equations*, *Banach Center Publ., Polish Acad. Sci. Inst. Math.*, **81** (2008), 105–117.
- [5] E. Espejo and T. Suzuki, [Global existence and blow-up for a system describing the aggregation of microglia](#), *Appl. Math. Lett.*, **35** (2014), 29–34.
- [6] D. Horstmann, [From 1970 until present: the Keller-Segel model in chemotaxis and its consequences](#), *I. Jahresber. Deutsch. Math. Verien.*, **105** (2003), 103–165.
- [7] D. Horstmann and G. Wang, [Blow-up in a chemotaxis model without symmetry assumptions](#), *European J. Appl. Math.*, **12** (2001), 159–177.
- [8] H. Y. Jin, [Boundedness of the attraction-repulsion Keller-Segel system](#), *J. Math. Anal. Appl.*, **422** (2015), 1463–1478.
- [9] H. Y. Jin and Z. Liu, [Large time behavior of the full attraction-repulsion Keller-Segel system in the whole space](#), *Appl. Math. Lett.*, **47** (2015), 13–20.
- [10] H. Y. Jin and Z. A. Wang, [Asymptotic dynamics of the one-dimensional attraction-repulsion Keller-Segel model](#), *Math. Methods Appl. Sci.*, **38** (2015), 444–457.
- [11] H. Y. Jin and Z. A. Wang, [Boundedness, blowup and critical mass phenomenon in competing chemotaxis](#), *J. Differential Equations*, **260** (2016), 162–196.
- [12] R. Kowalczyk and Z. Szymańska, [On the global existence of solutions to an aggregation model](#), *J. Math. Anal. Appl.*, **343** (2008), 379–398.
- [13] O. Ladyzhenskaya, V. Solonnikov and N. Uralceva, *Linear and Quasilinear Equations of Parabolic Type*, (Russian) Translated from the Russian by S. Smith. Translations of Mathematical Monographs, American Mathematical Society, Providence, R.I., 1968.
- [14] Y. Li and Y. X. Li, [Blow-up of nonradial solutions to attraction-repulsion chemotaxis system in two dimensions](#), *Nonlinear Anal. Real World Appl.*, **30** (2016), 170–183.

- [15] K. Lin and C. Mu, [Global existence and convergence to steady states for an attraction-repulsion chemotaxis system](#), *Nonlinear Anal. Real World Appl.*, **31** (2016), 630–642.
- [16] K. Lin, C. Mu and L. Wang, [Large-time behavior of an attraction-repulsion chemotaxis system](#), *J. Math. Anal. Appl.*, **426** (2015), 105–124.
- [17] K. Lin, C. Mu and D. Zhou, [Stabilization in a higher-dimensional attraction-repulsion chemotaxis system if repulsion dominates over attraction](#), *Math. Models Methods Appl. Sci.*, **28** (2018), 1105–1134.
- [18] D. Liu and Y. S. Tao, [Global boundedness in a fully parabolic attraction-repulsion chemotaxis model](#), *Math. Methods Appl. Sci.*, **38** (2015), 2537–2546.
- [19] P. Liu, J. P. Shi and Z. A. Wang, [Pattern formation of the attraction-repulsion Keller-Segel system](#), *Discrete Contin. Dyn. Syst. Ser. B*, **18** (2013), 2597–2625.
- [20] J. Liu and Z. A. Wang, [Classical solutions and steady states of an attraction-repulsion chemotaxis model in one dimension](#), *J. Biol. Dyn.*, **6** (2012), 31–41.
- [21] M. Luca, A. Chavez-Ross, L. Edelstein-Keshet and A. Mogilner, [Chemotactic signalling, Microglia, and Alzheimer’s disease senile plaques: Is there a connection?](#) *Bull. Math. Biol.*, **65** (2003), 693–730.
- [22] T. Nagai, T. Senba and K. Yoshida, [Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis](#), *Funkcial. Ekvac.*, **40** (1997), 411–433.
- [23] K. J. Painter and T. Hillen, [Volume-filling quorum-sensing in models for chemosensitive movement](#), *Can. Appl. Math. Q.*, **10** (2002), 501–543.
- [24] M. M. Porzio and V. Vespri, [Hölder estimates for local solutions of some doubly nonlinear degenerate parabolic equations](#), *J. Differential Equations*, **103** (1993), 146–178.
- [25] R. Shi and W. Wang, [Well-posedness for a model derived from an attraction-repulsion chemotaxis system](#), *J. Math. Anal. Appl.*, **423** (2015), 497–520.
- [26] Y. S. Tao, [Global dynamics in a higher-dimensional repulsion chemotaxis model with nonlinear sensitivity](#), *Discrete Contin. Dyn. Syst. Ser. B*, **18** (2013), 2705–2722.
- [27] Y. S. Tao and Z. A. Wang, [Competing effects of attraction vs. repulsion in chemotaxis](#), *Math. Models Methods Appl. Sci.*, **23** (2013), 1–36.
- [28] Y. S. Tao and M. Winkler, [Large time behavior in a multidimensional chemotaxis-haptotaxis model with slow signal diffusion](#), *SIAM J. Math. Anal.*, **47** (2015), 4229–4250.
- [29] M. Winkler, [Aggregation vs. global diffusive behavior in the higher-dimensional Keller-Segel model](#), *J. Differential Equations*, **248** (2010), 2889–2905.
- [30] M. Winkler, [Finite-time blow-up in the higher-dimensional parabolic-parabolic Keller-Segel system](#), *J. Math. Pures Appl.*, **100** (2013), 748–767.

Received May 2018; revised March 2019.

E-mail address: mahyjin@scut.edu.cn

E-mail address: mawza@polyu.edu.hk